

RESEARCH ARTICLE

View Article Online

View Journal | View Issue

Cite this: *Org. Chem. Front.*, 2023, **10**, 3000

Rhodium-catalyzed regioselective C–H activation/Lossen rearrangement/annulation for the green synthesis of trisubstituted 2-pyridones†

Yidi Li,^{‡a,e} Huiying Xu,^{‡c} Lin Huang,^{‡b,d} Zhi Zhou,^{‡c} Zhenhao Tang,^b Haifang Meng,^b Wei Zhang,^b Wei Yi,^{‡c} and Xiaowei Wu^{‡*a,b,d,e}

Multisubstituted 2-pyridones are prevalent in pharmaceuticals and bioactive molecules. We herein report an efficient and regioselective approach for the synthesis of trisubstituted 2-pyridone derivatives by a rhodium-catalyzed C–H activation/Lossen rearrangement/cyclization cascade reaction between acrylamides and propargyl alcohols. The desirable features of this protocol include a reusable catalytic system, high regioselectivity, uncommon Lossen rearrangement, good functional group tolerance, operation at room temperature, simple purification by filtration in most cases, and scale-up synthesis with as low as 1 mol% catalyst loading. Additionally, deuterium labeling and KIE assays were performed to investigate the reaction mechanism. The vital effect of the hydroxyl group on propargyl alcohols in determining the regioselectivity was demonstrated by control experiments and DFT calculations. In addition, Mulliken atomic charge analysis of the key intermediates was also carried out to probe the origin of the observed preference for the Lossen rearrangement.

Received 31st March 2023,

Accepted 10th May 2023

DOI: 10.1039/d3qo00469d

rsc.li/frontiers-organic

Introduction

2-Pyridone is a prevalent scaffold in organic compounds and bioactive molecules.¹ As a class of six-membered aza-heterocycles, the 2-pyridone ring possesses a nitrogen heteroatom and a carbonyl group which can act as a hydrogen bond donor/acceptor in medicinal chemistry. In this regard, 2-pyridone usually serves as a bioisostere for pyridine, amide, and N/O-containing heterocycles. Owing to their unique structures, 2-pyridones have been utilized as effective ligands for C–H

functionalization as well as kinase hinge binding motifs.^{2,3} In addition, the use of a 2-pyridone moiety as bioisosteres commonly has evident influences on the solubility, lipophilicity, and metabolic stability of bioactive compounds. As shown in Fig. 1, 2-pyridone derivatives exhibit a variety of pharmacological activities, such as antifungal, antiepileptic, anticancer, anti-fibrotic, cardiostonic, and anti-HIV activities.^{3b,4} Considering the prevalence and importance of the 2-pyridone scaffold, it is appealing to develop efficient methods for the synthesis of 2-pyridones.

Over the past decades, as a direct and step-economical strategy to construct heterocycles, C–H activation reactions by tran-

^aDrug Discovery and Development Center, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai 201203, China.

E-mail: wuxiaowei@simm.ac.cn

^bZhongshan Institute for Drug Discovery, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Zhongshan 528400, China

^cGuangzhou Municipal and Guangdong Provincial Key Laboratory of Protein Modification and Degradation & Molecular Target and Clinical Pharmacology, State Key Laboratory of Respiratory Disease, School of Pharmaceutical Sciences & the Fifth Affiliated Hospital, Guangzhou Medical University, Guangzhou, Guangdong 511436, China. E-mail: zhouzhi@gzhmu.edu.cn, yiwei@gzhmu.edu.cn

^dSchool of Pharmaceutical Sciences, Southern Medical University, Guangzhou 510515, China

^eUniversity of Chinese Academy of Sciences, No. 19A Yuquan Road, Beijing 100049, China

†Electronic supplementary information (ESI) available. CCDC 2213771 (3e) and 2234440 (3w). For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d3qo00469d>

‡These authors contributed equally.




Fig. 1 Selected bioactive molecules containing the 2-pyridone motif.



sition-metal catalysis have received much attention.⁵ In this respect, transition-metal catalyzed C–H activation/annulation of benzamides or acrylamides with alkynes has become an efficient method for the preparation of isoquinolones or pyridones (Scheme 1a).^{6–8} In recent years, propargyl alcohols have been frequently utilized in reactions because of their versatile reactivities.⁹ Additionally, regioselectivity and chemoselectivity are usually controlled by both the hydroxyl group on propargyl alcohols and the directing groups (DGs) in these C–H activation reactions. For example, a ruthenium-catalyzed C–H activation/[4 + 1] annulation of benzamides and propargyl alcohols was pioneered by Liu and coworkers, in which only one carbon of propargyl alcohols was involved in the cyclization (Scheme 1b).^{9b} Very recently, by employing the pivaloyl group to replace the ethyl group of directing groups, we reported a green and efficient rhodium-catalyzed C–H activation/annulation of *N*-(pivaloyloxy)benzamides and propargyl alcohols for the synthesis of isoquinolones (Scheme 1c).¹⁰


Given the prevalence and importance of 2-pyridones, we wish to develop an efficient and novel method to prepare novel

multi-substituted 2-pyridone derivatives. Owing to the presence of a hydroxyl group on propargyl alcohols, unique chemoselectivity and regioselectivity are usually achieved. Nevertheless, propargyl alcohols are frequently coupled with aromatic substrates in C–H activation reactions. The C–H activation/annulation of alkenyl substrates and propargyl alcohols for the synthesis of multi-substituted 2-pyridones remains elusive. Inspired by previous works, we speculated whether propargyl alcohols can facilitate some novel transformations when reacted with alkenyl substrates. Thus, we herein report a green and efficient rhodium-catalyzed C–H activation/Lossen rearrangement of acrylamides and propargyl alcohols for the synthesis of novel 2-pyridone derivatives at ambient temperature (Scheme 1d).


Results and discussion

Optimization studies towards the synthesis of 4-cyclopropyl-3-(hydroxy(phenyl)methyl)-6-phenylpyridin-2(1*H*)-one **3a** are shown in Table 1. In order to develop a green chemical reaction, we first focused on the use of ethyl acetate as the solvent. Then, various transition-metal catalysts were screened in the presence of KOAc at room temperature. No reaction of 2-phenyl-*N*-(pivaloyloxy)acrylamide **1a** with 3-cyclopropyl-1-phenylprop-2-yn-1-ol **2a** was promoted by 4 mol% of MnBr(CO)₅, [Cp*IrCl₂]₂, Cp*Co(CO)₂, and [RuCl₂(*p*-cym)₂], while the desired product **3a** was obtained in 64% isolated yield when [Cp*RhCl₂]₂ was used (Table 1, entries 1–5). To our delight,

Previous work:



This work:




Scheme 1 C–H activation reactions of benzamides or acrylamides with alkynes.

Table 1 Optimization of reaction conditions^a

Entry	Catalyst	Additive	Solvent	Yield ^{b,c} (%)
1	MnBr(CO) ₅	KOAc	EA	0
2	[Cp*IrCl ₂] ₂	KOAc	EA	0
3	Cp*Co(CO) ₂	KOAc	EA	0
4	[Ru(<i>p</i> -cym)Cl ₂] ₂	KOAc	EA	0
5	[Cp*RhCl ₂] ₂	KOAc	EA	64 (59)
6	[Cp*RhCl ₂] ₂	KOAc	EtOH	22 (16)
7	[Cp*RhCl ₂] ₂	KOAc	TFE	40
8	[Cp*RhCl ₂] ₂	KOAc	DCM	44
9	[Cp*RhCl ₂] ₂	KOAc	Acetone	44 (36)
10	[Cp*RhCl ₂] ₂	NaOAc	EA	69 (67)
11	[Cp*RhCl ₂] ₂	Na ₂ CO ₃	EA	58 (47)
12	[Cp*RhCl ₂] ₂	NaHCO ₃	EA	69 (58)
13	[Cp*RhCl ₂] ₂	CsOAc	EA	44 (37)
14 ^d	[Cp*RhCl ₂] ₂	KOAc	EA	57 (35)
15 ^d	[Cp*RhCl ₂] ₂	NaOAc	EA	59 (51)
16 ^e	[Cp*RhCl ₂] ₂	NaOAc	EA	73 (70)

^a Reaction conditions: **1a** (0.2 mmol), **2a** (0.3 mmol), catalyst (4 mol%), additive (0.2 mmol), under air, solvent (1.0 mL), room temperature, 12 h. ^b Total isolated yield. ^c Isolated yield by filtration is shown in parenthesis. ^d [Cp*RhCl₂]₂ (2.5 mol%). ^e Reaction time was 6.0 h.





Scheme 2 Substrate scope. Reaction conditions: **1** (0.25 mmol), **2** (0.375 mmol), catalyst (4 mmol%), NaOAc (1.0 equiv.), under air, ethyl acetate (1.0 mL), rt, 6–12 h. Isolated yields by simple filtration are shown in parentheses.




most parts of the product could be isolated by simple filtration, which avoided the use of large amounts of organic solvents for purification. Different solvents were also investigated. Unfortunately, none was superior to our initial choice, EA (entries 6–9). Additive screening revealed that NaOAc and NaHCO₃ were the best bases in terms of total isolated yields (entries 10–13). However, the product was obtained in 67% filtration yield in the presence of NaOAc while 58% filtration yield was obtained when NaHCO₃ was added. The yield was decreased when KOAc and 2.5 mol% [Cp*RhCl₂]₂ were used, when compared to entry 5 (entry 14). Similarly, when the base was changed to NaOAc, the total yield also declined to 59% by using 2.5 mol% [Cp*RhCl₂]₂ (entry 15). The above-mentioned reactions were conducted for 12 hours. Subsequently, the reaction time was decreased to 6 hours. Surprisingly, the yield of **3a** improved slightly compared with that obtained in 12 hours (entry 16).

The optimal reaction conditions were then applied to various substrates, as summarized in Scheme 2. Acrylamides reacted well with different propargyl alcohols. When the cyclopropyl group was placed at R², both electron-donating (Me and OMe) and electron-withdrawing (F, Cl, Br, I, CN, NO₂ and CF₃) substituents were compatible, with the isolated yields ranging from 39% to 75% (**3b–3k**). The structure of **3c** was confirmed by X-ray crystallographic analysis (CCDC number 2213771 for **3c**, see the ESI† for more details). In addition, alkynols containing cyclohexyl, *N*-Boc protected piperidyl, and a series of heteroaromatic rings such as thienyl, naphthyl and furyl rings also provided the corresponding products smoothly (**3m–3q**). It is also noteworthy that the corresponding products showed


excellent results when using tertiary alcohols, especially toward those naphthenic groups whose yields even surpassed that of the template reaction (**3r–3u**). This suggested that the reaction could overcome the issue of steric effect. We next investigated phenyl, methyl and *n*-butyl at R² and the reactions also took place smoothly with yields of 42%, 78% and 31%, respectively (**3v–3x**). Besides, the structure of **3w** was also con-

a) Deuterium incorporation experiments




b) Kinetic isotope effect study

parallel reactions




Scheme 3 Deuteration and KIE experiments.


a) Scale-up synthesis




b) Control experiments



c) Recycling experiments



Flow diagram of recycling experiments



Scheme 4 Scale-up preparation, control experiments, and recycling experiments.




firmed by X-ray crystallographic analysis (CCDC number 2234440 for **3w**, see the ESI† for more details). Subsequently, the introduction of other groups at R¹ was examined. Obviously, several substituents with different electron perturbations were tolerated and the yields were moderate (**3y–3zb**).


In addition, it was difficult to obtain a few products (**3b**, **3j**, **3n**, **3w**, **3x**, and **3y**) just by simple filtration probably due to their good solubility in ethyl acetate.

To probe the reaction mechanism, deuterium labeling and KIE assays were carried out (Scheme 3). In the presence of **1a**,

a) computed Gibbs free energy changes of the reaction path



b) IGMH analysis of TS-1 and TS-1i ($\delta g^{\text{inter}} = 0.008$)



Scheme 5 DFT calculations.




NaOAc, $[\text{Cp}^*\text{RhCl}_2]_2$, and 1.0 mL methanol- d^4 , about 10% deuterium incorporation occurred at the olefinic bond of acrylamide **1a**. The result indicated that the cleavage of the $\text{C}(\text{sp}^2)\text{--H}$ bond could be reversible. Approximately 100% deuteration occurred at the hydroxyl group of **3a** when **2a** was added, and no obvious deuteration was observed at other positions. In addition, two parallel reactions were performed to give a KIE value of 1.12 (Scheme 3b), suggesting that the step of C–H activation was not the rate-determining step.

Besides, to explore the synthetic utility of the methodology, several scale-up experiments were performed with lower catalyst loading. As shown in Scheme 4a, product **3a** was obtained in an ideal isolated yield (57%) with 2.5 mol% rhodium catalyst. Then, when 1.0 mol% rhodium catalyst was used, the corresponding product **3a** was afforded in 60% yield, respectively. Inspired by these results, we further explored another substrate **2p**, and the reaction afforded product **3p** in 79% yield. To explore the effect of the hydroxyl group on propargyl alcohols in determining the regioselectivity, two control experiments were conducted (Scheme 4b). The hydroxyl group of **2a** was removed to prepare **2a-1** and **2a-2**. The annulation of **1a** and **2a-1** gave two rearrangement products **4a** and **4a'** at the same time with comparable yields. Similarly, the reaction between **1a** and 3-cyclopropyl-1-phenylprop-2-yn-1-one **2a-2** also afforded two regioisomers **5a** and **5a'** in low yields. The structures of these compounds were determined by using NOESY spectra. A relatively higher regioselectivity could be detected for **2a-2** in comparison with **2a-1**; we proposed that the observed regioselectivity might be due to the electron-withdrawing character of the carbonyl moiety. These results indicated that the hydroxyl group plays a vital role in controlling the regioselectivity. Significantly, the recycling experiments of the catalytic system were carried out four times with desirable isolated yields under the standard conditions (Scheme 4c).

Having established the $\text{Rh}(\text{III})$ -catalyzed sequential C–H activation/Lossen rearrangement/[4 + 2] annulation cascade of acrylamides with propargyl alcohols, we were next interested to clarify the deep origin of the unconventional regio-/chemoselectivity by detailed DFT calculations. As shown in Scheme 5a, the five-membered rhodacycle **INT-0** was rationally selected as the starting point, which coordinated with propargyl alcohol **2a** followed by a regioselective migratory alkyne insertion. The calculated results revealed that an additional hydrogen bond affinity between the hydroxyl group and the DG was involved in **TS-1** ($\Delta G^\ddagger = 3.6 \text{ kcal mol}^{-1}$) to give **INT-2** with a free energy of $-25.2 \text{ kcal mol}^{-1}$, while a relatively higher energy barrier was involved in the converse regioselectivity via **TS-1i** ($\Delta G^\ddagger = 4.8 \text{ kcal mol}^{-1}$). Further IGMH analysis showed obvious hydrogen bond and van der Waals force interactions in **TS-1**, while only van der Waals force interaction was observed in **TS-1i** (Scheme 5b). Subsequent coordination change via **TS-2** ($\Delta G^\ddagger = -14.5 \text{ kcal mol}^{-1}$) afforded **INT-3**, from which different reaction paths were calculated. The concerted Lossen rearrangement/N–O bond cleavage from **INT-3** occurred via **TS-3** with an energy barrier of $7.0 \text{ kcal mol}^{-1}$ (from **INT-3** to **TS-3**) to furnish the isocyanate intermediate **INT-4** with an

obvious exothermic process. Alternatively, the classic C–N bond reductive elimination via **TS-3ii** ($\Delta G^\ddagger = -0.2 \text{ kcal mol}^{-1}$) involved a higher energy barrier of $23.0 \text{ kcal mol}^{-1}$ (from **INT-3** to **TS-3ii**), which was in line with the experimental result that no 3-phenylpyridin-2(1*H*)-one framework was formed. Moreover, the $\text{Rh}(\text{III})\text{--Rh}(\text{V})\text{--Rh}(\text{III})$ reaction pathway involving an oxidative addition process from **INT-3** was also ruled out owing to the relatively higher free energies of **TS-4i/TS-4ii**. Taken together, the DFT calculations illustrated a hydrogen bond assisted regioselective alkyne insertion/Lossen rearrangement/intramolecular [4 + 2] cyclization reaction pathway for the developed protocol. In addition, further Mulliken atomic charge analysis of the key intermediates was also carried out to probe the origin of the observed preference for Lossen rearrangement rather than other reaction manifolds (see the ESI† for details). The results suggested that the C2 atom in **INT-3** occupied a relatively more positive charge in comparison with the similar benzamide substrate (0.0346 vs. 0.0216). Thus, it can be inferred that there was an inclination to undergo a nucleophilic attack from the N atom, ultimately leading to the Lossen rearrangement process.

On the basis of the above mechanistic studies and literature precedents, we proposed a plausible catalytic cycle for the developed transformation (Scheme 6). Initially, the active $\text{Cp}^*\text{Rh}(\text{OAc})_2$ species was generated by anion ligand exchange in the presence of NaOAc, which coordinated with the acrylamide substrate and participated in the alkenylic C–H bond activation to afford intermediate **A**. Subsequent regioselective alkyne insertion led to the formation of intermediate **B**, in which the hydrogen bond affinity played a crucial role in determining the regioselectivity. Furthermore, the Lossen rearrangement process occurred smoothly to give the isocyanate intermediate **C**, which underwent an intramolecular [4 + 2] cyclization to deliver the 2-pyridone skeleton **D**. Finally, the protonolysis of **D** with the assistance of HOAc released the



Scheme 6 Proposed catalytic cycle.



desired product **3a** along with the regeneration of the Rh(III) catalyst.

Conclusions

In summary, a green and efficient rhodium(III)-catalyzed C–H activation/Lossen rearrangement of acrylamides and propargyl alcohols for the synthesis of novel 2-pyridone derivatives at ambient temperature was developed. This protocol features a reusable catalytic system, high regioselectivity, uncommon Lossen rearrangement, good functional group tolerance, metal oxidant-free process, operation at room temperature, simple purification by filtration in most cases, scale-up synthesis, and air compatibility. Additionally, deuterium labeling and KIE assays were performed to investigate the reaction mechanism. The vital effect of the hydroxyl group on propargyl alcohols in determining the regioselectivity was also demonstrated by control experiments and DFT calculations. In addition, Mulliken atomic charge analysis of the key intermediates was also carried out to probe the origin of the observed preference for Lossen rearrangement.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Financial support from the Shanghai Pujiang Program (21PJ1415800), Basic and Applied Basic Research Foundation of Guangdong Province (2021A1515110468), Natural Science Foundation of Guangdong Province (2019A151010935), High-level New R&D Institute (2019B090904008), High-level Innovative Research Institute (2021B0909050003), and NSFC (21877020, 22007020) is gratefully acknowledged.

Notes and references

- (a) M. Hagimori, T. Temma, N. Mizuyama, T. Uto, Y. Yamaguchi, Y. Tominaga, T. Mukai and H. Saji, A high-affinity fluorescent Zn^{2+} sensor improved by the suppression of pyridine-pyridone tautomerism and its application in living cells, *Sens. Actuators, B*, 2015, **213**, 45–52; (b) G. Li, H. M. Duong, Z. Zhang, J. Xiao, L. Liu, Y. Zhao, H. Zhang, F. Huo, S. Li, J. Ma, F. Wudl and Q. Zhang, Approaching a stable, green twisted heteroacene through “clean reaction” strategy, *Chem. Commun.*, 2012, **48**, 5974–5976; (c) S. Hibi, K. Ueno, S. Nagato, K. Kawano, K. Ito, Y. Norimine, O. Takenaka, T. Hanada and M. Yonaga, Discovery of 2-(2-oxo-1-phenyl-5-pyridin-2-yl-1,2-dihydropyridin-3-yl)benzonitrile (perampanel): a novel, noncompetitive α -amino-3-hydroxy-5-methyl-4-isoxazolepropanoic acid (AMPA) receptor antagonist, *J. Med. Chem.*, 2012, **55**, 10584–10600;
- (d) M. Bialer and H. S. White, Key factors in the discovery and development of new antiepileptic drugs, *Nat. Rev. Drug Discovery*, 2010, **9**, 68–82; (e) K. Ishiuchi, T. Kubota, H. Ishiyama, S. Hayashi, T. Shibata and J. Kobayashi, Lycnadins C and F, new *Lycopodium* alkaloids from *Lycopodium complanatum*, *Tetrahedron Lett.*, 2011, **52**, 289–292; (f) M. Torres, S. Gil and M. Parra, New Synthetic Methods to 2-Pyridone Rings, *Curr. Org. Chem.*, 2005, **9**, 1757–1779; (g) I. M. Lagoja, Pyrimidine as Constituent of Natural Biologically Active Compounds, *Chem. Biodiversity*, 2005, **2**, 1–50; (h) P. Hajek, H. McRobbie and K. Myers, *Thorax*, 2013, **68**, 1037–1042.
- (a) P. Wang, P. Verma, G. Xia, J. Shi, J. Qiao, S. Tao, P. T. W. Cheng, M. A. Poss, M. E. Farmer, K.-S. Yeung and J.-Q. Yu, Ligand-accelerated non-directed C–H functionalization of arenes, *Nature*, 2017, **551**, 489–493; (b) Y.-Q. Chen, Z. Wang, Y. Wu, S. R. Wisniewski, J. Qiao, W. R. Ewing, M. D. Eastgate and J.-Q. Yu, Overcoming the Limitations of γ - and δ -C–H Arylation of Amines through Ligand Development, *J. Am. Chem. Soc.*, 2018, **140**, 17884–17894; (c) L.-Y. Liu, K.-S. Yeung and J.-Q. Yu, Ligand-Promoted Non-Directed C–H Cyanation of Arenes, *Chem. – Eur. J.*, 2019, **25**, 2199–2202; (d) X.-Y. Chen, Y. Wu, J. Zhou, P. Wang and J.-Q. Yu, Synthesis of β -Arylethenesulfonyl Fluoride via Pd-Catalyzed Nondirected C–H Alkenylation, *Org. Lett.*, 2019, **21**, 1426–1429; (e) P. Wang, G.-C. Li, P. Jain, M. E. Farmer, J. He, P.-X. Shen and J.-Q. Yu, Ligand-Promoted meta-C–H Amination and Alkynylation, *J. Am. Chem. Soc.*, 2016, **138**, 14092–14099; (f) P. Wang, M. E. Farmer, X. Huo, P. Jain, P. Shen, M. Ishoe, J. E. Bradner, S. R. Wisniewski, M. D. Eastgate and J.-Q. Yu, Ligand-Promoted Meta-C–H Arylation of Anilines, Phenols, and Heterocycles, *J. Am. Chem. Soc.*, 2016, **138**, 9269–9276; (g) R.-Y. Zhu, Z.-Q. Li, H. S. Park, C. H. Senanayake and J.-Q. Yu, Ligand-Enabled γ -C(sp³)–H Activation of Ketones, *J. Am. Chem. Soc.*, 2018, **140**, 3564–3568.
- (a) T. Siu, E. S. Kozina, J. Jung, C. Rosenstein, A. Mathur, M. D. Altman, G. Chan, L. Xu, E. Bachman, J. R. Mo, M. Bouthillette, T. Rush, C. J. Dinsmore, C. G. Marshall and J. R. Young, The discovery of tricyclic pyridone JAK2 inhibitors. Part 1: Hit to lead, *Bioorg. Med. Chem. Lett.*, 2010, **20**, 7421–7425; (b) V. Simov, S. V. Deshmukh, C. J. Dinsmore, F. Elwood, R. B. Fernandez, Y. Garcia, C. Gibeau, H. Gunaydin, J. Jung, J. D. Katz, B. Kraybill, B. Lapointe, S. B. Patel, T. Siu, H. Su and J. R. Young, Structure-based design and development of (benz)imidazole pyridones as JAK1-selective kinase inhibitors, *Bioorg. Med. Chem. Lett.*, 2016, **26**, 1803–1808; (c) J. Bach, P. Eastwood, J. Gonzalez, E. Gómez, J. A. Alonso, S. Fonquerna, E. Lozoya, A. Orellana, M. Maldonado, E. Calaf, J. Albertí, J. Pérez, A. Andrés, N. Prats, C. Carreño, E. Calama, J. D. Alba, M. Calbet, M. Miralpeix and I. Ramis, Identification of 2-Imidazopyridine and 2-Aminopyridone Purinones as Potent Pan-Janus Kinase (JAK) Inhibitors for the Inhaled Treatment of Respiratory Diseases, *J. Med. Chem.*, 2019, **62**, 9045–9060; (d) H. S. Eidam, J. Russell,



- K. Raha, M. DeMartino, D. Qin, H. A. Guan, Z. Zhang, G. Zhen, H. Yu, C. Wu, Y. Pan, G. Joberty, N. Zinn, S. Laquerre, S. Robinson, A. White, A. Giddings, E. Mohammadi, B. G.-V. Meerveld, A. Oliff, S. Kumar and M. Cheung, Discovery of a First-in-Class Gut-Restricted RET Kinase Inhibitor as a Clinical Candidate for the Treatment of IBS, *ACS Med. Chem. Lett.*, 2018, **9**, 623–628; (e) A. J. Massey, S. Stokes, H. Browne, N. Foloppe, A. Fiumana, S. Scrace, M. Fallowfield, S. Bedford, P. Webb, L. Baker, M. Christie, M. J. Drysdale and M. Wood, Identification of novel, in vivo active Chk1 inhibitors utilizing structure guided drug design, *Oncotarget*, 2015, **6**, 35797–35812; (f) M. C. Bryan, D. J. Burdick, B. K. Chan, Y. Chen, S. Clausen, J. Dotson, C. Eigenbrot, R. Elliott, E. J. Hanan, R. Heald, P. Jackson, H. La, M. Lainchbury, S. Malek, S. E. Mann, H. E. Purkey, G. Schaefer, S. Schmidt, E. Seward, S. Sideris, S. Wang, I. Yen, C. Yu and T. P. Heffron, Pyridones as Highly Selective, Noncovalent Inhibitors of T790M Double Mutants of EGFR, *ACS Med. Chem. Lett.*, 2016, **7**, 100–104.
- 4 (a) M. Niewerth, D. Kunze, M. Seibold, M. Schaller, H. C. Korting and B. Hube, Ciclopirox olamine treatment affects the expression pattern of *Candida albicans* genes encoding virulence factors, iron metabolism proteins, and drug resistance factors, *Antimicrob. Agents Chemother.*, 2003, **47**, 1805–1817; (b) T. Chen, S.-H. Dai, Z.-Q. Jiang, P. Luo, X.-F. Jiang, Z. Fei, S.-B. Gui and Y.-L. Qi, The AMPAR Antagonist Perampanel Attenuates Traumatic Brain Injury Through Anti-Oxidative and Anti-Inflammatory Activity, *Cell. Mol. Neurobiol.*, 2017, **37**, 43–52; (c) H. Nakazato, H. Oku, S. Yamane, Y. Tsuruta and R. Suzuki, A novel anti-fibrotic agent pirfenidone suppresses tumor necrosis factor- α at the translational level, *Eur. J. Pharmacol.*, 2002, **20**, 177–185; (d) Z. A. Kalme, R. A. Zhalubovskis, A. Shmidlers, J. Celmins and G. Duburs, Synthesis of 6-Bromomethyl-Substituted D Derivatives of Pyridin-2(1H)-Ones and Their Reaction with Nucleophiles, *Chem. Heterocycl. Compd.*, 2004, **40**, 862–868; (e) M.-T. Lai, M. Feng, J.-P. Falgoutyret, P. Tawa, M. Witmer, D. DiStefano, Y. Li, J. Burch, N. Sachs, M. Lu, E. Cauchon, L.-C. Campeau, J. Grobler, Y. Yan, Y. Ducharme, B. Côté, E. Asante-Appiah, D. J. Hazuda and M. D. Miller, In vitro characterization of MK-1439, a novel HIV-1 nonnucleoside reverse transcriptase inhibitor, *Antimicrob. Agents Chemother.*, 2014, **58**, 1652–1663.
- 5 Selected recent reviews: (a) P. Gandeepan, T. Müller, D. Zell, G. Cera, S. Warratz and L. Ackermann, 3d Transition Metals for C–H Activation, *Chem. Rev.*, 2019, **119**, 2192–2452; (b) D. J. Abrams, P. A. Provencher and E. J. Sorensen, Recent applications of C–H functionalization in complex natural product synthesis, *Chem. Soc. Rev.*, 2018, **47**, 8925–8967; (c) C. Sambigao, D. Schönbauer, R. Blicke, T. D. Huy, G. Pototschnig, P. Schaaf, T. Wiesinger, M. F. Zia, J. W. Delord, T. Besset, B. U. W. Maes and M. Schnürch, A comprehensive overview of directing groups applied in metal-catalysed C–H functionalisation chemistry, *Chem. Soc. Rev.*, 2018, **47**, 6603–6743; (d) Y. Yang, J. Lan and J. You, Oxidative C–H/C–H Coupling Reactions between Two (Hetero)arenes, *Chem. Rev.*, 2017, **117**, 8787–8863; (e) T. Gensch, M. N. Hopkinson, F. Glorius and J. W. Delord, Mild metal-catalyzed C–H activation: examples and concepts, *Chem. Soc. Rev.*, 2016, **45**, 2900–2936; (f) T. Yamada, Y. Shibata, S. Kawauchi, S. Yoshizaki and K. Tanaka, Formal Lossen Rearrangement/[3+2] Annulation Cascade Catalyzed by a Modified Cyclopentadienyl Rh^{III} Complex, *Chem. – Eur. J.*, 2018, **24**, 5723–5727; (g) C.-Q. Wang, Y. Zhang and C. Feng, Fluorine Effects on Group Migration via a Rhodium (V) Nitrenoid Intermediate, *Angew. Chem., Int. Ed.*, 2017, **56**, 14918–14922; (h) T. Yamada, Y. Shibata and K. Tanaka, Formal Lossen Rearrangement/Alkenylation or Annulation Cascade of Heterole Carboxamides with Alkynes Catalyzed by CpRh^{III} Complexes with Pendant Amides, *Chem. – Eur. J.*, 2019, **25**, 16022–16031; (i) M. Bian, H. Mawjuda, H. Gao, H. Xu, Z. Zhou and W. Yi, Lossen Rearrangement vs C–N Reductive Elimination Enabled by Rh(III)-Catalyzed C–H Activation/Selective Lactone Ring-Opening: Chemodivergent Synthesis of Quinolinones and Dihydroisoquinolinones, *Org. Lett.*, 2020, **22**, 9677–9682.
- 6 (a) S. Mochida, N. Umeda, K. Hirano, T. Satoh and M. Miura, Rhodium-catalyzed Oxidative Coupling/Cyclization of Benzamides with Alkynes via C–H Bond Cleavage, *Chem. Lett.*, 2010, **39**, 744–746; (b) T. K. Hyster and T. Rovis, Rhodium-Catalyzed Oxidative Cycloaddition of Benzamides and Alkynes via C–H/N–H Activation, *J. Am. Chem. Soc.*, 2010, **132**, 10565–10569; (c) G. Song, D. Chen, C. Pan, R. H. Crabtree and X. Li, Rh-Catalyzed Oxidative Coupling between Primary and Secondary Benzamides and Alkynes: Synthesis of Polycyclic Amides, *J. Org. Chem.*, 2010, **75**, 7487–7490; (d) N. Guimond, C. Gouliaras and K. Fagnou, Rhodium(III)-catalyzed isoquinolone synthesis: the N–O bond as a handle for C–N bond formation and catalyst turnover, *J. Am. Chem. Soc.*, 2010, **132**, 6908–6909; (e) N. Guimond, S. I. Gorelsky and K. Fagnou, Rhodium (III)-catalyzed heterocycle synthesis using an internal oxidant: improved reactivity and mechanistic studies, *J. Am. Chem. Soc.*, 2011, **133**, 6449–6457.
- 7 (a) Y. Su, M. Zhao, K. Han, G. Song and X. Li, Synthesis of 2-pyridones and iminoesters via Rh(III)-catalyzed oxidative coupling between acrylamides and alkynes, *Org. Lett.*, 2010, **12**, 5462–5465; (b) T. K. Hyster and T. Rovis, An Improved Catalyst Architecture for Rhodium(III) Catalyzed C–H Activation and its Application to Pyridone Synthesis, *Chem. Sci.*, 2011, **2**, 1606–1610; (c) X. Xu, Y. Liu and C.-M. Park, Rhodium(III)-catalyzed intramolecular annulation through C–H activation: total synthesis of (±)-antofine, (±)-septicine, (±)-tylophorine, and rosettacin, *Angew. Chem., Int. Ed.*, 2012, **51**, 9372–9376; (d) N. Quiñones, A. Seoane, R. G. Fandiño, J. L. Mascareñas and M. Gulías, Rhodium (III)-catalyzed intramolecular annulations involving amide-directed C–H activations: synthetic scope and mechanistic studies, *Chem. Sci.*, 2013, **4**, 2874–2879; (e) J. P. Krieger, D. Lesuisse, G. Ricci, M. A. Perrin, C. Meyer and J. Cossy,



- Rhodium(III)-Catalyzed C-H Activation/Heterocyclization as a Macrocyclization Strategy. Synthesis of Macrocyclic Pyridones, *Org. Lett.*, 2017, **19**, 2706–2709; (f) L. Song, G. Tian, J. van der Eycken and E. V. Van der Eycken, Intramolecular cascade annulation triggered by rhodium (III)-catalyzed sequential C(sp²)-H activation and C(sp³)-H amination, *Beilstein J. Org. Chem.*, 2019, **15**, 571–576.
- 8 (a) L. Ackermann, A. V. Lygin and N. Hofmann, Ruthenium-catalyzed oxidative synthesis of 2-pyridones through C-H/N-H bond functionalizations, *Org. Lett.*, 2011, **13**, 3278–3281; (b) R. N. P. Tulichala, M. Shankar and K. C. K. Swamy, Ruthenium-Catalyzed Oxidative Annulation and Hydroarylation of Chromene-3-carboxamides with Alkynes via Double C-H Functionalization, *J. Org. Chem.*, 2017, **82**, 5068–5079; (c) D. N. Garad and S. B. Mhaske, Ruthenium-Catalyzed Regioselective Cascade Annulation of Acrylamides with 2-Alkynoates for the Synthesis of Various 6-Oxo Nicotinic Acid Esters, *J. Org. Chem.*, 2019, **84**, 1863–1870; (d) L. Grigorjeva and O. Daugulis, Cobalt-catalyzed, aminoquinoline-directed C(sp²)-H bond alkenylation by alkynes, *Angew. Chem., Int. Ed.*, 2014, **53**, 10209–10212; (e) A. Lerchen, T. Knecht, M. Koy, C. G. Daniliuc and F. Glorius, A General Cp*Co^{III}-Catalyzed Intramolecular C-H Activation Approach for the Efficient Total Syntheses of Aromathecine, Protoberberine, and Tylophora Alkaloids, *Chem. – Eur. J.*, 2017, **23**, 12149–12152; (f) Y. Yu, L. Huang, W. Wu and H. Jiang, Palladium-catalyzed oxidative annulation of acrylic acid and amide with alkynes: a practical route to synthesize α -pyrones and pyridines, *Org. Lett.*, 2014, **16**, 2146–2149; (g) T. Matsubara, L. Ilies and E. Nakamura, Oxidative C-H Activation Approach to Pyridone and Isoquinolone through an Iron-Catalyzed Coupling of Amides with Alkynes, *Chem. – Asian J.*, 2016, **11**, 380–384; (h) B. Ma, P. Wu, X. Wang, Z. Wang, H.-X. Lin and H.-X. Dai, Efficient synthesis of Spirooxindole Pyrrolones by a Rhodium(III)-Catalyzed C-H Activation/Carbene Insertion/Lossen Rearrangement Sequence, *Angew. Chem., Int. Ed.*, 2019, **58**, 13335–13339; (i) J.-F. Tan, C. T. Bormann, K. Severin and N. Cramer, Alkynyl Triazenes as Fluoroalkyne Surrogates: Regioselective Access to 4-Fluoro-2-pyridones by a Rh(III)-Catalyzed C-H Activation–Lossen Rearrangement–Wallach Reaction, *ACS Catal.*, 2020, **10**, 3790–3796.
- 9 (a) F. Wang, Z. Qi, J. Sun, X. Zhang and X. Li, Rh(III)-catalyzed coupling of benzamides with propargyl alcohols via hydroarylation-lactonization, *Org. Lett.*, 2013, **15**, 6290–6293; (b) X. Wu, B. Wang, S. Zhou, Y. Zhou and H. Liu, Ruthenium-Catalyzed Redox-Neutral [4 + 1] Annulation of Benzamides and Propargyl Alcohols via C-H Bond Activation, *ACS Catal.*, 2017, **4**, 2494–2499; (c) W. Yi, W. Chen, F.-X. Liu, Y. Zhong, D. Wu, Z. Zhou and H. Gao, Rh(III)-Catalyzed and Solvent-Controlled Chemoselective Synthesis of Chalcone and Benzofuran Frameworks via Synergistic Dual Directing Groups Enabled Regioselective C-H Functionalization: A Combined Experimental and Computational Study, *ACS Catal.*, 2018, **8**, 9508–9519; (d) W. Gong, Z. Zhou, J. Shi, B. Wu, B. Huang and W. Yi, Catalyst-Controlled [3 + 2] and [4 + 2] Annulations of Oximes with Propargyl Alcohols: Divergent Access to Indenamines and Isoquinolines, *Org. Lett.*, 2018, **20**, 182–185; (e) X. Wu, B. Wang, Y. Zhou and H. Liu, Propargyl Alcohols as One-Carbon Synthons: Redox-Neutral Rhodium(III)-Catalyzed C-H Bond Activation for the Synthesis of Isoindolinones Bearing a Quaternary Carbon, *Org. Lett.*, 2017, **19**, 1294–1297; (f) W. Chen, F.-X. Liu, W. Gong, Z. Zhou, H. Gao, J. Shi, B. Wu and W. Yi, Hydroxyl Group-Prompted and Iridium(III)-Catalyzed Regioselective C-H Annulation of N-phenoxyacetamides with Propargyl Alcohols, *Adv. Synth. Catal.*, 2018, **360**, 2470–2475; (g) X. Wu and H. Ji, Rhodium-Catalyzed [4 + 1] Cyclization via C-H Activation for the Synthesis of Divergent Heterocycles Bearing a Quaternary Carbon, *J. Org. Chem.*, 2018, **83**, 4650–4656; (h) W. Zhou, Y.-L. Mei, B. Li, Z.-Y. Guan and Q.-H. Deng, Synthesis of β -Alkyl 2-Hydroxychalcones by Rhodium-Catalyzed Coupling of N-Phenoxyacetamides and Nonterminal Propargyl Alcohols, *Org. Lett.*, 2018, **20**, 5808–5812; (i) X. Yan, R. Ye, H. Sun, J. Zhong, H. Xiang and X. Zhou, Synthesis of 2-Arylindoles by Rhodium-Catalyzed/Copper-Mediated Annulative Coupling of N-Aryl-2-aminopyridines and Propargyl Alcohols via Selective C-H/C-C Activation, *Org. Lett.*, 2019, **21**, 7455–7479; (j) X. Hu, X. Chen, Y. Zhu, Y. Deng, H. Zeng, H. Jiang and W. Zeng, Rh(III)-Catalyzed Carboamination of Propargyl Cycloalkanols with Arylamines via Csp²-H/Csp³-Csp³ Activation, *Org. Lett.*, 2017, **19**, 3474–3477; (k) J.-L. Pan, C. Liu, C. Chen, T.-Q. Liu, M. Wang, Z. Sun and S.-Y. Zhang, Dual Directing-Groups-Assisted Redox-Neutral Annulation and Ring Opening of N-Aryloxyacetamides with 1-Alkynylcyclobutanols via Rhodium(III)-Catalyzed C-H/C-C Activations, *Org. Lett.*, 2019, **21**, 2823–2827; (l) A. Anukumar, M. Tamizmani and M. Jeganmohan, Ruthenium(II)-Catalyzed Regioselective-Controlled Allenylation/Cyclization of Benzimides with Propargyl Alcohols, *J. Org. Chem.*, 2018, **83**, 8567–8580; (m) P. Sihag and M. Jeganmohan, Regioselective Synthesis of Isocoumarins via Iridium(III)-Catalyzed Oxidative Cyclization of Aromatic Acids with Propargyl Alcohols, *J. Org. Chem.*, 2019, **84**, 2699–2712; (n) Y. Xu, M. Shen, X. Zhang and X. Fan, Selective Synthesis of Pyrazolo[1,2-a]pyrazolones and 2-Acylindoles via Rh(III)-Catalyzed Tunable Redox-Neutral Coupling of 1-Phenylpyrazolidinones with Alkynyl Cyclobutanols, *Org. Lett.*, 2020, **22**, 4697–4702; (o) L. Zhang, J. Chen, X. Chen, X. Zheng, J. Zhou, T. Zhong, Z. Chen, Y.-F. Yang, X. Jiang, Y.-B. She and C. Yu, Rh(III)-catalyzed, hydrazine-directed C-H functionalization with 1-alkynylcyclobutanols: a new strategy for 1H-indazoles, *Chem. Commun.*, 2020, **56**, 7415–7418; (p) X. Song, B. N. D. Doan, X. Zhang, R. Lee and X. Fan, Complementary C-H Functionalization Mode of Benzoylacetonitriles: Computer-Augmented Study of a Regio- and Stereoselective Synthesis of Functionalized Benzofulvenes, *Org. Lett.*, 2020, **22**, 46–51; (q) Y. Xu, B. Li, X. Zhang and X. Fan, One-Pot Synthesis of Fused N,O-Heterocycles through Rh(III)-Catalyzed Cascade



Reactions of Aromatic/Vinyl N-Alkoxyamides with 4-Hydroxy-2-alkynoates, *Adv. Synth. Catal.*, 2018, **360**, 2613–2620; (r) L. Zhang, Y. Xu, X. Zhang, X. Zhang and X. Fan, Synthesis of pyrazolone fused benzodiazepines via Rh(III)-catalyzed [4 + 3] annulation of 1-phenylpyrazolidinones with propargyl alcohols, *Org. Chem. Front.*, 2020, **7**, 2284–2290; (s) M. Sen, P. Dahiya, J. R. Premkumar and B. Sundararaju, Dehydrative Cp*Co(III)-Catalyzed C-H Bond Allenylation, *Org. Lett.*, 2017, **19**, 3699–3702; (t) W. Kong, L. H. Finger, A. M. Messinis, R. Kuniyil, J. C. A. Oliveira and L. Ackermann, Flow Rhodaelectro-Catalyzed Alkyne Annulations by Versatile C-H Activation: Mechanistic Support for Rhodium(III/IV), *J. Am. Chem. Soc.*, 2019, **141**, 17198–17206; (u) X. Wu, P. Li, Y. Lu, J. Qiao, J. Zhao, X. Jia, H. Ni, L. Kong, X. Zhang and F. Zhao, Rhodium-Catalyzed Cascade Reactions of Indoles with 4-Hydroxy-2-Alkynoates for the Synthesis of Indole-Fused

Polyheterocycles, *Adv. Synth. Catal.*, 2020, **362**, 2953–2960; (v) F. Zhao, Z. Zhou, Y. Lu, J. Qiao, X. Zhang, X. Gong, S. Liu, S. Lin, X. Wu and W. Yi, Chemo-, Regio-, and Stereoselective Assembly of Polysubstituted Furan-2(5H)-ones Enabled by Rh(III)-Catalyzed Domino C-H Alkenylation/Directing Group Migration/Lactonization: A Combined Experimental and Computational Study, *ACS Catal.*, 2021, **11**, 13921–13934; (w) Y. Xu, F. Wang, S. Yu and X. Li, Rhodium(III)-catalyzed selective access to isoindolinones via formal [4+1] annulation of arylamides and propargyl alcohols, *Chin. J. Catal.*, 2017, **38**, 1390–1398.

- 10 H. Meng, H. Xu, Z. Zhou, Z. Tang, Y. Li, Y. Zhou, W. Yi and X. Wu, Recyclable rhodium-catalyzed C-H activation/[4+2] annulation with unconventional regioselectivity at ambient temperature: experimental development and mechanistic insight, *Green Chem.*, 2022, **24**, 7012–7021.

