ChemComm

COMMUNICATION



View Article Online View Journal | View Issue

Check for updates

Cite this: Chem. Commun., 2023, 59, 3751

Received 19th December 2022, Accepted 1st March 2023

DOI: 10.1039/d2cc06901f

rsc.li/chemcomm

Chemical control of phase separation in DNA solutions[†]

Samuel Hauf 🝺 and Yohei Yokobayashi 🕩 *

We designed a series of DNA sequences comprising a trinucleotide repeat segment and a small molecule-binding aptamer. Optimization of the DNA sequences and reaction conditions enabled chemical control of phase separation of DNA condensates. Our results demonstrate a new strategy to regulate biomolecular phase transition.

Phase separation is the process by which a solution of previously miscible chemical species becomes unmixed.¹ Liquid– liquid phase separation (LLPS) has recently gained recognition as an important process in life, involved in regulatory processes,^{2,3} stress response,⁴ or certain diseases,⁵ among others. Besides its importance in natural phenomena, LLPS is also finding applications in synthetic biology.^{6,7} The process is usually driven by biopolymers such as proteins and/or nucleic acids forming dense intermolecular networks.

Nucleic acids, especially RNA, are often involved in phase separation processes observed in cells. For example, RNA involved in repeat expansion disorders can form ribonucleoprotein bodies in the cellular nucleus.⁵ In the nucleolus, DNA, RNA, and proteins phase separate from the surrounding environment to regulate transcription.⁸ Stress granules are phase separated aggregates that increase fitness under stress conditions.⁴ How exactly phase separation occurs and how it is influenced by the sequence of the biopolymers are among the subjects of ongoing research. The process is assumed to be tightly controlled by factors such as the concentration and sequence of the phase separating species, posttranslational protein modifications (*e.g.*, SUMOylation),^{4,9} temperature, and cation species.^{10,11}

For RNA, it has been shown that phase separation depends on sequence, length, and strength of RNA–RNA interaction.⁵ It has been hypothesized that the phase separation process is driven by intermolecular base pairing resulting in a complex network of RNA molecules.⁵ Because nucleic acid interactions based on Watson–Crick base pairing are relatively straightforward, it provides an opportunity to design nucleic acid sequences with programmable LLPS properties. Control of LLPS was previously shown using temperature,^{12,13} enzymatic pH-modulation,¹⁴ light,¹⁵ small molecules,^{16,17} and sizespecific diffusion of ATP through pores in lipid bilayers.¹⁸

We aimed to investigate if sequence-driven phase separation of nucleic acids can be controlled using aptamers. Aptamers are nucleic acid sequences that selectively bind target molecules (ligands) such as small molecules and proteins. Here, we show for the first time that single-stranded oligonucleotide sequences that combine the ability to form phase separated aggregates with the capability to sense a small molecule recognized by an aptamer can be designed. Our system can provide insights into chemical regulation of phase separation, and it can lead to applications in biosensing and drug delivery.

Jain and Vale showed that RNAs consisting of CAG or GGGGCC repeats can phase separate when reaching a minimum length.⁵ Their results suggested that a minimum valency and strength of interaction is necessary for nucleic acids phase separation without other factors (*e.g.*, proteins). It was hypothesized that CAG repeats drive aggregation by forming intermolecular base pairs and the GGGGCC repeats by forming intermolecular G-quadruplexes.⁵ G-quadruplex-forming RNAs have also been shown to drive phase separation in the presence of an arginine-rich peptide.¹⁹

We started by evaluating if it is possible to combine a DNA aptamer with repeat sequences to control phase separation behavior by the aptamer ligand. For this purpose, we chose the well-studied ATP aptamer²⁰ and combined it with different numbers of CAG repeats. The ATP aptamer sequence can form G-quadruplexes²⁰ which can contribute to phase separation.⁵ However, its short length may not provide sufficient intermolecular interactions to induce phase separation. We presumed that adding CAG-repeats to the 5' end of the aptamer would

Nucleic Acid Chemistry and Engineering Unit, Okinawa Institute of Science and Technology Graduate University, Onna, Okinawa, Japan 904-0495, Japan. E-mail: yohei.yokobayashi@oist.jp

[†] Electronic supplementary information (ESI) available: Materials, methods, and supplementary figures. See DOI: https://doi.org/10.1039/d2cc06901f



Fig. 1 Proposed mechanism for aptamer-controlled phase separation. CAG repeats and the ATP aptamer can phase separate in the absence of ATP due to their combined capacity for intermolecular interactions through base pairing and G-quadruplex formation. In the presence of ATP, the aptamer adopts a hairpin-like fold and can no longer engage in intermolecular interactions. The strength and valency of the CAG repeats alone are not sufficient to keep the aggregates together resulting in their dissolution.

increase the valency of interstrand interactions to trigger phase separation. Upon binding of ATP,²¹ however, the intermolecular network in the condensed phase is compromised and cannot maintain separate phases (Fig. 1).

Under the conditions determined by Jain and Vale in their RNA experiments (10 mM MgCl₂), we did not observe phase separation with the ATP aptamer fused to up to 14 CAG repeats (Fig. 2A). However, increasing MgCl₂ concentration to 100 mM or 250 mM yielded aggregates for constructs with at least 5 CAG repeats (Fig. 2A). At 100 mM MgCl₂, the aggregates were small but became clearly visible at 250 mM. This is consistent with the recent observation that higher concentrations of divalent cations enhance phase separation of ssDNA.^{11,22} The ATP aptamer alone could not form aggregates (Fig. 2A). It thus appears that both the ATP aptamer and the CAG repeats contribute to the observed phase separation. In agreement with previous results,^{5,22} a minimum length or minimum number of intermolecular interactions appears to be necessary.

The presence of 200 μ M ATP in the solution of (CAG)₁₄-ApATP (14 CAG repeats fused to ATP aptamer) with 250 mM MgCl₂ prevented the formation of visible aggregates, while presence of other nucleotides (CTP, GTP, UTP) did not affect phase separation (Fig. 2B). This suggests that the structural rearrangement of the ATP aptamer upon ATP binding²⁰ prevents phase separation. The degree of phase separation was quantified by monitoring the absorbance at 500 nm (turbidity) of the solution, and by calculating the dispersion of pixel intensities from the microscopic images (Fig. 2B). Phase



Fig. 2 Phase separated aggregates generated by combining the ATP aptamer and CAG repeats of different lengths (20 μ M) in a buffer containing 25 mM NaCl, 10 mM Tris–HCl pH 7.0, and varying amounts of MgCl₂. (A) Aggregates can be observed at 100 mM or 250 mM MgCl₂ for oligos with at least 5 CAG repeats, while no aggregates were observed at 10 mM MgCl₂ or with \leq 2 CAG repeats. Scale bars indicate 10 μ m. (B) The aggregates of (CAG)₁₄-ApATP do not form in the presence of 200 μ M ATP while other nucleotides had no effects. Phase separation was quantified by measuring the absorbance of the solution at 500 nm blanked against itself before heating or by calculating dispersion from the pixel intensities of microscopic images from three independent experiments. Individual values with mean (and standard deviation for the dispersion) are shown.

separation (turbidity) strongly depends on the Mg^{2+} as well as the DNA concentrations used (Fig. S1, ESI[†]). The aggregates were stained by SYBR Green I, indicating that they contain DNA (Fig. S2, ESI[†]). Preformed aggregates were rapidly dissolved upon addition of ATP, while addition of other NTPs had no effect (Movies S1-S5, ESI⁺). In agreement with previous reports on DNA-based systems,^{13,23} fluorescence recovery after photobleaching (FRAP) experiments showed that the aggregates behave like gels at room temperature (23 °C) and more like liquid at 43 °C (Fig. S3 and S4, ESI⁺). At 38 °C, fusion of droplets could be observed in the presence of 5% ethylene glycol indicating a liquid-like state, while aggregate formation was still prevented by addition of 1 mM ATP (Fig. S5, ESI[†]). Taken together, these results suggest that it is possible to chemically control the phase separation behavior of short DNA by incorporating an aptamer into the sequence.

Thus far, DNA phase separation was enhanced by a high Mg^{2+} concentration. Molecular crowding is also known to influence LLPS.^{24,25} Therefore, effects of the addition of uncharged crowding agents PEG200 and DMSO were investigated. Indeed, it was possible to enhance phase separation at 100 mM MgCl₂ by adding $\geq 2.5\%$ PEG200 or $\geq 5\%$ DMSO (Fig. S6A, ESI[†]). PEG200 was very efficient, but the aggregates

formed no longer responded to ATP. Aggregates formed in the presence of DMSO, however, remained sensitive to ATP (Fig. S6B, ESI[†]). PEG200 is known to enhance phase separation and stabilize G-quadruplex structures.²⁶ Therefore, it is possible that the aggregates are too stable to respond to ATP in the presence of PEG200. Alternatively, PEG200 may result in DNA precipitation.²⁷ Next, we tested the effect of Tris-HCl (pH 7.0) and NaCl concentration on aggregate formation in the presence of 10% DMSO. High concentration of Na⁺ is known to inhibit phase separation of nucleic acids.^{5,15} In our case, though, Na⁺ concentration had a negligible effect, while Tris-HCl at concentrations >100 mM inhibited phase separation (Fig. S6C, ESI[†]). The identity of the alkali cation used can influence the stability of the aggregates by affecting G-quadruplex stability.²⁸ We therefore tested if the aggregates behave differently in the presence of Li⁺ or K⁺ compared to Na⁺. We found that K⁺ stabilizes the aggregates so that they no longer respond to ATP, while Li⁺ and Na⁺ destabilize the aggregates at the highest concentration tested (200 mM, Fig. S7, ESI⁺).

Next, the concentration dependence on ATP was investigated. To minimize pH variation due to the different ATP and high Mg²⁺ concentrations, the experiments were performed at a higher Tris–HCl concentration (100 mM each of Tris–HCl (pH 7.0), NaCl, MgCl₂, 10% DMSO). A marked decrease in aggregate formation was observed between 100–1000 μ M ATP, while CTP, GTP, and UTP had no effect at the highest concentration tested (1000 μ M) (Fig. 3A, B, and D). Additionally, inactivation of the aptamer by mutation of the conserved residues²⁹ abolished the reaction to ATP, but it also negatively impacted the rate of aggregation probably due to negative effects on intermolecular interactions (Fig. 3C). These results demonstrate that the phase separation process can be finetuned by the addition of the aptamer ligand.

When working with DNA (instead of RNA), Jain and Vale used spermine to drive DNA LLPS.⁵ Spermine is a tetravalent cation at neutral pH and has been shown to drive aggregation of G-quadruplex-forming sequences.³⁰ It was also shown that the properties of the resulting aggregates depend on the sequences used.^{5,30} In our case, LLPS of (CAG)₁₄-ApATP could be triggered by 5 mM spermine, but the resulting aggregates did not react to ATP (Fig. 4A, top). The ATP aptamer alone did not phase separate in the presence of spermine (data not shown), but the construct (CAG)₂-ApATP did form aggregates that respond to ATP (Fig. 4A, middle). The change in turbidity is clearly visible to the naked eye offering a convenient read-out of the chemical regulation of phase separation (Fig. 4A, bottom) which may be useful for analytical applications.

Shorter DNA sequences can undergo phase separation in the presence of spermine compared to Mg²⁺. This suggests that the tetravalent cation spermine is a powerful driver of DNA LLPS. Therefore, a window in which the DNA aggregates respond to the aptamer ligand appears at shorter lengths.

It was recently shown that Ca^{2+} can also trigger the formation of phase separated aggregates that are more stable than aggregates formed with Mg^{2+} .²² Indeed, our construct also formed aggregates with 100 mM CaCl₂ (Fig. 4B). Interestingly,



Fig. 3 ATP concentration dependence in 100 mM Tris-HCl pH 7.0, 100 mM NaCl, 100 mM MgCl₂, and 10% DMSO. (A) (CAG)₁₄-ApATP aggregate dissolution by increasing concentrations of ATP. (B) Addition of CTP, GTP, and UTP at 1 mM did not affect the (CAG)₁₄-ApATP aggregates. (C) Aggregates formed by (CAG)₁₄-ApATP with mutations T14A and G18C ((CAG)₁₄-ApATP*) in the ATP aptamer were notably smaller, and they did not react to the addition of 1 mM ATP. (D) Sequences of the oligos used with the aptamer mutations underlined, and quantification of phase separation in panels (A) and (B) by calculating dispersion from the pixel intensities of microscopic images from three independent experiments. Individual values with mean and standard deviation are shown. Scale bars indicate 10 μ m.



aggregates in the presence of 5 mM spermine, 10 mM Tris-HCl pH 7.0 but does not react to ATP. The shorter $(CAG)_{14}$ -ApATP (20 μ M) reacts to ATP under the same conditions. This reaction is easily visualized in a cuvette (bottom). (B) Aggregate formation of $(CAG)_{14}$ -ApATP with 100 mM CaCl₂ and its reaction to ATP but not to CTP. (C) The $(CAG)_{11}$ -ApTheo (30 μ M) formed aggregates with MgCl₂ (250 mM) and DMSO (10%) that disappeared when theophylline was added but not with caffeine. Scale bars indicate 10 μ m.

these aggregates still react to ATP. Nucleic acid aggregates produced by phase separation with calcium thus seem to be more stable than with magnesium, but they can still react to sequence-encoded triggers. This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

pen Access Article. Published on 06 March 2023. Downloaded on 7/26/2025 10:57:55 PM.

Finally, we tested whether the approach is extendable to other aptamers. For this purpose, a theophylline DNA aptamer was chosen.³¹ A construct with 11 CAG repeats fused to the aptamer ((CAG)₁₁-ApTheo) phase separate when heated to 46 °C even in the presence of 1 mM caffeine, but not in the presence of 1 mM theophylline (Fig. 4C). In this case, preformed aggregates were not dissolved by the addition of the ligand. When heated to 95 $^{\circ}$ C, the difference between the condition with or without theophylline was not as pronounced (data not shown). This suggests that the ligand might affect the second transition temperature at which the phase separation occurs.²² The findings with the theophylline aptamer highlight the highly sequence- and condition-dependent behavior of nucleic acids and suggest that the chemical response may be improved through further optimization, although more work is needed to explain the underlying mechanism that causes sequence-dependent phase separation of DNA.

In summary, we have shown that it is possible to regulate the phase separation process of DNA by a small molecule by combining a trinucleotide repeat sequence and a small molecule aptamer. The oligo DNAs designed here form chemically controllable microdroplets under appropriate conditions. The chemically responsive phase separation behavior observed depends on a number of factors. First, the DNA sequences must be able to form sufficient intermolecular base pairing or tertiary interactions (e.g. Gquadruplex) to drive phase separation. Second, aptamer-ligand interaction must be able to substantially compromise the intermolecular network to dissolve or prevent phase separation. This objective can be achieved through DNA sequence design and/or optimizing reaction conditions. For example, longer DNA can favor phase separation²² but may inhibit sensitivity to aptamer-ligand binding. Similarly, a chemical environment that enhances DNA phase separation may negatively affect ligand response, but it can also allow shorter DNA sequences to form aggregates that respond to the ligand. While our systems are based on DNA, the results may also apply to RNA. It is also worth noting that Deng and Walther recently designed ATP-fuelled DNA coacervates that use ATP as a fuel for DNA ligation.32

Although previously not reported, it is possible that small molecule metabolites regulate LLPS in living cells through interaction with nucleic acids. Chemical regulation of microdroplets containing nucleic acids may find applications in synthetic biology and medicine, for example, for diagnostic (aptamer-based sensors) or therapeutic (drug delivery) tools. Alternatively, nucleic acid-based aggregates may be useful for the study of protocells or artificial cell models.^{7,12} Further refinements of the nucleic acid sequence design and analysis methods toward these goals are the subjects of future research.

We would like to thank the OIST Imaging Section, in particular Paolo Barzaghi, for support with FRAP experiments and video acquisition. This work was supported by funds from Okinawa Institute of Science and Technology Graduate University and Deutsche Forschungsgemeinschaft (DFG) - Projektnummer 452628014, Geschätszeichen: HA9374 to S. H.

Conflicts of interest

There are no conflicts to declare.

References

- 1 J. B. Clarke, J. W. Hastie, L. H. E. Kihlborg, R. Metselaar and M. M. Thackeray, *Pure Appl. Chem.*, 1994, **66**, 577–594.
- 2 S. Chong, C. Dugast-Darzacq, Z. Liu, P. Dong, G. M. Dailey, C. Cattoglio, A. Heckert, S. Banala, L. Lavis, X. Darzacq and R. Tjian, *Science*, 2018, 361, eaar2555.
- 3 D. Hnisz, K. Shrinivas, R. A. Young, A. K. Chakraborty and P. A. Sharp, *Cell*, 2017, **169**, 13–23.
- 4 J. A. Riback, C. D. Katanski, J. L. Kear-Scott, E. V. Pilipenko, A. E. Rojek, T. R. Sosnick and D. A. Drummond, *Cell*, 2017, 168, 1028–1040.
- 5 A. Jain and R. D. Vale, Nature, 2017, 546, 243-247.
- 6 C. D. Crowe and C. D. Keating, Interface Focus, 2018, 8, 20180032.
- 7 C. Xu, N. Martin, M. Li and S. Mann, Nature, 2022, 609, 1029-1037.
- 8 S. Ide, R. Imai, H. Ochi and K. Maeshima, *Sci. Adv.*, 2020, 6. eabb5953.
- 9 S. F. Banani, A. M. Rice, W. B. Peeples, Y. Lin, S. Jain, R. Parker and M. K. Rosen, *Cell*, 2016, **166**, 651–663.
- 10 J. H. Jung, A. D. Barbosa, S. Hutin, J. R. Kumita, M. Gao, D. Derwort, C. S. Silva, X. Lai, E. Pierre, F. Geng, S. B. Kim, S. Baek, C. Zubieta, K. E. Jaeger and P. A. Wigge, *Nature*, 2020, 585, 256–260.
- 11 P. Pullara, I. Alshareedah and P. R. Banerjee, *Soft Matter*, 2022, 18, 1342–1349.
- 12 A. Samanta, V. Sabatino, T. R. Ward and A. Walther, *Nat. Nanotechnol.*, 2020, **15**, 914–921.
- 13 Y. Sato, T. Sakamoto and M. Takinoue, Sci. Adv., 2020, 6, eaba3471.
- 14 H. Karoui, M. J. Seck and N. Martin, Chem. Sci., 2021, 12, 2794-2802.
- 15 N. Martin, L. Tian, D. Spencer, A. Coutable-Pennarun, J. L. R. Anderson and S. Mann, Angew. Chem., Int. Ed., 2019, 58, 14594–14598.
- 16 W. M. Babinchak, B. K. Dumm, S. Venus, S. Boyko, A. A. Putnam, E. Jankowsky and W. K. Surewicz, *Nat. Commun.*, 2020, **11**, 5574.
- 17 M. Ramesh, C. Balachandra, P. Baruah and T. Govindaraju, *J. Pept. Sci.*, 2022, e3465.
- 18 S. Deshpande, F. Brandenburg, A. Lau, M. G. F. Last, W. K. Spoelstra, L. Reese, S. Wunnava, M. Dogterom and C. Dekker, *Nat. Commun.*, 2019, 10, 1800.
- 19 M. Tsuruta, T. Torii, K. Kohata, K. Kawauchi, H. Tateishi-Karimata, N. Sugimoto and D. Miyoshi, *Chem. Commun.*, 2022, 58, 12931–12934.
- 20 D. E. Huizenga and J. W. Szostak, Biochemistry, 1995, 34, 656-665.
- 21 C. H. Lin and D. J. Patel, Chem. Biol., 1997, 4, 817-832.
- 22 W. Liu, A. Samanta, J. Deng, C. O. Akintayo and A. Walther, *Angew. Chem., Int. Ed.*, 2022, **61**, e202208951.
- 23 R. Merindol, S. Loescher, A. Samanta and A. Walther, *Nat. Nano*technol., 2018, 13, 730–738.
- 24 J. Hochmair, C. Exner, M. Franck, A. Dominguez-Baquero, L. Diez, H. Brognaro, M. L. Kraushar, T. Mielke, H. Radbruch, S. Kaniyappan, S. Falke, E. Mandelkow, C. Betzel and S. Wegmann, *EMBO J.*, 2022, 41, e108882.
- 25 S. Park, R. Barnes, Y. Lin, B.-J. Jeon, S. Najafi, K. T. Delaney, G. H. Fredrickson, J.-E. Shea, D. S. Hwang and S. Han, *Commun. Chem.*, 2020, 3, 83.
- 26 R. Hansel, F. Lohr, S. Foldynova-Trantirkova, E. Bamberg, L. Trantirek and V. Dotsch, *Nucleic Acids Res.*, 2011, 39, 5768–5775.
- 27 A. Schmitz and D. Riesner, Anal. Biochem., 2006, 354, 311-313.
- 28 E. Largy, J.-L. Mergny and V. Gabelica, in *The Alkali Metal Ions: Their Role for Life*, ed. A. Sigel, H. Sigel and R. K. O. Sigel, Springer International Publishing, Cham, 2016, pp. 203–258, DOI: 10.1007/978-3-319-21756-7_7.
- 29 R. Nutiu and Y. Li, J. Am. Chem. Soc., 2003, 125, 4771-4778.
- 30 A. M. Williams, R. R. Poudyal and P. C. Bevilacqua, *Biochemistry*, 2021, 60, 2715–2726.
- 31 P. J. Huang and J. Liu, ACS Chem. Biol., 2022, 17, 2121-2129.
- 32 J. Deng and A. Walther, Chem, 2020, 6, 3329-3343.