

## PAPER

[View Article Online](#)  
[View Journal](#) | [View Issue](#)Cite this: *Dalton Trans.*, 2023, **52**,  
3308The coordination chemistry of 2,4,6-oxy  
functionalised 1,3,5-triphosphinines†Anne Sofie Abels,<sup>a</sup> Frederik Eiler,<sup>a</sup> Grégoire Le Corre,<sup>a</sup> Pascal Jurt,<sup>a</sup>  
Michael Wörle,<sup>a</sup> René Verel,<sup>a</sup> Zoltan Benkő<sup>b</sup> and Hansjörg Grützmacher<sup>\*,a</sup>

A number of stable group 6 metal complexes bearing 2,4,6-oxy functionalised 1,3,5-triphosphinines, phosphorus containing heterocyclic ligands with a central  $C_3P_3$  core, were synthesised such that a complete series of  $[M(P_3C_3(OX)_3)(CO)_3]$  compounds is obtained [ $M = Cr(0), Mo(0), W(0)$ ;  $X = H, Si^tBuPh_2, B(ipc)_2$ ]. In all complexes, the triphosphinine coordinates in a  $\eta^6$ -binding mode *via* the delocalized  $6\pi$ -system of the ring. The ligand properties can be tuned by changing the substituent on the oxygen centre. The  $\pi$ -electron accepting properties of the ligand increases in the following order:  $P_3C_3(OH)_3 < P_3C_3(OSi^tBuPh_2)_3 < P_3C_3(OB(ipc)_2)_3$ . This trend is reflected in the structures determined by X-ray crystallography, and the  $\nu(CO)$  stretching frequencies determined by IR spectroscopy. The collected data raise questions with respect to the frequently made assumption that phosphinines act as stronger  $\pi$ -acceptors with respect to arenes and thereby deplete electron density at the metal centres. With  $P_3C_3(OH)_3$  as an  $\eta^6$ -coordinated ligand further molecules can be coordinated in the second coordination sphere *via* hydrogen bonds, which may be of interest for the construction of coordination polymers.

Received 7th January 2023,  
Accepted 2nd February 2023

DOI: 10.1039/d3dt00057e

rsc.li/dalton

## Introduction

Since Märkl and Ashe successfully prepared the first monophosphinines,<sup>1,2</sup> these phosphorus heterocycles and their coordination chemistry have been studied extensively.<sup>3–10</sup> Nowadays, these compounds serve as ligands in transition metal complexes that are active homogeneous catalysts, for example in hydroformylation, hydrogenation and oligomerisation reactions.<sup>6,9,11</sup> In contrast to monophosphinines, the chemistry of phosphinines containing three phosphorus centres, namely triphosphinines, is much less developed.<sup>12,13</sup> Generally, 1,3,5-triphosphinines are obtained by transition metal mediated trimerisation of kinetically stabilised phosphalkynes.<sup>14–17</sup> The range of reported 1,3,5-triphosphinines is rather limited and most carry tertiary butyl, *t*Bu, substituents or related bulky hydrocarbon groups attached to the C atom of the ring. In combination, these studies led to the assumption that phosphorus-containing heterocycles have

strong  $\pi$ -accepting properties with respect to arenes, making them excellent ligands for binding low-valent electron rich metal centres in a  $\eta^6$ -coordination mode.<sup>4,18–26</sup>


In 2017, Grützmacher and co-workers reported on a new synthetic route to access functionalized triphosphinines by the reaction of sodium phosphoethynolate,  $[Na(OCP)]$ , with commercially available (–)-B-chlorodi(isopinocampheyl)borane,  $(ipc)_2B-Cl$ , forming the boryloxy-functionalised phosphalkyne  $(ipc)_2B-O-C\equiv P$  as slightly thermodynamically preferred product that trimerises in absence of a catalyst to form a 1,3,5-triphosphinine with a central six-membered  $P_3C_3$  ring (Scheme 1,  $P_3C_3-B$ ) on a multi-gram scale.<sup>27</sup> This compound is the first triphosphinine that can readily be derivatised and thereby the electronic properties can be tuned by changing the substituent at the oxygen atom. Treatment of triphosphinine  $P_3C_3-B$  with an excess of alcohol yields the deprotected tris (hydroxyl) triphosphinine  $P_3C_3(OH)_3$  ( $P_3C_3-H$ ), the phosphorus analogue of cyanuric acid. But unlike cyanuric acid,  $P_3C_3(OH)_3$  is rather labile, therefore could not be isolated and was only stable in solution.

Cyanuric acid is an industrially useful chemical and finds use in the production of herbicides,<sup>28,29</sup> dyes<sup>30</sup> and flame retardants.<sup>31–33</sup> Furthermore, its trimeric structure and the ability to form hydrogen bonds makes cyanuric acid an attractive precursor for cross linkers in polymers<sup>34,35</sup> and an building block in supramolecular networks.<sup>36–41</sup> The formation of a stable  $P_3C_3(OH)_3$  compound may therefore be of interest. Furthermore, these type of  $P_3C_3$  rings have shown that they are

<sup>a</sup>Department of Chemistry and Applied Biosciences, ETH Zürich, Vladimir-Prelog Weg 1, Hönggerberg, 8093 Zürich, Switzerland. E-mail: hgruetzmacher@ethz.ch, abelsa@inorg.chem.ethz.ch

<sup>b</sup>Department of Inorganic and Analytical Chemistry, Budapest University of Technology and Economics, 1111 Budapest, Műegyetem rakpart 3, Hungary. E-mail: benko.zoltan@vbk.bme.hu

† Electronic supplementary information (ESI) available. CCDC 2234729–2234735. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d3dt00057e>



**Scheme 1** Previous synthesis of borylated compound P<sub>3</sub>C<sub>3</sub>-B from the reaction of Na[OCp] with (ipc)<sub>2</sub>BCl, the formation of the phosphorus analogue of cyanuric acid (P<sub>3</sub>C<sub>3</sub>-H) and the synthesis of molybdenum 2,4,6-oxy functionalised 1,3,5-triposphinine complexes.

excellent ligands and coordinate in a  $\eta^6$ -binding mode through the  $\pi$ -system of the ring to a molybdenum centre (Scheme 1). However, the molybdenum complex with P<sub>3</sub>C<sub>3</sub>(OH)<sub>3</sub> as ligand could not be isolated, therefore key structural information and spectroscopic data is missing.

In this study we investigated the coordination chemistry of these 2,4,6-oxy functionalised 1,3,5-triposphinines with group 6 metals in depth and we were able to isolate and analyse the phosphorus analogue of cyanuric acid as ligand in coordination compounds. This series of group 6 metal 2,4,6-tri-oxy-1,3,5-triposphinine complexes bearing different substituents on the oxygen centre allowed us to study the physical properties of this class of compounds in detail by single-crystal X-ray structure determinations (XRD's), IR-, and NMR-spectroscopic analyses.

## Results and discussion

The Mo(0) complexes [Mo{P<sub>3</sub>C<sub>3</sub>[OB(ipc)<sub>2</sub>]<sub>3</sub>}(CO)<sub>3</sub>] **Mo-P<sub>3</sub>C<sub>3</sub>-B** and [Mo{P<sub>3</sub>C<sub>3</sub>[O(SitBuPh<sub>2</sub>)<sub>3</sub>](CO)<sub>3</sub>] **Mo-P<sub>3</sub>C<sub>3</sub>-Si** were previously synthesised from [Mo(Mes)(CO)<sub>3</sub>] and isolated in crystalline form. The complex [Mo{P<sub>3</sub>C<sub>3</sub>(OH)<sub>3</sub>}(CO)<sub>3</sub>] **Mo-P<sub>3</sub>C<sub>3</sub>-H** could only be spectroscopically observed in solution.<sup>27</sup> In straightforward ligand exchange reactions, the analogous tricarbonyl chromium and tungsten complexes of P<sub>3</sub>C<sub>3</sub>-B were synthesised from [M(NCMe)<sub>3</sub>(CO)<sub>3</sub>] (M = Cr and W) (Fig. 1). Single crystals of Cr-P<sub>3</sub>C<sub>3</sub>-B and W-P<sub>3</sub>C<sub>3</sub>-B were grown by slow evaporation of *n*-hexane solutions and analysed by single crystal diffraction. The silyloxy-substituted compound P<sub>3</sub>C<sub>3</sub>[O(SitBuPh<sub>2</sub>)<sub>3</sub>], P<sub>3</sub>C<sub>3</sub>-Si, was prepared in solution<sup>27</sup> and addition of the group 6 metal precursor to the reaction mixture yielded compounds Cr-P<sub>3</sub>C<sub>3</sub>-Si and W-P<sub>3</sub>C<sub>3</sub>-Si (Fig. 1). The reaction time varied from 1 hour for Cr-P<sub>3</sub>C<sub>3</sub>-Si to 5 days for W-P<sub>3</sub>C<sub>3</sub>-Si, the latter was heated to 40 °C to accelerate the reaction. The use of multiple reagents explains the rather low isolated yields


(10–11%) in these one-pot syntheses of M-P<sub>3</sub>C<sub>3</sub>-Si (M = Cr, W) and it proved to be difficult to separate the desired complexes from a number of side products. Slow evaporation of *n*-hexane solutions of Cr-P<sub>3</sub>C<sub>3</sub>-Si and W-P<sub>3</sub>C<sub>3</sub>-Si gave single crystals suitable for X-ray diffraction studies.

To obtain the tris(hydroxyl)triposphinine complexes, the group 6 metal tricarbonyl complexes of P<sub>3</sub>C<sub>3</sub>-B were dissolved in *n*-hexane and an excess of methanol was added to give a biphasic system. After shaking the reaction flask, the subsequently formed P<sub>3</sub>C<sub>3</sub>(OH)<sub>3</sub> complexes could be isolated as orange (M = Cr) or yellow (M = Mo, W) solids from the methanol phase. The *n*-hexane phase contains the methoxyborane-(ipc)<sub>2</sub>B(OMe), which could easily be separated. Coordination of the P<sub>3</sub>C<sub>3</sub>(OX)<sub>3</sub> rings [X = B(ipc)<sub>2</sub>:  $\delta(^{31}\text{P}) = 182.7$  ppm, X = SitBuPh<sub>2</sub>:  $\delta(^{31}\text{P}) = 177.5$  ppm; X = H:  $\delta(^{31}\text{P}) = 133.7$  ppm]<sup>27</sup> leads to a significant coordination shift  $\Delta\delta = \delta(\text{P}_3\text{C}_3[\text{OX}]_3) - \delta([\text{M}\{\text{P}_3\text{C}_3[\text{OX}]_3(\text{CO})_3\}])$  of the <sup>31</sup>P NMR resonances to lower frequencies in the range of 161 to 198 ppm, which is especially pronounced in case of the parent tris(hydroxyl)triposphinine P<sub>3</sub>C<sub>3</sub>(OH)<sub>3</sub> (see Table 1, first entry). The  $\Delta\delta$  values increase slightly in the order Cr < Mo < W within each series of ligand and SitBuPh<sub>2</sub> < B(ipc)<sub>2</sub> < H with respect of the group X bound to the oxygen centres of the P<sub>3</sub>C<sub>3</sub>(OX)<sub>3</sub> ring. Comparable strong coordination shifts ( $\Delta\delta > 160$  ppm to lower frequencies) were also observed when 2,4,6-triphenyl-phosphinine, PC<sub>5</sub>H<sub>2</sub>(Ph)<sub>3</sub>, or 2,4,6-*tert*-butyltriposphinine, P<sub>3</sub>C<sub>3</sub>(*t*Bu)<sub>3</sub>, are  $\eta^6$ -coordinated to a M(CO)<sub>3</sub> fragment.<sup>42</sup>

Single crystals of Cr-P<sub>3</sub>C<sub>3</sub>-B, W-P<sub>3</sub>C<sub>3</sub>-B, Cr-P<sub>3</sub>C<sub>3</sub>-Si, and W-P<sub>3</sub>C<sub>3</sub>-Si were grown by slow evaporation of *n*-hexane solutions and analysed by single crystal diffraction (XRD) (Fig. 1). Single crystals of the coordinated P<sub>3</sub>C<sub>3</sub>(OH)<sub>3</sub> group 6 complexes were obtained by cooling a concentrated *n*-hexane solution of Cr-P<sub>3</sub>C<sub>3</sub>-H, Mo-P<sub>3</sub>C<sub>3</sub>-H, or W-P<sub>3</sub>C<sub>3</sub>-H in the presence of *t*BuOH. The structures of the P<sub>3</sub>C<sub>3</sub>-H metal complexes determined by XRD show interactions of the OH groups of the ring with the OH groups of *t*BuOH *via* hydrogen bonding. Related O-H...O hydrogen bonding was previously postulated for Mo-P<sub>3</sub>C<sub>3</sub>-H in THF solution.<sup>27</sup> Table 1 summarises selected spectroscopic data and bond lengths and angles of the solid state structures of the nine different chromium, molybdenum and tungsten triphosphinine complexes, bearing different functionalities at the oxygen centres in 1,3,5 position of the P<sub>3</sub>C<sub>3</sub> ring.

In these piano stool complexes, the P<sub>3</sub>C<sub>3</sub> ring is planar and binds in an  $\eta^6$ -coordination mode to the metal centre (Fig. 1). The P-C bond lengths (1.76–1.78 Å) of the coordinated P<sub>3</sub>C<sub>3</sub> rings are slightly elongated compared to the uncoordinated triphosphinine ring (1.736 Å).<sup>27</sup> Compared to P<sub>3</sub>C<sub>3</sub>[OB(ipc)<sub>2</sub>]<sub>3</sub> (C-P-C<sub>av</sub>: 104°; P-C-P<sub>av</sub> 136°), the C-P-C angles become slightly more acute (P-C-P<sub>av</sub> 101.9°) while the P-C-P angles widen up slightly (P-C-P<sub>av</sub> 138.0°) upon coordination to a M(CO)<sub>3</sub> fragment. Within the complexes, these values vary rather little [C-P-C 100.6(3)–102.9(3)°; P-C-P 137.0(3)–140.2(4)], but independently of M, the boryloxy substituted complexes show the smallest C-P-C (largest P-C-P) angle, while the hydroxyl substituted show the largest C-P-C (smallest P-C-P) angle.





**Fig. 1** Synthesis of complexes  $M-P_3C_3-B$  ( $M = Cr, W$ ),  $M-P_3C_3-Si$  ( $M = Cr, W$ ) and  $M-P_3C_3-H$  ( $M = Cr, Mo, W$ ). The crystal structures of  $W-P_3C_3-B$ ,  $W-P_3C_3-Si$  and  $W-P_3C_3-H$  are given below the respective chemical structures. Hydrogen atoms and solvent molecules are omitted for clarity. Average interatomic distances are summarised in Table 1. The crystal structures of  $Cr-P_3C_3-B$ ,  $Cr-P_3C_3-Si$ ,  $Cr-P_3C_3-H$  and  $Mo-P_3C_3-H$  are included in the ESI.†

**Table 1** Selected average bond distances [Å] and angles [°] for  $M-P_3C_3-B$ ,  $M-P_3C_3-Si$  and  $M-P_3C_3-H$  ( $M = Cr, Mo, W$ ) obtained from XRD data with single crystals. Coordination shifts  $\Delta\delta$  in the  $^{31}P$  NMR spectra are defined by  $\delta(P_3C_3[OX]_3) - \delta([M(P_3C_3[OX]_3)(CO)_3])$  and are given in ppm

	$Cr-P_3C_3-B$	$Cr-P_3C_3-Si$	$Cr-P_3C_3-H$	$Mo-P_3C_3-B$	$Mo-P_3C_3-Si$	$Mo-P_3C_3-H$	$W-P_3C_3-B$	$W-P_3C_3-Si$	$W-P_3C_3-H$
$\Delta\delta$	166	161	191	176	167	193	185	177	198
M-ct	1.6407(8)	1.6499(5)	1.6817(8)	1.753(1)	1.778(6)	1.8157(11)	1.7462(11)	1.7743(19)	1.802(5)
P-C	1.7578(13)	1.7577(7)	1.7597(9)	1.765(4)	1.769(2)	1.7717(18)	1.7687(18)	1.767(3)	1.773(8)
M-CO	1.8947(18)	1.8714(11)	1.8667(12)	2.022(4)	2.005(2)	1.989(3)	2.024(3)	2.009(5)	2.001(12)
C-O	1.1370(18)	1.1453(12)	1.1487(16)	1.139(5)	1.143(2)	1.156(4)	1.129(3)	1.137(5)	1.130(13)
C-P-C	101.46(7)	102.25(5)	102.69(8)	101.19(2)	101.93(8)	102.7(3)	101.13(11)	101.37(18)	102.2(6)
P-C-P	138.53(10)	137.7(6)	137.28(8)	138.7(2)	137.88(2)	137.3(18)	138.8(12)	138.4(3)	137.7(6)

For every metal, the M-ct distance is the longest with  $P_3C_3(OH)_3$  as ligand ( $Cr-P_3C_3-H = 1.682$  Å,  $Mo-P_3C_3-H = 1.816$  Å,  $W-P_3C_3-H = 1.802$  Å), followed by the siloxylated complexes ( $Cr-P_3C_3-Si = 1.650$  Å,  $Mo-P_3C_3-Si = 1.778$  Å,  $W-P_3C_3-Si = 1.774$  Å) and the boryloxylated  $P_3C_3$  metal complexes ( $Cr-P_3C_3-B = 1.619$  Å,  $Mo-P_3C_3-B = 1.753$  Å,  $W-P_3C_3-B = 1.746$  Å). Although the differences are small (within  $3\sigma$ ), the Mo-complexes show always slightly longer Mo-ct distances compared to the corresponding W-ct distances despite the increasing covalent radii [ $r_{cov}(Mo) = 1.54$  Å;  $r_{cov}(W) = 1.62$  Å]. Complexes with chromium show expectedly significantly shorter Cr-ct distances [ $r_{cov}(Cr) = 1.39$  Å].<sup>43</sup> (The atomic van der Waals radii of Cr, Mo, and W [ $r_{vdw}(Cr) = 1.89$  Å;  $r_{vdw}(Mo) = 2.09$  Å,  $r_{vdw}(W) = 2.10$  Å] reflect better the observed trends in the  $M-P_3C_3-X$  complexes with M in the oxidation state 0, but also suggest

that the M-ct bond lengths should steadily increase in the order  $W > Mo > Cr$ ).<sup>44</sup> Indeed this trend is observed for  $[M(P_3C_3(tBu)_3)(CO)_3]$  ( $M = Cr, Mo, W$ ), whose structures were calculated by DFT methods. Here the increase in M-ct distances corresponds to the van der Waals radii of the metals.<sup>23,44</sup>

In  $[M(L)_x(CO)_y]$  complexes, a ligand L with  $\pi$ -accepting properties competes with carbonyl co-ligands for  $M \rightarrow \pi^*(CO)$  electron back donation. Consequently the electronic properties of a ligand L in  $[M(L)_x(CO)_y]$  complexes are reflected in the M-C and C-O distances as well as in the CO stretching frequencies  $\nu(CO)$  of the carbonyl co-ligands.<sup>45</sup> A ligand L of high  $\pi$ -electron accepting property will lead to long M-C bonds to the CO ligand, short C-O bonds and  $\nu(CO)$  at high wave numbers. In case of complexes with  $Cr(CO)_3$  or  $Mo(CO)_3$  fragments, the ones with  $P_3C_3-B$  rings show the longest M-CO dis-



tances while, on the contrary, the M–CO distances are the shortest in complexes with the  $P_3C_3$ –H ligand. As expected, the opposite trends are observed with respect to the C–O bond lengths.

Another frequently used instrument in order to evaluate the electronic properties of L in  $[M(L)_x(CO)_y]$  complexes are the  $\nu(CO)$  of the carbonyl co-ligands. It is generally accepted that the relative ratio of ligand-to-metal electron donation d ( $M \leftarrow L$ ) to metal-to-ligand back donation b ( $M \rightarrow L$ ), denoted here as d/b, decreases in the order  $CO \approx$  alkynes  $<$  olefins  $<$  carbenes  $<$  phosphines  $<$  amines (simplified meaning that the  $\pi$ -electron accepting property of the respective ligand decreases in this order).<sup>46–48</sup> Consequently a decrease of  $\nu(CO)$  is expected along this ligand line. In Tables 2–4 a compilation of IR data is given for selected, comparable chromium, molybdenum, and tungsten tricarbonyl complexes with simple monodentate  $\sigma$ -donor/ $\pi$ -acceptor ligands, arene ligands, and phosphinine ligands. In all complexes the  $M(CO)_3$  fragment

**Table 2** IR stretching frequencies in  $cm^{-1}$  for the CO groups in various chromium tricarbonyl complexes with  $\sigma$ -donor/ $\pi$ -acceptor, arene, and phosphinine ligands. The stretching frequencies  $\nu(CO)$  of  $Cr-P_3C_3-B$ ,  $Cr-P_3C_3-Si$ , and  $Cr-P_3C_3-H$  were recorded by ATR-IR spectroscopy from crystalline products. Substituents in the arene ring are in 1,3,5-position, in phosphines in 2,4,6-position

	$\nu_{CO}$ ( $cm^{-1}$ )
1	$[Cr(CO)_6]^{49}$ 1985
2	$[Cr\{P(OMe)_3\}(CO)_3]^{50}$ 1962, 1888, 1874
3	$[Cr\{PMe_2Ph\}(CO)_3]^{51}$ 1920, 1820
4	$[Cr(NCMe)_3(CO)_3]^{52}$ 1910, 1782
5	$[Cr(C_6H_6)(CO)_3]^{53}$ 1971, 1860
6	$[Cr(Mes)(CO)_3]^{54}$ 1965, 1852
7	$[Cr\{C_6H_3(tBu)_3\}(CO)_3]^{55}$ 1947, 1871
8	$[Cr\{C_6H_3(OH)_3\}(CO)_3]^{56}$ 1945, 1850
9	$[Cr\{C_6H_3(CO_2Me)_3\}(CO)_3]^{57}$ 2003, 1947
10	$[Cr\{C_6H_3(OSO_2CF_3)_3\}(CO)_3]^{56}$ 2019, 1947
11	$[Cr\{PC_5H_2(Ph)_3\}(CO)_3]^{58}$ 1976, 1915, 1907
12	$[Cr\{P_3C_3(tBu)_3\}(CO)_3]^{23}$ 1979, 1926
13	$[Cr\{P_3C_3(OH)_3\}(CO)_3]$ $Cr-P_3C_3-H$ 1980, 1921
14	$[Cr\{P_3C_3[O(SiBuPh_2)_3]_3\}(CO)_3]$ $Cr-P_3C_3-Si$ 1983, 1927
15	$[Cr\{P_3C_3[OB(ipc)_2]_3\}(CO)_3]$ $Cr-P_3C_3-B$ 2012, 1967

**Table 3** IR stretching frequencies in  $cm^{-1}$  for the CO groups in various molybdenum tricarbonyl complexes with  $\sigma$ -donor/ $\pi$ -acceptor, arene, and phosphinine ligands. The stretching frequencies  $\nu(CO)$  of  $Mo-P_3C_3-B$ ,  $Mo-P_3C_3-Si$ , and  $Mo-P_3C_3-H$  were recorded by ATR-IR spectroscopy from crystalline products. Substituents in the arene ring are in 1,3,5-position, in phosphines in 2,4,6-position

	$\nu_{CO}$ ( $cm^{-1}$ )
1	$[Mo(CO)_6]^{49}$ 1985
2	$[Mo\{P(OMe)_3\}(CO)_3]^{50}$ 1974, 1898, 1884
3	$[Mo\{PMe_2Ph\}(CO)_3]^{51}$ 1939, 1832, 1824
4	$[Mo(NCMe)_3(CO)_3]^{52}$ 1915, 1783
5	$[Mo(C_6H_6)(CO)_3]^{49}$ 1987, 1916
6	$[Mo(Mes)(CO)_3]^{49}$ 1973, 1901
7	$[Mo\{PC_5H_2(Ph)_3\}(CO)_3]^{42}$ 1996, 1942, 1933
8	$[Mo\{P_3C_3(OH)_3\}(CO)_3]$ $Mo-P_3C_3-H$ 1988, 1925
9	$[Mo\{P_3C_3[O(SiBuPh_2)_3]_3\}(CO)_3]$ $Mo-P_3C_3-Si$ 1993, 1941
10	$[Mo\{P_3C_3[OB(ipc)_2]_3\}(CO)_3]$ $Mo-P_3C_3-B$ 2022, 1972

**Table 4** IR stretching frequencies in  $cm^{-1}$  for the CO groups in various tungsten tricarbonyl complexes with  $\sigma$ -donor/ $\pi$ -acceptor, arene, and phosphinine ligands. The stretching frequencies  $\nu(CO)$  of  $W-P_3C_3-B$ ,  $W-P_3C_3-Si$ , and  $W-P_3C_3-H$  were recorded by ATR-IR spectroscopy from crystalline products. Substituents in the arene ring are in 1,3,5-position, in phosphines in 2,4,6-position

	$\nu_{CO}$ ( $cm^{-1}$ )
1	$[W(CO)_6]^{49}$ 1981
2	$[W\{P(OMe)_3\}(CO)_3]^{50}$ 1970, 1893, 1878
3	$[W\{PMe_2Ph\}(CO)_3]^{51}$ 1932, 1926
4	$[W(NCMe)_3(CO)_3]^{52}$ 1915, 1793
5	$[W(C_6H_6)(CO)_3]^{59}$ 1990, 1916
6	$[W(Mes)(CO)_3]^{59}$ 1975, 1902
7	$[W\{P_3C_3(OH)_3\}(CO)_3]$ $W-P_3C_3-H$ 1987, 1921
8	$[W\{P_3C_3[O(SiBuPh_2)_3]_3\}(CO)_3]$ $W-P_3C_3-Si$ 1989, 1921
9	$[W\{P_3C_3[OB(ipc)_2]_3\}(CO)_3]$ $W-P_3C_3-B$ 2019, 1963

( $M = Cr, Mo, W$ ) adopts a facial coordination mode and reference is made only to the symmetrical  $\nu(CO)$  stretching frequency of  $A_1$  symmetry at higher wave numbers (in case of deviation from local  $C_{3v}$  symmetry of the  $M(CO)_3$  fragment, reference is made to  $\nu(CO)$  at the highest wave number). As reference compounds the hexacarbonyl complexes  $[M(CO)_6]$  are chosen, for which  $\nu(CO)$  is quite similar independently on the metal ( $M = Cr$ : 1985  $cm^{-1}$ ;  $Mo = 1985$   $cm^{-1}$ ;  $M = W$ : 1981  $cm^{-1}$ ). As expected, the replacement of three CO groups in facial position by ligands with an increasingly stronger  $\sigma$  or  $\pi$ -donating but poorer  $\pi$ -accepting property – that is an increasingly higher d/b ratio – as in the series  $P(OMe)_3 < PMe_2Ph < NCMe$  – will increasingly lower the  $\nu(CO)$  stretching frequency as a result of increasing electron density on the metal centre (see entries 2–4 in each table). This effect is remarkably similar for each metal and the variation  $\Delta\nu(CO)$  from  $[M(CO)_6]$  to  $[M(NCMe)_3(CO)_3]$  is about 70  $cm^{-1}$  to lower wave numbers. In the benzene complexes  $[M(C_6H_6)(CO)_3]$  a slight blue-shift of  $\nu(CO)$  to higher wave numbers is observed for  $M = Mo$  (1987  $cm^{-1}$ ) or  $W$  (1990  $cm^{-1}$ ) (entries 5 in Tables 3 and 4), while in  $[Cr(C_6H_6)(CO)_3]$  (1971  $cm^{-1}$ ) a red-shift by 14  $cm^{-1}$  to lower wave numbers occurs (entry 5 in Table 2).  $\sigma$ -Donating substituents like methyl groups in 1,3,5-position of the arene ring increase the d/b ratio and lower the  $\nu(CO)$  by 6 to 15  $cm^{-1}$  (entries 6 in Tables 2–4). The IR data for a range of Cr(0) complexes with arene ligands functionalised in 1,3,5-position are available (see entries 7–10 in Table 2).  $\sigma$ -Electron and  $\pi$ -electron donating substituents like *t*Bu or hydroxyl groups shift  $\nu(CO)$  by about 25  $cm^{-1}$  to lower wave numbers. Electron accepting groups like carboxy groups,  $CO_2Me$ , or triflate groups,  $OSO_2CF_3$ , in 1,3,5-position shift  $\nu(CO)$  to higher wave numbers by 32  $cm^{-1}$  or 48  $cm^{-1}$ , respectively. Hence, the range by which  $A_1$   $\nu(CO)$  can be changed in these type of complexes is about 70  $cm^{-1}$ .

What effect will be evoked when one CH unit is replaced by a phosphorus centre? In the monophosphinine complexes  $[Cr\{PC_5H_2(2,4,6-Ph)_3\}(CO)_3]$  (1976  $cm^{-1}$ , entry 11 in Table 2) and  $[Mo\{PC_5H_2(2,4,6-Ph)_3\}(CO)_3]$  (1996  $cm^{-1}$ , entry 7 in Table 3)  $\nu(CO)$  shifts slightly (5, 9  $cm^{-1}$ ) with respect to the benzene








Fig. 2  $M(\text{CO})_3$  complex of bis(imidazolium)-1,3-diphosphete-diide (IPD) ( $M = \text{Cr}, \text{Mo}$ ).

complexes, or modestly ( $11, 23 \text{ cm}^{-1}$ ) with respect to the mesitylene complexes to higher wave numbers indicating indeed some depletion of the electron density at the metal centre with respect to an arene ligand. The further substitution of two additional CH units by phosphorus leading to triphosphinine ligands raises the  $A_1 \nu(\text{CO})$  to even higher frequencies. A significant blue-shift of  $32 \text{ cm}^{-1}$  is observed comparing  $[\text{Cr}\{\text{C}_6\text{H}_3(\text{tBu})_3\}(\text{CO})_3]$  ( $1947 \text{ cm}^{-1}$ ; entry 7 in Table 2) and  $[\text{Cr}\{\text{P}_3\text{C}_3(\text{tBu})_3\}(\text{CO})_3]$  ( $1979 \text{ cm}^{-1}$ ; entry 13 in Table 2). As in arene complexes,  $\sigma$ -donating *t*Bu substituents in 1,3,5-position of a triphosphinine ligand have a very similar effect than  $\pi$ -donating hydroxyl groups as a comparison between  $[\text{Cr}\{\text{P}_3\text{C}_3(\text{OH})_3\}(\text{CO})_3]$  ( $\text{Cr}-\text{P}_3\text{C}_3-\text{H}$ ) ( $1980 \text{ cm}^{-1}$ , entry 14 Table 2) and  $[\text{Cr}\{\text{P}_3\text{C}_3(\text{tBu})_3\}(\text{CO})_3]$  ( $1979 \text{ cm}^{-1}$ , entry 13 Table 2) shows.

How is  $\nu(\text{CO})$  influenced by electronically different substituents in 2,4,6-position of a triphosphinine ligand? This question can be partly answered by inspecting the last three entries 14–16 in Table 2. While replacing hydroxyl groups in  $\text{Cr}-\text{P}_3\text{C}_3-\text{H}$  by siloxy groups has a marginal influence, the replacement by boryloxy groups has a significant impact. The complex  $[\text{Cr}\{\text{P}_3\text{C}_3[\text{OB}(\text{ipc})_2]_3\}(\text{CO})_3]$  ( $\text{Cr}-\text{P}_3\text{C}_3-\text{B}$ ) has the highest  $\nu(\text{CO}) = 2012 \text{ cm}^{-1}$  (entry 16 Table 2) of all triphosphinine complexes indicating that a  $\text{B}(\text{ipc})_2$  group efficiently abstracts electron density from the  $\pi$ -electron donating oxygen centres connected to the  $\text{P}_3\text{C}_3$  ring thereby depleting the overall electron density at the chromium centre. But the variation between the complex with an “electron rich”  $\text{P}_3\text{C}_3(\text{tBu})_3$  [or  $\text{P}_3\text{C}_3(\text{OH})_3$ ] ligand and the one with an “electron poor”  $\text{P}_3\text{C}_3[\text{OB}(\text{ipc})_2]_3$  ligand is smaller [ $\Delta\nu(\text{CO}) = 32 \text{ cm}^{-1}$ ] than the one seen between “electron rich” and “electron poor” arene ligands ( $70 \text{ cm}^{-1}$ ; *vide supra*). This span of  $\Delta\nu(\text{CO})$  is seen for all  $[\text{M}\{\text{P}_3\text{C}_3(\text{OX})_3\}(\text{CO})_3]$  complexes independently on the metal centre  $M = \text{Cr}, \text{Mo}, \text{W}$  (see the last three entries in every Tables 2–4).

The nature of the metal centre has also an influence on the  $\nu(\text{CO})$  stretching frequencies but this is rather small and for all type of ligands the largest step is seen between the  $\text{Cr}(0)$  and  $\text{Mo}(0)$  complexes (shift of about  $10 \text{ cm}^{-1}$  to higher wave numbers) while there is almost no change between  $\text{Mo}(0)$  and  $\text{W}(0)$  complexes.

That for every  $[\text{M}\{\text{P}_3\text{C}_3(\text{OH})_3\}(\text{CO})_3]$  (entry 14, entry 8, entry 7 in Tables 2, 3 and 4, respectively) or  $[\text{M}\{\text{P}_3\text{C}_3[\text{O}(\text{Si}^t\text{BuPh}_2)]_3\}(\text{CO})_3]$  complex ( $M = \text{Cr}, \text{Mo}, \text{W}$ ; entry 15, entry 9, entry 8 in Tables 2, 3 and 4, respectively) the  $A_1 \nu(\text{CO})$  is quite similar to the one observed in the corresponding parent benzene complex  $[\text{M}(\text{C}_6\text{H}_6)(\text{CO})_3]$  (entry 5 in Tables 2, 3 and 4, respectively) may be due to counterbalancing effect of the  $\pi$ -electron donating hydroxyl groups and  $\pi$ -electron withdrawing  $\text{P}_3\text{C}_3$  core.

$^{183}\text{W}$  satellites (natural abundance 14.3%)<sup>60</sup> were observed in the  $^{31}\text{P}\{^1\text{H}\}$  NMR spectra of the  $\text{W}(0)$  triphosphinine complexes prepared in this work. The  $J_{\text{WP}}$  couplings decrease very slightly from  $\text{W}-\text{P}_3\text{C}_3-\text{B}$  ( $21.4 \text{ Hz}$ ) to  $\text{W}-\text{P}_3\text{C}_3-\text{Si}$  ( $19.1 \text{ Hz}$ ) and  $\text{W}-\text{P}_3\text{C}_3-\text{H}$  ( $16.4 \text{ Hz}$ ). These data demonstrate that, as expected, a shorter  $\text{W}-\text{ct}$  distance correlates with a larger  $^{31}\text{P}-^{183}\text{W}$  coupling constant as a result of a stronger

$\pi$ -accepting ligand (see Fig. 1 and 2, in the ESI† for a correlation of  $\text{W}-\text{ct}$ ,  $J_{\text{WP}}$ ,  $\nu(\text{CO})$ , and  $\delta^{183}\text{W}$ ). No  $J_{\text{WP}}$  coupling has been reported for  $[\text{W}(\text{P}_3\text{C}_3^t\text{Bu}_3)(\text{CO})_3]$  and a  $\eta^6$ -coordinated monophosphinine tungsten complex has not been prepared yet. Only the  $J_{\text{WP}}$  coupling constants of  $\eta^1$ -phosphinine tungsten complexes are reported in the literature, as for example in the hexakis( $\eta^1$ -phosphinine)tungsten complex  $[\text{W}(\eta^1\text{-PC}_5\text{H}_5)_6]$ , which has a much larger direct  $\text{P}-\text{W}$  coupling of  $304 \text{ Hz}$ .<sup>61</sup>

Lastly,  $^{183}\text{W}$  resonances were measured using a triple resonance broad-band (TBI) probe with  $^1\text{H}/\text{D}/^{31}\text{P}/\text{BB}$  channels. A  $^{31}\text{P}-^{183}\text{W}$  heteronuclear multiple-quantum coherence (HMQC) experiment was performed, enhancing the sensitivity of the low  $\gamma^{183}\text{W}$  nucleus. The resonances in the  $^{183}\text{W}$  NMR spectrum shift very slightly to higher frequencies in the order  $\text{W}-\text{P}_3\text{C}_3-\text{B}$  ( $\delta^{183}\text{W} = -3146 \text{ ppm}$ ) <  $\text{W}-\text{P}_3\text{C}_3-\text{Si}$  ( $\delta^{183}\text{W} = -3140 \text{ ppm}$ ) <  $\text{W}-\text{P}_3\text{C}_3-\text{H}$  ( $\delta^{183}\text{W} = -3110 \text{ ppm}$ ). The number of reported  $^{183}\text{W}$  NMR chemical shifts for tungsten(0) complexes is rather limited and span from about  $-3500$  to  $-2500 \text{ ppm}$ . The most shielded  $^{183}\text{W}$  NMR shift is observed for  $[\text{W}(\text{CO})_6]$ , which shows a  $^{183}\text{W}$  NMR resonance at  $\delta^{183}\text{W} = -3446 \text{ ppm}$  and is chosen as reference compound here.<sup>62</sup> Replacement of three CO groups by a phosphite ligand like  $\text{P}(\text{OMe})_3$  (a strong  $\sigma$ -donor and medium  $\pi$ -acceptor) leads to a high frequency shift of about  $150 \text{ ppm}$  to  $\delta^{183}\text{W} = -3297 \text{ ppm}$  in  $[\text{W}\{\text{P}(\text{OMe})_3\}_3(\text{CO})_3]$ .<sup>63</sup> This deshielding effect becomes even more pronounced ( $630 \text{ ppm}$ ) in  $[\text{W}(\text{PMe}_2\text{Ph})_3(\text{CO})_3]$  ( $\delta^{183}\text{W} = -2818 \text{ ppm}$ ) where three strong  $\sigma$ -electron donor but relatively weak  $\pi$ -acceptor ligands were incorporated in the ligand sphere.<sup>64</sup> To the best of our knowledge, no  $\delta^{183}\text{W}$  NMR shifts are reported for any arene tungsten carbonyl complex. The shifts of the  $^{183}\text{W}$  resonances of  $[\text{W}\{\text{P}_3\text{C}_3(\text{OX})_3\}(\text{CO})_3]$  to higher frequencies by about  $300\text{--}336 \text{ ppm}$  places them in between  $[\text{W}(\text{CO})_6]$  and  $[\text{W}(\text{PMe}_2\text{Ph})_3(\text{CO})_3]$  indicating that to some extent the electronic effect on the  $^{183}\text{W}$  resonance caused by the replacement of three CO groups in facial position by one  $\text{P}_3\text{C}_3(\text{OX})_3$  ligand may be comparable to three  $\text{P}(\text{OMe})_3$  ligands.

## Conclusions

The successful preparation, and isolation of a  $3 \times 3$  matrix of complexes  $[\text{M}\{\text{P}_3\text{C}_3(\text{OX})_3\}(\text{CO})_3]$  using three different metals  $M = \text{Cr}, \text{Mo}, \text{W}$  and three different substituents  $X = \text{B}(\text{ipc})_2$ ,  $\text{Si}^t\text{BuPh}_2$ ,  $\text{H}$  allowed to systematically study and evaluate the effects provoked by these variations on a variety of spectroscopic data. The  $\eta^6$ -binding mode entails a large coordination shift of  $>160 \text{ ppm}$  to lower frequencies regardless the nature of the phosphinine. As has been observed in the coordination



chemistry of 2,4,6-*t*Bu-1,3,5-triphenylphosphine, the C–P–C angles are narrowed while the P–C–P angles are widened upon binding to a  $M(CO)_3$  fragment (note that the bulky and  $\sigma$ -electron donating *t*Bu groups lead to slightly larger C–P–C and more acute P–C–P angles in both the uncoordinated molecule as well as in the complexes).<sup>23</sup> As expected, the distances between the Cr centre and the centroid of the  $P_3C_3$  rings in all complexes are significantly smaller (by about 8%) than the Mo–Ct and W–ct distances, which are almost identical. This is attributed to the contracted d-orbital shell of Cr *versus* those of Mo and W.<sup>65,66</sup> This contraction may also be seen as reason for the fact that in all Cr(0) complexes listed in Table 2 independently of the type of ligand the  $\nu(CO)$  is somewhat smaller than in analogous Mo and W complexes because efficient  $M \rightarrow L$  electron back donation (which raises  $\nu(CO)$ ) requires an efficient orbital overlap.

As mentioned in the introduction, phosphinines are considered as especially good  $\pi$ -electron density accepting ligands. We recalculated the HOMO–LUMO gap of  $C_6H_6$  (6.60 eV),  $PC_5H_4$  (5.23 eV),  $1,3-P_2C_4H_4$  (4.88 eV), and  $1,3,5-P_3C_3H_3$  (4.63 eV) with density functional theory (DFT) methods (B3LYP/6-311+G\*\*) (see ref. 27 and the ESI† for details). While indeed there is a significant closing of the gap by  $\Delta E = 1.37$  eV when converting  $C_6H_6$  to  $PC_5H_5$ , additional substitution leads to much smaller changes (<0.4 eV). That  $P_3C_3(OH)_3$  has an especially small HOMO–LUMO gap of 4.27 eV is mainly due to the fact that the three hydroxyl groups in 1,3,5-position lift the HOMO in energy. Given these data, it is not surprising that the effect of replacing a monophosphinine by a triphosphinine has only a modest effect on the  $\nu(CO)$  stretching frequencies and thereby the electron density at the metal centre. The variation of the substituents in 1,3,5-position of an arene ligand or 2,4,6-position of a 1,3,5-triphenylphosphine allows to modify  $\nu(CO)$  over an range of about  $70\text{ cm}^{-1}$  or  $30\text{ cm}^{-1}$ , respectively, fully in accord with expectations (that is an electron donating ligand lowers  $\nu(CO)$ ; an electron accepting ligand raises  $\nu(CO)$ ). However, it remains surprising that in comparison to benzene complexes phosphinines provoke only modest (< $10\text{ cm}^{-1}$  when the parent molecules without strong electronic substituent effects are taken into account) blue-shifts of  $\nu(CO)$ . Under the assumption that these heterocycles are really “strong  $\pi$ -electron acceptor” ligands more pronounced deviations might have been expected. For example, we observed that the coordination of bis(imidazolium)-1,3-diphosphete-diide (IPD) – an unusual molecule with a central  $6\pi$ -aromatic  $C_2P_2$  ring of biradicaloid character – to form  $M(CO)_3$  complexes **A** (Fig. 2) in which  $\nu(CO)$  is shifted by about  $100\text{ cm}^{-1}$  to lower frequencies indicating strong electron donation to the metal centre by these cycles, indeed.<sup>67</sup> There is no question that phosphinines act as strongly binding ligands to metals (in their synthesis frequently an arene ligand in a precursor complex is displaced by a phosphinine). But we feel that the characterization of phosphinines as electron accepting ligands is an oversimplification and the electronic structure of phosphinine complexes is not fully understood yet. High-level quantum chemical methods should be applied to gain a better insight into electrostatic

interactions and the electron donating properties of phosphinines as well.

## Conflicts of interest

There are no conflicts to declare.

## References

- 1 G. Märkl, *Angew. Chem.*, 1966, **78**, 907–908.
- 2 A. J. Ashe, *J. Am. Chem. Soc.*, 1971, **93**, 3293–3295.
- 3 P. L. Floch, *Coord. Chem. Rev.*, 2006, **250**, 627–681.
- 4 P. Le Floch and F. Mathey, *Coord. Chem. Rev.*, 1998, **178–180**, 771–791.
- 5 N. Mezailles, F. Mathey and P. Le Floch, *Prog. Inorg. Chem.*, 2001, **49**, 455–550.
- 6 L. Weber, *Angew. Chem., Int. Ed.*, 2002, **41**, 563–572.
- 7 L. Kollar and G. Keglevich, *Chem. Rev.*, 2010, **110**, 4257–4302.
- 8 C. Müller, L. E. E. Broeckx, I. de Krom and J. J. M. Weemers, *Eur. J. Inorg. Chem.*, 2013, **2**, 187–202.
- 9 C. Müller and D. Vogt, *Dalton Trans.*, 2007, **47**, 5505–5523.
- 10 C. Müller and D. Vogt, *C. R. Chim.*, 2010, **13**, 1127–1143.
- 11 N. T. Coles, A. S. Abels, J. Leitz, R. Wolf, H. Grützmacher and C. Müller, *Coord. Chem. Rev.*, 2021, **433**, 213729–213756.
- 12 H. Heydt, *New Asp. Phosphorus Chem. II*, 223, 2003, vol. 223, pp. 215–249.
- 13 R. L. Falconer and C. A. Russell, *Coord. Chem. Rev.*, 2015, **297–298**, 146–167.
- 14 A. R. Barron and A. H. Cowley, *Angew. Chem., Int. Ed. Engl.*, 1987, **26**, 907–908.
- 15 R. Milczarek, W. Rüsseler, P. Binger, K. Jonas, K. Angermund, C. Krüger and M. Regitz, *Angew. Chem., Int. Ed. Engl.*, 1987, **26**, 908–909.
- 16 P. Binger, J. Stannek, B. Gabor, R. Mynott, J. Bruckmann, C. Krüger and S. Leininger, *Angew. Chem., Int. Ed. Engl.*, 1995, **34**, 2227–2230.
- 17 F. Tabellion, C. Peters, U. Fischbeck, M. Regitz and F. Preuss, *Chem. – Eur. J.*, 2000, **6**, 4558–4566.
- 18 P. L. Arnold, F. G. N. Cloke, P. B. Hitchcock and J. F. Nixon, *J. Am. Chem. Soc.*, 1996, **118**, 7630–7631.
- 19 P. Binger, S. Stutzmann, J. Stannek, B. Gabor and R. Mynott, *Eur. J. Inorg. Chem.*, 1999, **1**, 83–86.
- 20 C. W. Tate, P. B. Hitchcock, G. A. Lawless, Z. Benkő, L. Nyulászi and J. F. Nixon, *C. R. Chim.*, 2010, **13**, 1063–1072.
- 21 M. D. Francis, C. Holtel, C. Jones and R. P. Rose, *Organometallics*, 2005, **24**, 4216–4225.
- 22 K. Eggers, F. W. Heinemann, M. Hennemann, T. Clark, P. Binger and U. Zenneck, *C. R. Chim.*, 2010, **13**, 1203–1212.
- 23 S. B. Clendenning, J. C. Green and J. F. Nixon, *J. Chem. Soc., Dalton Trans.*, 2000, **9**, 1507–1512.



- 24 J. Waluk, H. P. Klein, A. J. Ashe and J. Michl, *Organometallics*, 1989, **8**, 2804–2808.
- 25 C. Elschenbroich, M. Nowotny, B. Metz, W. Massa, J. Graulich, K. Biehler and W. Sauer, *Angew. Chem., Int. Ed. Engl.*, 1991, **30**, 547–550.
- 26 R. Gleiter, H. Lange, P. Binger, J. Stannek, C. Krüger, J. Bruckmann, U. Zenneck and S. Kummer, *Eur. J. Inorg. Chem.*, 1998, **1998**, 1619–1621.
- 27 R. Suter, Y. Mei, M. Baker, Z. Benkő, Z. Li and H. Grützmacher, *Angew. Chem., Int. Ed.*, 2017, **56**, 1356–1360.
- 28 A. M. Cook and R. Huetter, *J. Agric. Food Chem.*, 1981, **29**, 1135–1143.
- 29 H. J. Sanders, *Chem. Eng. News*, 1981, **59**, 20–36.
- 30 Y. Yu, J. C. Yu, C.-Y. Chan, Y.-K. Che, J.-C. Zhao, L. Ding, W.-K. Ge and P.-K. Wong, *Appl. Catal., B*, 2005, **61**, 1–11.
- 31 P. Gijsman, R. Steenbakkers, C. Fürst and J. Kersjes, *Polym. Degrad. Stab.*, 2002, **78**, 219–224.
- 32 P. Zhao, M. Zhang, D. Wu and Y. Liu, *Korean J. Chem. Eng.*, 2013, **30**, 1687–1690.
- 33 D. Jiang, C. Sun, Y. Zhou, H. Wang, X. Yan, Q. He, J. Guo and Z. Guo, *Fibers Polym.*, 2015, **16**, 388–396.
- 34 K. P. Nair, V. Breedveld and M. Weck, *Macromolecules*, 2008, **41**, 3429–3438.
- 35 K. P. Nair, V. Breedveld and M. Weck, *Soft Matter*, 2011, **7**, 553–559.
- 36 B. Roy, P. Baire and A. K. Nandi, *RSC Adv.*, 2014, **4**, 1708–1734.
- 37 E. E. Simanek, M. Mammen, D. M. Gordon, D. Chin, J. P. Mathias, C. T. Seto and G. M. Whitesides, *Tetrahedron*, 1995, **51**, 607–619.
- 38 D. S. Lawrence, T. Jiang and M. Levett, *Chem. Rev.*, 1995, **95**, 2229–2260.
- 39 D. Philp and J. F. Stoddart, *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 1154–1196.
- 40 L. J. Prins, D. N. Reinhoudt and P. Timmerman, *Angew. Chem., Int. Ed.*, 2001, **40**, 2382–2426.
- 41 H. Fenniri, B. L. Deng and A. E. Ribbe, *J. Am. Chem. Soc.*, 2002, **124**, 11064–11072.
- 42 J. Deberitz and H. Nöth, *Chem. Ber.*, 1973, **106**, 2222–2226.
- 43 B. Cordero, V. Gomez, A. E. Platero-Prats, M. Reyes, J. Echeverria, E. Cremades, F. Barragan and S. Alvarez, *Dalton Trans.*, 2008, 2832–2838.
- 44 S. S. Batsanov, *Inorg. Mater.*, 2001, **37**, 871–885.
- 45 R. H. Crabtree, *The Organometallic Chemistry of the Transition Metals*, 2019, ch. 4, pp. 98–133.
- 46 G. Frenking and N. Frohlich, *Chem. Rev.*, 2000, **100**, 717–774.
- 47 G. Frenking, *J. Organomet. Chem.*, 2001, **635**, 9–23.
- 48 G. Frenking, *Coord. Chem. Rev.*, 2003, **238–239**, 55–82.
- 49 C. J. Breheny, J. M. Kelly, C. Long, S. O’Keeffe, M. T. Pryce, G. Russell and M. M. Walsh, *Organometallics*, 1998, **17**, 3690–3695.
- 50 J. A. S. Howell, P. C. Yates, N. F. Ashford, D. T. Dixon and R. Warren, *J. Chem. Soc., Dalton Trans.*, 1996, 3959–3966.
- 51 J. M. Jenkins, J. R. Moss and B. L. Shaw, *J. Chem. Soc. A*, 1969, 2796–2800.
- 52 B. L. Ross, J. G. Grasselli, W. M. Ritchey and H. D. Kaesz, *Inorg. Chem.*, 1963, **2**, 1023–1030.
- 53 R. D. Fischer, *Chem. Ber.*, 1960, **93**, 165–175.
- 54 G. Davidson and E. M. Riley, *J. Organomet. Chem.*, 1969, **19**, 101–114.
- 55 J. A. S. Howell, C. J. Beddows, P. J. O’Leary, P. C. Yates, P. McArdle, D. Cunningham and H. E. Gottlieb, *J. Organomet. Chem.*, 1997, **527**, 21–27.
- 56 Z. Zhao, J. Messinger, U. Schon, R. Wartchow and H. Butenschön, *Chem. Commun.*, 2006, 3007–3009.
- 57 A. D. Hunter, V. Mozol and S. D. Tsai, *Organometallics*, 1992, **11**, 2251–2262.
- 58 J. Deberitz and H. Nöth, *Chem. Ber.*, 1970, **103**, 2541–2547.
- 59 R. B. King and A. Fronzaglia, *Inorg. Chem.*, 1966, **5**, 1837–1846.
- 60 P. S. Pregosin, in *Encyclopedia of Magnetic Resonance*, 2011.
- 61 C. Elschenbroich, S. Voss, O. Schiemann, A. Lippek and K. Harms, *Organometallics*, 1998, **17**, 4417–4424.
- 62 R. L. Keiter and D. G. Vander Velde, *J. Organomet. Chem.*, 1983, **258**, C34–C36.
- 63 G. T. Andrews, I. J. Colquhoun, W. McFarlane and S. O. Grim, *J. Chem. Soc., Dalton Trans.*, 1982, **12**, 2353–2358.
- 64 H. C. E. McFarlane, W. McFarlane and D. S. Rycroft, *J. Chem. Soc., Dalton Trans.*, 1976, **1**, 1616–1622.
- 65 M. Kaupp, *J. Comput. Chem.*, 2007, **28**, 320–325.
- 66 F. Weinhold and C. R. Landis, *Valency and bonding: a natural bond orbital donor-acceptor perspective*, Cambridge University Press, 2005.
- 67 Z. Li, X. Chen, L. L. Liu, M. T. Scharnholtz and H. Grützmacher, *Angew. Chem., Int. Ed.*, 2020, **59**, 4288–4293.

