Dalton Transactions



COMMUNICATION

View Article Online



Cite this: *Dalton Trans.*, 2023, **52**, 10672

Received 4th July 2023, Accepted 19th July 2023 DOI: 10.1039/d3dt02094k

rsc.li/dalton

Interaction of germanium analogue of organic isonitrile with Cu(1) imide in side-on mode†

[NHC \rightarrow GeN(Ar)Cu₂NAr]₂, the formal adduct of germanium analogue of organic isonitrile [GeNAr] with Cu(1) imide [(Cu₂NAr)₂] (Ar = 2,6-*i*Pr₂C₆H₃) was prepared from the N-heterocyclic carbene (NHC) stabilized acyclic germylene Ge[N(H)Ar]₂ by reacting with two equivalents of *n*BuLi and CuCl(PPh₃)₃. As elucidated by X-ray crystal structural characterization, the two separated [GeNAr] moieties interacted with [(Cu₂NAr)₂] core in the side-on mode.

The synthesis of the heavier group 14 element analogues of organic isonitriles, in the form of MNR (M = Ge, Sn, Pb; R = organic group), is both structurally interesting and reactively attractive. Due to the large size of the M(II) elements as well as the weak π bonds, compared to the congeneric C, such heavier analogues mostly exist as oligomers $(MNR)_n$ (n = 2, 3, 4), rather than monomers like the organic isonitriles C=NR. For example, using the moderately sterically hindered diisopropyl phenyl ligand, Power and co-workers synthesized the trimeric cyclic (GeNR)₃ (R = 2,6-iPr₂C₆H₃), which contained a planar Ge₃N₃ six-membered ring.¹ Other instances included cubanes $(M_4N_4, M = Ge, ^2 Sn, ^{2-7} and Pb^5)$ or the double cubane (Sn₇N₈).8 However, germanium derivatives with lower aggregation, like dimers, which could involve multiple bonding of Ge (II) and N, are still rare. Limited available examples included the dimeric (GeNR)2, containing the Ge2N2 four-membered ring, supported by exceptionally bulky R ligands (R = 2,4,6- $2,4,6-tBu_3C_6H_2,^{10}$ $(CF_3)_3C_6H_2$, and Me₃C₆H₂)₂C₆H₃ ¹¹). In relevant reactivity studies, ¹⁰ the dimeric Ge(II) imido complexes (GeNR)2 were found to be reactive in oxidative addition or coordination to transition metals. So far, It is noteworthy that in the above reports of dimeric $(GeNR)_2$, 10,11 the key precursor $Ge[N(H)R]_2$ was specially mentioned, which was found to spontaneously generate dimeric $(GeNR)_2$ and release the primary amine H_2NR , as shown in the reaction eqn (1).

$$2Ge[N(H)R]_2 \longrightarrow (GeNR)_2 + 2H_2NR$$
 (1)

$$2Ge[N(M)R]_2 \longrightarrow 2[GeNR] + 2[M_2NR] \longrightarrow ?$$
 (2)

This gives us a hint of whether a similar reaction would occur if the amine proton H in (1) is replaced by the monovalent metal M(1) (as shown by reaction eqn (2))? If so, would the in situ generated [GeNR] moiety preferentially coordinate with the in situ formal $[M(1)_2NR]$ to bring about the possibility of the two [GeNR] species being split? For this end, we planned to carry out the reaction exploration outlined in eqn (2) using Cu(1) of group 11 metals as the costar metal element, which has a good affinity for both germanium and nitrogen.¹³ To make this methodology more representative and comparable, we proposed to use the common commercially available primary amine H₂NAr with large steric hindrance (typically $Ar = 2,6-iPr_2C_6H_3$) rather than extra bulky ones⁹⁻¹¹ to realize our synthetic intention. However, as known, it is more challenging to carry out the reaction using 2,6-diisopropyl aniline as the starting material due to two possible difficulties derived from the literature work. First, if the route of intermolecular deamination of primary amines is used, i.e., if the synthesis starts from $Ge[N(SiMe_3)_2]_2$ with H_2NAr (Ar = 2,6- $iPr_2C_6H_3$), the more stable trimeric (GeNAr)3 rather than the expected Ge[N (H)Ar₂ is obtained regardless of the ratio of reactants. Second, if the route of intermolecular elimination of salt is used, it could result in very low yields of impure products, as suggested by the previous unsuccessful attempt in which GeCl₂·dioxane

neither example of splitting the dimer (GeNR)₂ into separated GeNR species has been reported, nor the interaction of GeNR with metals through both Ge and N sites. So far, the interaction mode in the side-on bonded isonitrile complex [Mo₂(μ - η^2 -CNtBu)-(CO)₄Cp₂],¹² is known.

^aSchool of Chemistry and Chemical Engineering, Central South University, Lushannan Road 932, 410083 Changsha, China. E-mail: yangy@csu.edu.cn ^bInstitut für Anorganische Chemie, Universität Göttingen, Tammannstrasse 4, 37077 Göttingen, Germany. E-mail: hroesky@gwdg.de

[†] Electronic supplementary information (ESI) available: Experimental section, calculation details and outputs, characterisation data, NMR spectra, Cartesian coordinates, crystal data and refinement. CCDC 2201109-2201112. For ESI and crystallographic data in CIF or other electronic format see DOI: https://doi.org/10.1039/d3dt02094k

was treated with even greatly hindered LiN(H)Ar[#] (Ar[#] = 2,6- $(2,4,6\text{-Me}_3\text{C}_6\text{H}_2)_2\text{C}_6\text{H}_3^{-11}$). Thus, the two-electron-donor groups (D) was considered, which may offer additional steric protection and help the adduct D \rightarrow Ge(NHAr)₂ to be a much stronger Lewis base. Among such donors, N-heterocyclic carbene (NHC) has been applied in a number of previous studies on the synthesis and reactions of germylenes. ¹⁴⁻¹⁹ Therefore, our

work started with the synthesis of NHC \rightarrow Ge[N(H)Ar]₂.

Dalton Transactions

The primary amine ArNH₂ (Ar = 2,6-iPr₂C₆H₃) was treated with the equivalent molar of nBuLi to generate the lithiated product [LiN(H)Ar] (Scheme 1). The latter with or without separation was further reacted with GeCl₂·1,4-dioxane in the presence of NHC (:C[N(iPr)C(Me)]₂) to give the expected product NHC \rightarrow Ge[N(H)Ar]₂ (1, Scheme 1) as yellow solids in relatively high yield (80%). Alternatively, 1 can also be obtained in moderated yield (74%) by the treatment of GeCl₂·1,4-dioxane with equimolar [NHC \rightarrow LiN(H)Ar]₂ (2), which was formed by the addition of NHC to the solution of the intermediate [LiN(H) Ar].

The melting point of NHC \rightarrow Ge[N(H)Ar]₂ (1) is 135 °C, which is much lower than that of Ge[N(H)R]₂ (R = 2,6-(2,4,6-Me₃C₆H₂)₂C₆H₃) (198 °C).¹¹ In the FT-IR spectrum of 1, the strong adsorption at 3349 cm⁻¹ was attributed to NH stretching. The ¹H NMR spectrum of 1 exhibited resonances for amide protons observed at δ 5.98 ppm as a broad signal, while the septet at δ 3.51 ppm for the methine protons of isopropyl groups of Ar groups showed a correct integral ratio relative to that of NHC methine protons (2:1), consistent with the theoretically chemical formulation. The elemental analysis of 1 further confirmed the chemical compositions by comparing the measured elemental ratios with the theoretically calculated results.

The saturated solution of **1** in Et₂O was stored at room temperature for one day to produce yellow crystals suitable for single-crystal X-ray diffraction measurement. Compound **1** crystallizes in the triclinic space group $P\bar{1}$. Its molecular structure is depicted in Fig. 1. Due to the steric hindrance among NHC and amine ligands, the molecular structure of **1** is not symmetric. The pyramidal geometry of the Ge(II) centre was completed by two amide nitrogen atoms and one carbene carbon atom, where Ge(II) was 1.0267(13) Å away from the C(35)–N(1)–N(2) plane. The sum of bond angles around Ge(II) is $288.19(25)^{\circ}$, indicating the presence of the lone pair of electrons on the vertex Ge(II) atom. In contrast, the carbene C(35) atom is located in an almost trigonal planar environment, as revealed by the sum of the angles at C(35) (360.00(47)°). The

$$ArNH_{2} \xrightarrow{nBuLi} ArNHLi = \begin{cases} 1) \text{ NHC} \\ 2) \text{ GeCl}_{2} \\ 2:1:1 \\ -nBuH \end{cases}$$

$$Ar = \begin{cases} Pr \\ Pr \end{cases} \qquad NHC = \begin{cases} Pr \\ NHAr \end{cases}$$

$$NHC = \begin{cases} Pr \\ NHAr \end{cases}$$

$$NHC = \begin{cases} Pr \\ NHAr \end{cases}$$

Scheme 1 Synthesis of NHC \rightarrow Ge[N(H)Ar]₂ (1).




Fig. 1 Molecular structure of NHC \rightarrow Ge[N(H)Ar]₂ (1). Ellipsoids are drawn at 30% probability. H atoms except for NH groups have been omitted for clarity. Selected bond lengths (Å) and bond angles (°): Ge(1)–C(35) 2.144(2), Ge(1)–N(1) 1.951(2), Ge(1)–N(2) 1.950(2); N(1)–Ge(1)–N(2) 97.04(9), N(1)–Ge(1)–C(35) 89.75(8), N(2)–Ge(1)–C(35) 101.40(8).

Ge(1)–C(35) bond length is 2.144(2) Å, which is longer than the Ge–C single bond in general (\sim 2.04 Å).²⁰ (Fig. S1c†). The Ge–N bond lengths (1.950(2) and 1.951(2) Å) and N–Ge–N bond angle (97.04(9)°) of 1 are found to be larger than those of Ge[N (H)Ar $^{\#}$]₂ (1.896(2) Å), (88.6(2)°).¹¹

Interestingly, the Ge(II) atom of the acyclic compound 1 possesses a highly similar environment to that in a cyclic germylene L'Ge \leftarrow NHC (L' = HC[C(CH₂)N(Ar)]C(Me)N(Ar), Ar = 2,6-iPr₂C₆H₃; NHC = :C[N(Me)C(Me)]₂), in terms of the bond lengths and bond angles of CGeN₂ core with angle around Ge (II) (288.01°), Ge–C bond length (2.147(3) Å), Ge–N bond lengths (1.927(3) and 1.929(3) Å), and N–Ge–N bond angle (95.1(1)°). ¹⁵

To gain a deeper understanding of the electronic nature of 1, the quantum chemical calculations were performed using PBE1PBE/genecp (def2-TZVP basis set for Ge and def2SVP basis set for C, N, H) with Gaussian 16 program suite, ²¹ and analysed with Multiwfn. ²² The optimized structure of 1' is the global minimum and Cartesian coordinates were listed in Table S1 (ESI)† with a geometry similar to that determined experimentally. The frontier molecular orbitals, Ge(II)–C bond, and the interaction between carbene and germylene fragments were investigated. The HOMO and LUMO surfaces are depicted in Fig. 2.

The HOMO and LUMO energies are -4.73 and -0.30 eV, respectively, and the energy gap is 4.43 eV (426.99 kJ mol⁻¹), which falls in the common range for germylenes.²³

To investigate the electron transfer between the NHC and $[ArN(H)]_2$ Ge fragments, the extended charge decomposition analysis (ECDA) was performed, which suggested the 0.38 net electrons obtained by $Ge[N(H)Ar]_2$ from NHC (Fig. S11†).

The lithiation of 1 was carried out in situ and the subsequent transmetallation was performed with $CuCl(PPh_3)_3$ to

Communication Dalton Transactions





Fig. 2 HOMO (a) and LUMO (b) of NHC \rightarrow Ge[N(H)Ar]₂ (1') (H atoms are omitted for clarity). Optimized geometry: Ge(1)–C(35) 2.14101 Å, Ge(1)–N(1) 1.9537 Å, Ge(1)–N(2) 1.9467 Å; N(1)–Ge(1)–N(2) 97.2133°, N(1)–Ge (1)–C(35) 87.3829°, N(2)–Ge(1)–C(35) 100.3288°.



Scheme 2 Synthesis of $[NHC \rightarrow GeN(Ar)Cu_2NAr]_2$ (3) and the possible intermediates (3A and 3B).

afford the separation of 3 as dark yellow solid in 40% yield (Scheme 2).

NMR characterizations and elemental analysis have shown that compound 3 may preclude the presence of PPh₃ ligand and suggested a 2:1 composition of Ar and NHC moieties in it. As expected, X-ray single crystal diffraction proved this composition and further indicated the cleavage of Ge–N bonds and the concurrent formation Ge–Cu and N–Cu bonds (Fig. 3).

It is therefore proposed as illustrated in Scheme 2 that, the reaction may proceed four steps, *i.e.* lithiation, transmetallation (3A), dimerization (3B), and rearrangement, to give [NHC \rightarrow GeN(Ar)Cu₂NAr]₂ (3), as a unique adduct of (HNC \rightarrow GeNAr) with (Cu₂NAr)₂ in a 2:1 ratio.

As shown in Fig. 3, compound [NHC \rightarrow GeN(Ar)Cu₂NAr]₂ (3) crystallizes in the monoclinic $P2_1/n$ space group with an inversion center, in which Ge(1) centre is located in a four-coordinated environments by bonding to C(1), N(3) and two Cu atoms. The Cu-linked N atoms adopted two different coordinate environments, in which the amino N(3) is four-coordinated in tetrahedral geometry, while the imino N(4) is three-coordinated in trigonal planar geometry; each of them bonded to two Cu atoms with shorter average Cu-N bond lengths (1.867(4), 1.880(4) Å) for the former and longer




Fig. 3 The molecular structure of [NHC \rightarrow GeN(Ar)Cu₂NAr]₂ (3). Ellipsoids are drawn at 30% probability. H atoms have been omitted for clarity. Selected bond lengths (Å) and bond angles (°): Ge(1)–C(1) 1.910 (5), Ge(1)–N(3) 1.878(4), Ge(1)–Cu(1) 2.8909(8), Ge(1)–Cu(2) 2.6988(9), Cu(1)–N(3) 1.867(4), Cu(2)–N(3) 1.880(4), Cu(1)–N(4) 1.893(4), Cu(2)–N (4A) 1.927(4), Cu(1)–Cu(2) 2.6333(8), Cu(1)–Cu(2A) 2.7573(8); N(3)–Ge (1)–C(1) 175.68(19), Cu(1)–N(3)–Cu(2) 89.30(16), Cu(1)–N(4)–Cu(2A) 92.41(16); A=1 – X, –Y, 1 – Z.

averages for the latter (1.893(4), 1.926(4) Å). The Cu_4N_4 core is highly planar, with nearly right angles of Cu-Cu-Cu (88.75(2)° and 91.25(2)°) and Cu-N-Cu (89.3(2)° and 92.4(2)°). This is significantly different from the previous related examples, where either ligands used were homoleptic, 24,25 or the Cu₄N₄ plane was very distorted. 26 The Cu-Cu separations (2.6334(9) Å and 2.7572(7) Å) in 3 are in the common range, ²⁴⁻²⁶ while Ge-C (1.909(6) Å) and Ge–N (1.878(4) Å) bond lengths are greatly shortened compared to those in 1 (2.146(3) Å and 1.946(2)-1.950(2) Å, respectively), and can be seen to fall between the single and double Ge-C/N bonds. 27,28 Interestingly, Ge(1) and N(3) were found to be right on the NHC backbone plane (0.002 (9) Å and 0.078(15) Å deviation, respectively), and the C(1), Ge(1), and N(3) atoms were almost arranged in a straight line $(C(1)-Ge(1)-N(3) 175.6(2)^{\circ})$. Compound 3 represents the first example separating the GeNR monomers by interaction with metals in the side-on bonded isonitrile mode.

Using the density functional method PBE1PBE/genecp (def2-TZVP basis set for Ge and Cu and def2SVP basis set for C, N, H), the optimized structure 3' was obtained with constraints of heavy-atom positions. Calculation of the vibrational frequencies revealed no imaginary frequencies, indicating that this structure is at a minimum on the potential energy surface. The optimized Cartesian coordinates of 3' were listed in Table S2.† Fig. 4 presented the HOMO and LUMO surfaces of 3'.

The HOMO/LUMO energies 3' were -3.21 and -2.29 eV, respectively, leading to a HOMO-LUMO gap of 0.92 eV (89.14 kJ mol⁻¹). The Wiberg bond order of the Ge(π)-N (0.92) and Ge(π)-C (0.98) bonds (Fig. S13†) in 3' suggested the inter-

a) b)

Dalton Transactions

Fig. 4 HOMO (a) and LUMO (b) of $[NHC \rightarrow GeN(Ar)Cu_2NAr]_2$ (3') (H atoms are omitted for clarity).

mediate nature between single and double bonds, when compared with those in 1′ (0.56 and 0.62). The Ge(1)–N(3) bond is largely occupied (1.89e) (Fig. S14†), in which the Ge(π) centre uses sp^{0.78} hybrid orbital bonding to N (sp^{8.46}) with high p character. The lone pair of electrons (1.33e) on Ge(1) is found somewhat reduced, in a natural bond orbital of high p character (sp^{29.91}). The ECDA (notes in Fig. S15†) showed that each NHC gave 0.69 net electrons to the rest fragment of 3′, while each NHC \rightarrow GeN(Ar) moiety rendered 0.35 net electrons, evidencing the donor role of NHC in carbene-germylene tandem interaction.

In conclusion, we have synthesized and characterized the N-heterocyclic carbene stabilized acyclic germylene $Ge[N(H) Ar]_2$ (Ar = 2,6- $iPr_2C_6H_3$) (1). The introduction of an extra auxiliary carbene donor proved to be effective in decreasing the bulky demand of the Ar group. The reaction of lithiated 1 with $CuCl(PPh_3)_3$ underwent rearrangement to form $[NHC \rightarrow GeN(Ar)Cu_2NAr]_2$ (3), the formal adduct of germanium analogue of organic isonitrile GeNAr with Cu(I) imide Cu_2NAr . Further exploration of the reactivity and properties of compound 3 can be expected to lead to certain discoveries. In addition, acyclic germylene 1 can be considered as an important and versatile precursor for the preparation of the title product and possibly more other M(I), M(II) or M(III) derivatives to bring out interesting compounds.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was financially supported by the National Natural Science Foundation of China (21771194). We are grateful for resources from the High-Performance Computing Center of Central South University.

References

- 1 R. A. Bartlett and P. P. Power, *J. Am. Chem. Soc.*, 1990, **112**, 3660–3662.
- 2 W. J. Grigsby, T. Hascall, J. J. Ellison, M. M. Olmstead and P. P. Power, *Inorg. Chem.*, 1996, 35, 3254–3261.
- 3 M. Veith, Angew. Chem., Int. Ed. Engl., 1987, 26, 1-14.
- 4 R. E. Allan, M. A. Beswick, A. J. Edwards, M. A. Paver, M.-A. Rennie, P. R. Raithby and D. S. Wright, *J. Chem. Soc., Dalton Trans.*, 1995, 1991–1994.
- 5 H. Chen, R. A. Bartlett, H. V. R. Dias, M. M. Olmstead and P. P. Power, *Inorg. Chem.*, 1991, 30, 3390–3394.
- 6 M. Veith and O. Recktenwald, Z. Naturforsch., B: Chem. Sci., 1983, 38, 1054–1061.
- 7 M. Veith, M.-L. Sommer and D. Jäger, *Chem. Ber.*, 1979, 112, 2581–2587.
- 8 F. Benevelli, E. L. Doyle, E. A. Harron, N. Feeder, E. A. Quadrelli, D. Sáez and D. S. Wright, *Angew. Chem., Int. Ed.*, 2000, **39**, 1501–1503.
- 9 J.-T. Ahlemann, H. W. Roesky, R. Murugavel, E. Parisini, M. Noltemeyer, H.-G. Schmidt, O. Müller, R. Herbst-Irmer, L. N. Markovskii and Y. G. Shermilovich, *Chem. Ber.*, 1997, 130, 1113–1121.
- 10 P. B. Hitchcock, M. F. Lappert and A. J. Thorne, *J. Chem. Soc., Chem. Commun.*, 1990, 1587–1589.
- 11 W. A. Merrill, R. J. Wright, C. S. Stanciu, M. M. Olmstead, J. C. Fettinger and P. P. Power, *Inorg. Chem.*, 2010, 49, 7097–7105.
- 12 H. Adams, N. A. Bailey, C. Bannister, M. A. Faers, P. Fedorko, V. A. Osborn and M. J. Winter, *J. Chem. Soc., Dalton Trans.*, 1987, 341–348.
- 13 J. Hlina, H. Arp, M. Walewska, U. Flörke, K. Zangger, C. Marschner and J. Baumgartner, *Organometallics*, 2014, 33, 7069–7077.
- 14 J. A. Cabeza, P. Garcia-Alvarez and C. J. Laglera-Gandara, Eur. J. Inorg. Chem., 2020, 784–795.
- 15 S. Yao, Y. Xiong and M. Driess, *Chem. Commun.*, 2009, 6466–6468.
- 16 C. Jones, A. Sidiropoulos, N. Holzmann, G. Frenking and A. Stasch, *Chem. Commun.*, 2012, 48, 9855–9857.
- 17 N. Katir, D. Matioszek, S. Ladeira, J. Escudié and A. Castel, *Angew. Chem., Int. Ed.*, 2011, **50**, 5352–5355.
- 18 A. Jana, V. Huch and D. Scheschkewitz, *Angew. Chem., Int. Ed.*, 2013, **52**, 12179–12182.
- 19 Y. Xiong, S. Yao, G. Tan, S. Inoue and M. Driess, *J. Am. Chem. Soc.*, 2013, **135**, 5004–5007.
- 20 P. Jutzi, S. Keitemeyer, B. Neumann and H. G. Stammler, *Organometallics*, 1999, **18**, 4778–4784.
- 21 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb and J. R. Cheeseman, *Gaussian 16 Rev. C.01*, Wallingford, CT, 2016.
- 22 T. Lu and F. Chen, J. Comput. Chem., 2012, 33, 580-592.
- 23 L. Wang, Y. S. Lim, Y. Li, R. Ganguly and R. Kinjo, *Molecules*, 2016, 21, 990.
- 24 P. Reiß and D. Fenske, Z. Anorg. Allg. Chem., 2000, 626, 1317–1331.

25 M. Veith and K. L. Woll, *Chem. Ber.*, 1993, **126**, 2383-2388.

Communication

- 26 M. K. Davies, P. R. Raithby, M.-A. Rennie, A. Steiner and D. S. Wright, *J. Chem. Soc., Dalton Trans.*, 1995, 2707–2709.
- 27 S. Kundu, P. P. Samuel, A. Luebben, D. M. Andrada, G. Frenking, B. Dittrich and H. W. Roesky, *Dalton Trans.*, 2017, 46, 7947–7952.
- 28 M. J. Evans, F. M. Burke, P. M. Chapple and J. R. Fulton, *Inorg. Chem.*, 2021, **60**, 8293–8303.