

CORRECTION

[View Article Online](#)
[View Journal](#) | [View Issue](#)



Cite this: *Food Funct.*, 2023, **14**, 9947

Correction: Long-chain polyunsaturated fatty acids and extensively hydrolyzed casein-induced browning in a *Ucp-1* reporter mouse model of obesity

Liufeng Mao,^{a,b} Jiwen Lei,^{a,b} Marieke H. Schoemaker,^c Bingxiu Ma,^d Yan Zhong,^e Tim T. Lambers,^c Eric A. F. Van Tol,^c Yulai Zhou,^d Tao Nie^{*a,b,f} and Donghai Wu^{*g,h}

Correction for 'Long-chain polyunsaturated fatty acids and extensively hydrolyzed casein-induced browning in a *Ucp-1* reporter mouse model of obesity' by Liufeng Mao *et al.*, *Food Funct.*, 2018, **9**, 2362–2373, <https://doi.org/10.1039/C7FO01835E>.

DOI: 10.1039/d3fo90095a

rsc.li/food-function

The authors regret that there was an error in Fig. 3M where some images were duplicated. The corrected Fig. 3 is shown below.

^aCAS Key Laboratory of Regenerative Biology, Joint School of Life Sciences, Guangzhou Medical University, Guangzhou 511436, China

^bGuangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou 510530, China

^cMead Johnson Pediatric Nutrition Institute, Global R&D, Middenkampweg 2, 6545 CJ Nijmegen, The Netherlands

^dSchool of Pharmaceutical Sciences, Jilin University, Changchun, 130012 Jilin, China

^eMead Johnson Pediatric Nutrition Institute, Global R&D, 15th floor, East building of New Hualian Mansion, No. 755 Middle Huaihai Road, Shanghai 200020, China

^fCentral Laboratory of the First Affiliated Hospital of Jinan University, Guangzhou 510630, China. E-mail: nietaoly@gmail.com; Fax: +86-020-32015297;

Tel: +86-020-32015297

^gCAS Key Laboratory of Regenerative Biology, Joint School of Life Sciences, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou

510530, China

^hGuangzhou Medical University, Guangzhou 511436, China. E-mail: wu_donghai@gibh.ac.cn; Fax: +86-020-32015250; Tel: +86-020-32015250






Fig. 3 ARA + DHA, eHC or their combination enhanced *Ucp1* expression. (A) Representative luminescence images of *Ucp1*^{+/LUC} mice in response to control, AD, eHC and their combination CAD diets. (B–D) Quantification of luminescence for the dorsal, upper body and lower body view images in response to control, AD, eHC and their combination CAD diets. (E–G) Luciferase enzymatic activity in iBAT, iWAT and eWAT in response to control, AD, eHC and their combination CAD diets. (H–J) Relative RNA expression of UCP1 in iBAT, iWAT and eWAT in response to control, AD, eHC and their combination CAD diets. (K) Western blot analysis of UCP1 in iBAT and (L) the ratio of UCP1/β-actin expression as determined using NIH Image J software. (M) H&E staining and UCP1 immunohistochemistry in iBAT and iWAT and (N–O) relative cell area in iBAT and iWAT. H&E, hematoxylin and eosin; LUC, luciferase; UCP1, uncoupling protein 1. #*p* < 0.01 vs. LFD-CON, **p* < 0.05 vs. HFD-CON.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.