



Cite this: *New J. Chem.*, 2023,
47, 1576

DOI: 10.1039/d2nj90176e

rsc.li/njc

Correction: Visible light-triggered synthesis of oxime ethers using tetrabromomethane as a mediator

Meng Tu,^a Bo-Jiang Chen,^a Rukhsana,^a Wenying Cui,^a Wentao Fu,^a
Chengshuo Shen^{*a} and Yong-Miao Shen ^{*ab}


Correction for 'Visible light-triggered synthesis of oxime ethers using tetrabromomethane as a mediator'
by Meng Tu et al., *New J. Chem.*, 2022, **46**, 23276–23281, <https://doi.org/10.1039/D2NJ04615F>.

The authors regret that some substrates were missing from Table 2 in the original article. The correct Table 2 is shown below.
The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Department of Chemistry, Zhejiang Sci-Tech University, Hangzhou, Zhejiang 310018, P. R. China. E-mail: shenym@zstu.edu.cn, shenchengshuo@zstu.edu.cn

^b Key Laboratory of Excited-State Materials of Zhejiang Province, Zhejiang University, Hangzhou 310027, P. R. China



Table 2 Scope of various substrates for the reaction of oximes with tetrahydrofuran^{ab}^a Reaction conditions: **1** (0.1 mmol, 1 eq.), THF **2** (5.0 mmol, 50 eq.), CBr₄ (0.18 mmol, 1.8 eq.), BAC (2 mL), 6 W blue LED, argon, for 5 hours.^b Isolated yield. ^c **3ac**–**3ag** irradiated for 12 hours.