

## CORRECTION

[View Article Online](#)  
[View Journal](#) | [View Issue](#)

Cite this: *New J. Chem.*, 2023,  
47, 1576

DOI: 10.1039/d2nj90176e

rsc.li/njc

## Correction: Visible light-triggered synthesis of oxime ethers using tetrabromomethane as a mediator

Meng Tu,<sup>a</sup> Bo-Jiang Chen,<sup>a</sup> Rukhsana,<sup>a</sup> Wenying Cui,<sup>a</sup> Wentao Fu,<sup>a</sup> Chengshuo Shen<sup>\*a</sup> and Yong-Miao Shen<sup>ID \*ab</sup>


Correction for 'Visible light-triggered synthesis of oxime ethers using tetrabromomethane as a mediator' by Meng Tu et al., *New J. Chem.*, 2022, **46**, 23276–23281, <https://doi.org/10.1039/D2NJ04615F>.

The authors regret that some substrates were missing from Table 2 in the original article. The correct Table 2 is shown below.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

<sup>a</sup> Department of Chemistry, Zhejiang Sci-Tech University, Hangzhou, Zhejiang 310018, P. R. China. E-mail: shenym@zstu.edu.cn, shenchengshuo@zstu.edu.cn

<sup>b</sup> Key Laboratory of Excited-State Materials of Zhejiang Province, Zhejiang University, Hangzhou 310027, P. R. China

Table 2 Scope of various substrates for the reaction of oximes with tetrahydrofuran<sup>a,b</sup>

<sup>a</sup> Reaction conditions: **1** (0.1 mmol, 1 eq.), THF **2** (5.0 mmol, 50 eq.), CBr<sub>4</sub> (0.18 mmol, 1.8 eq.), BAC (2 mL), 6 W blue LED, argon, for 5 hours.

<sup>b</sup> Isolated yield. <sup>c</sup> 3ac–3ag irradiated for 12 hours.