ORGANIC CHEMISTRY

CCS 中国化学会 CHEMICAL SOCIETY





FRONTIERS

RESEARCH ARTICLE

View Article Online
View Journal | View Issue



Cite this: *Org. Chem. Front.*, 2023, **10**, 1738

Gold(1)-catalyzed cycloisomerization of alcohol or amine tethered-vinylidenecyclopropanes providing access to morpholine, piperazine or oxazepane derivatives: a carbene *versus* non-carbene process†

Jun-Sheng Wei, a Song Yang, a Yin Wei, b Sima Shamsaddinimotlagh, Hossein Tavakol* and Min Shi a ** sand Shamsaddinimotlagh, sand Shamsaddinimotla

A gold(i)-catalyzed intramolecular cyclization of alcohol or amine tethered-vinylidenecyclopropanes *via* a carbene or non-carbene process was developed to afford functionalized morpholines, piperazines and oxazepanes in good yields with a broad scope and excellent functional group tolerance under mild conditions. The steric bulkiness or chain length can modulate the reaction pathway. Substrates with a less sterically hindered group located at the allene moiety afford morpholines or piperazines containing an alkylidenecyclopropane *via* the non-carbene process, while sterically bulky ones give morpholines or piperazines containing a cyclobutene unit through the carbene process. Moreover, extending the carbon chain length of vinylidenecyclopropane enables the formation of seven-membered oxazepane *via* the carbene process. The synthetic utility of this protocol was also highlighted by its gram-scale synthesis and various transformations of the cyclization products.

Received 19th January 2023, Accepted 24th February 2023 DOI: 10.1039/d3qo00085k

rsc.li/frontiers-organic

Morpholine, piperazine and oxazepane derivatives have gained extensive attention due to their positive physicochemical, metabolic and biological properties. They are biologically active nitrogen-containing heterocycles involved in a variety of natural products and pharmaceuticals, such as the antibiotic drug finafloxacin, antianginal drug trimetazidine, anti-depressant agents reboxetine and befuraline, dopamine D4 receptor ligands, noradrenaline reuptake inhibitors and so on (Fig. 1). Although numerous methodologies have been established for the synthesis of these useful scaffolds, the construction of functionalized morpholine, piperazine or oxazepane derivatives still remains challenging.

Recently, gold(1)-catalyzed nucleophilic additions to C-C multiple bonds as a popular research branch of gold chemistry

[†] Electronic supplementary information (ESI) available: Experimental procedures and characterization data of new compounds. CCDC 1537103, 2178224, and 2178226. For ESI and crystallographic data in CIF or other electronic format see DOI: https://doi.org/10.1039/d3q000085k




Fig. 1 Selected biologically active compounds containing morpholine, piperazine and 1,4-oxazepane skeletons.

have made significant progress. Over the past two decades, considerable advancements have been made in gold(i)-catalyzed addition of X–H (X = O, N, C) bonds to unsaturated carbon–carbon bonds of olefins, 10 alkynes 11 and allenes. For instance, Widenhoefer $et\ al.$ reported the first example of gold(i)-catalyzed intramolecular hydroamination of unactivated alkenes, efficiently affording various tetrahydropyrrole and hexahydropiperidine skeletons (Scheme 1a). Later, Yamamoto $et\ al.$ reported the gold(i)-catalyzed intermolecular hydroamination between allenes and morpholine. The appropriate steric environment around gold catalysts plays a pivotal role in carrying out such aliphatic hydroamination. In addition,

^aKey Laboratory for Advanced Materials and Institute of Fine Chemicals, School of Chemistry & Molecular Engineering, East China University of Science and Technology, 130 Mei Long Road, Shanghai 200237, People's Republic of China ^bState Key Laboratory of Organometallic Chemistry, Center for Excellence in Molecular Synthesis, University of Chinese Academy of Sciences, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 345 Lingling Road, Shanghai 200032, China. E-mail: mshi@sioc.ac.cn


^cDepartment of Chemistry, Isfahan University of Technology, Isfahan 84156-83111, Iran. E-mail: h_tavakol@cc.iut.ac.ir

Scheme 1 Previous examples gold(ı)-catalyzed additions to C-C multiple bonds

different from monosubstituted allenes (Scheme 1b), 1,3-disubstituted allenes usually gave a regioisomeric mixture and its ratio also depends on the steric effects (Scheme 1c). Moreover, an analogous cyclization reaction of alkynes has also been developed by Plé's group recently, 11a giving N- and O-spirocycles in good to excellent yields (Scheme 1d).

In the meantime, it should be mentioned that catalytic transformations involving gold carbenes as key intermediates are also considered as some of the most important aspects of homogeneous gold catalysis.¹³ To enrich the chemistry of gold carbene, a gold(1)-catalyzed cycloisomerization or allyl transfer of vinylidenecyclopropane-ene derivatives via controllable carbene or non-carbene processes has been explored in our laboratory, 14 providing a new synthetic route for the construction of O-bearing heterocyclic scaffolds containing a fused tricyclic system and fused five- and six-membered ring systems. Substituents adjacent to the oxygen atom on the aryl groups could switch the reaction mode (Scheme 2a).

Therefore, vinylidenecyclopropanes (VDCPs) can be realized as excellent candidates for gold(1)-catalyzed cycloisomerization due to their multiple reaction sites. 14,15 During our ongoing efforts on gold-catalyzed cycloisomerization of VDCPs, we subsequently designed and synthesized a series of VDCPs tethered with an alcohol or amine moiety to explore their reaction outgold(1)-catalyzed intramolecular cyclization (Scheme 2b, this work). Interestingly, we found that the cyclization could proceed through a carbene or a non-carbene process as well depending on the substrate's steric effect or the connected carbon chain length, affording various morpholine, piperazine or oxazepane derivatives in good yields under mild conditions. Herein, we wish to report the details in this context.



Scheme 2 Our previous work of gold(ı)-catalyzed cycloisomerization of vinylidenecyclopropane-enes via a carbene or non-carbene process

At the beginning of our investigation, VDCP 1a was selected as the model substrate to optimize the reaction conditions of this gold(1)-catalyzed intramolecular cyclization. As shown in Table 1, a series of gold catalysts have been screened using AgNTf₂ as a silver salt additive and a pre-prepared JohnPhosAu (MeCN)SbF₆ complex upon heating in chlorobenzene at 90 °C (Table 1, entries 1-8), and it was found that the product 2a was obtained in poor to moderate yields with complete consumption of substrate 1a, and XphosAuCl was identified as the best gold(i) catalyst in this reaction, affording 2a in 43% yield. Lowering the reaction temperature to 50 °C gave 2a in 60% yield (Table 1, entry 9) and then, we found that by carrying out the reaction at room temperature and prolonging the reaction time, the yield of 2a could be improved to 76% (Table 1, entry 10). The examination of the solvent effect with various commonly used solvents revealed that THF was the best solvent, giving 2a in 95% NMR yield along with 87% isolated yield (Table 1, entries 11-16). During the screening of solvents, it is surmised that the use of acetonitrile as the solvent resulted in essentially complete recovery of the starting material without observing the formation of the product presumably due to its coordination with the monovalent gold(1) catalyst (Table 1, entry 12). Notably, compared to the *in situ* generation of the cationic gold catalyst, using cationic gold catalyst XPhosAuNTf₂ slightly reduced the yield of 2a (Table 1, entry 17). Furthermore, the effect of silver salts on the reaction outcome was explored, identifying that AgNTf2 led to a better result than other silver additives such as AgBF4, AgOTf, and AgSbF₆ (Table 1, entries 18-20). No conversion was observed under the same conditions when the silver salt additive was replaced with NaBArF (Table 1, entry 21). As a consequence, alkylidenecyclopropane functionalized morpholine derivative 2a was obtained in 87% isolated yield with excellent regioselectivity using 5 mol% XPhosAuCl as the catalyst and 5 mol% AgNTf₂ as the silver salt additive in THF at room temperature under an ambient atmosphere, which serve as the


Table 1 Optimization of the reaction conditions for the synthesis of 2a

Entry ^a	Catalyst	Time (h)	Solvent	Temp. (°C)	Yield ^b (%)
1	IPrAuCl, AgNTf ₂	12	PhCl	90	Trace
2	(4-CF ₃ Ph) ₃ PAuCl, AgNT ₂	12	PhCl	90	Trace
3	ⁱ Pr ₃ PAuCl, AgNTf ₂	12	PhCl	90	Trace
4	JohnPhosAuCl, AgNTf ₂	12	PhCl	90	11
5	PPh ₃ AuCl, AgNTf ₂	12	PhCl	90	35
6	^t Bu ₃ PAuCl, AgNTf ₂	12	PhCl	90	12
7	JohnPhosAu(MeCN)SbF ₆	12	PhCl	90	38
8	XphosAuCl, AgNTf ₂	12	PhCl	90	43
9	XphosAuCl, AgNTf ₂	12	PhCl	50	60
10	XphosAuCl, AgNTf ₂	16	PhCl	rt	76
11	XphosAuCl, AgNTf ₂	16	1,4-Dioxane	rt	54
12	XphosAuCl, AgNTf ₂	16	CH_3CN	rt	Trace
13	XphosAuCl, AgNTf ₂	16	DCE	rt	40
14	XphosAuCl, AgNTf ₂	16	THF	rt	95 (87) ^c
15	XphosAuCl, AgNTf ₂	16	DCM	rt	46
16	XphosAuCl, AgNTf ₂	16	Toluene	rt	68
17	XphosAuNTf ₂	16	THF	rt	82
18	XphosAuCl, AgBF ₄	16	THF	rt	50
19	XphosAuCl, AgOTf	16	THF	rt	83
20	XphosAuCl, AgSbF ₆	16	THF	rt	86
21	XphosAuCl, NaBArF	16	THF	rt	Trace

^aThe reaction scale is 0.1 mmol of 1a in untreated solvent (0.04 M). ^b Yield was determined from ¹H NMR spectroscopic data using 1,3,5trimethoxybenzene as an internal standard. ^c Isolated yield.

optimal conditions for the production of 2a. The control experiments illustrated that silver salts such as AgNTf₂, Brønsted acid HNTf₂, and XPhosAuCl did not catalyze the reaction at all, indicating that this cyclization is indeed catalyzed by the gold(1) complex (see S8 in the ESI†). The crystal structure of 2a was unequivocally determined by X-ray diffraction as shown in Table 1, and the ORTEP drawing and its CIF data are presented in the ESI.†

Next, we synthesized a variety of alcohol or amine tethered vinylidenecyclopropanes with different R groups located at allene moieties to explore the generality of this cyclization under the optimal conditions and the results are shown in Scheme 3. For alcohol tethered VDCPs 1b-1g, in which the R group was a linear alkyl substituent, an alkyl chain with a terminal protected aldehyde group, a homoallyl group or a benzyl group, the reaction was tolerated, affording the corresponding products 2b-2g in 58-87% yields along with a trace amount of inseparable cyclized compound 2' derived from a gold-carbene induced cyclization. Remarkably, in the case of VDCP 1h bearing a cyclopropane at the allene moiety, the desired product 2h was formed in 74% yield as a single isomer perhaps due to the less steric hindrance of the cyclopropyl ring. Besides N-Ts protected VCDPs, N-phenylsulfonyl pro-




Scheme 3 Substrate scope for the non-carbene process. Unless otherwise specified, all reactions were carried out using 1 (0.2 mmol), XPhosAuCl (5 mol%) and AgNTf₂ (5 mol%) in THF (4.0 mL) at rt. Isolated yields. ^a Inseparable structural isomers of 2 and 2' were formed and the ratio was determined on the basis of ¹H NMR spectroscopic data.

tected substrate 1i and N-Bs protected substrate 1j also performed very well, affording the desired products 2i and 2j in 98% and 87% yields, respectively. Substrates 1k and 1l connected by an oxygen atom or a carbon atom were also compatible, giving the corresponding products 2k and 2l in good yields ranging from 90% to 99% under the standard conditions. We were delighted to find that by replacing primary alcohol with a secondary alcohol, the substrate 1m delivered the desired product 2m in 92% yield as a single diastereoisomer and its stereochemical configuration was determined by ¹H NMR NOESY spectroscopic analysis. Moreover, amine tethered vinylidenecyclopropanes 1n and 1o gave the desired functionalized piperazines 2n and 2o as single isomers in 77% and 85% yields, respectively, illustrating that this reaction has excellent functional group tolerance. Interestingly, the use of VDCP 1p as the substrate, which had a cyclobutane at the allene moiety, resulted in an inseparable mixture of 2p and 2p', which contains a cyclobutene moiety, in an 8:5 ratio as determined by ¹H NMR analysis presumably due to its steric effect. Notably, when replacing the benzylamine unit with a tosylated amine group such as VDCP 1z, no corresponding product could be obtained, perhaps due to its weaker nucleophilicity. In addition, aryl-substituted VDCPs 1aa-1ac were not suitable substrates for this gold-catalyzed transformation, presumably because the aryl substituents on the allene moiety make them difficult to polarize by cationic gold species in our reaction system. Substrate 1ad also failed to give the corresponding products under the standard conditions.

In order to gain a better understanding of the reaction processes, several sterically bulky R groups were introduced into the substrates, and we found that all of them could give the cyclobutene containing cyclized products 2q'-2t' as major products in 56-86% yields (Scheme 4). The crystal structure of 2q' was unambiguously determined by X-ray diffraction. The ORTEP drawing is given in Scheme 4 and its CIF data are also summarized in the ESI.† Particularly, different from VDCP 1p, cyclobutyl containing amine tethered vinylidenecyclopropane 1v furnished the desired product 2v' as a sole product in 63% yield via a gold carbene process, indicating that the tethered nucleophile could also significantly affect the reaction pathway. To our surprise, further investigations revealed that our reaction system enables the formation of seven-membered 1,4-oxazepanes 2w'-2y' in 67-88% yields as a sole product through a gold carbene process, indicating that the connected alkyl chain length of the substrate can modulate the reaction process.

In order to elucidate the impact of substrate steric bulkiness on the selectivity of reaction pathways, substrates 1a and 1q were selected to investigate the energy differences of several gold-VDCP intermediates since the A-value16 of the methyl group is 1.8 kcal mol⁻¹ and that of the isopropyl group is above 3.3 kcal $\text{mol}^{-1.16d}$ The relative free energies of several key intermediates are shown in Fig. 2 (for details, see S135 in the ESI†). For substrate 1a, the gold catalyst activates the allene double-bond away from the cyclopropyl unit, which is favored by 0.96 kcal mol⁻¹, suggesting that the non-carbene process is a dominant process for groups with less steric hindrance. For substrate 1q, the gold species activates the allene double-bond connected to the cyclopropyl unit, which is favored by 0.86 kcal mol⁻¹, suggesting that the carbene process becomes a dominant process for groups with more steric hindrance. These results are in good agreement with our experimental findings (Schemes 3 and 4).



Scheme 4 Substrate scope for the carbene process. Unless otherwise specified, all reactions were carried out using 1 (0.2 mmol), XPhosAuCl (5 mol%) and AgNTf2 (5 mol%) in THF (4.0 mL) at rt. Isolated yields. ^a Inseparable structural isomers of 2' and 2 were formed and the ratios were determined on the basis of ¹H NMR spectroscopic data.

Relative free energy
$$\Delta G_{298, THF}$$
 (kcal mol⁻¹)

Relative free energy $\Delta G_{298, THF}$ (kcal mol⁻¹)


Relative free energy $\Delta G_{298, THF}$ (kcal mol⁻¹)

Int A-1q, 0.86

Int C-1q, 0.00

Fig. 2 The relative free energies of key intermediates.


On the basis of the previous literature 12e,14,17 and our above results, a plausible mechanism for the present cyclization reaction is depicted in Scheme 5. Under the conditions of gold(1) catalysis, substrate 1 can undergo two different reaction pathways (carbene and non-carbene processes) depending on the steric hindrance of the R groups or the connected alkyl chain length. As for the non-carbene process, the π -activation of the double bond in the allene moiety with the in situ generated gold(1) complex takes place to furnish intermediate A, which triggers a 6-exo-trig cyclization through intramolecular nucleophilic addition by the hydroxyl group onto the electrophilic carbon center to afford the alkenyl gold intermediate B. The protodeauration of intermediate B produces the corresponding morpholine derivative 2 along with the regeneration of the Au (1) catalytic species for the catalytic cycle. In this pathway, the sterically less hindered R groups in substrate 1 may result in a lower transition state energy barrier of 6-exo cyclization, giving the products 2 as the major ones. However, the carbene and non-carbene pathways are competing cyclization processes in these reactions. Sterically more bulky groups at substrate 1's allene moiety may delay the direct 6-exo cyclization since the transition energy barrier is quite high and the carbene process




Scheme 5 Proposed reaction mechanisms.

becomes the main cyclization process, affording products 2' as the major ones. The same reason can be applied for the prolonged alkyl chain since the direct cyclization will become a slow process and the carbene process will dominate the reaction pathway. Thus, coordination of the gold(1) complex with vinylidenecyclopropane 1 bearing a sterically bulky group gives intermediate C, which initiates a ring expansion to afford the gold carbenoid intermediate D. The intramolecular nucleophilic attack of the hydroxyl group onto the electrophilic carbon center in intermediate D and release of a proton furnish intermediate E, which undergoes protodeauration to produce the cyclized product 2' and regenerates the gold catalyst. More comprehensive investigations on the mechanistic manipulation of carbene and non-carbene processes and the related density functional theory (DFT) calculations will be published in due course.

To illustrate the synthetic utility of the obtained product, a gram-scale reaction of 1a, with 2.15 g (7.0 mmol), was performed and the reaction proceeded smoothly, delivering 1.8 g of 2a in 83% yield (Scheme 6). Then, a variety of transformations of 2a were conducted under different conditions (Scheme 6). Initially, treatment of 2a with HCl aqueous solution (4.0 M in dioxane) gave a chlorine-substituted vinylcyclopropane tethered alcohol, which could be further transformed to compound 3 in 78% overall yield in the presence of p-nitrobenzoyl chloride, and its structure has been clarified by X-ray diffraction as shown in Scheme 6. Its CIF data are also presented in the ESI.† Next, iodine-substituted vinylcyclopropane tethered alcohol 4 was formed in 67% yield upon treating 2a with 1.0 equiv. of BF₃·Et₂O and 1.5 equiv. of Bu₄NI in DCM at



Scheme 6 Gram-scale synthesis and transformations of the obtained products 2a and 2q'. All of the transformations were carried out using 0.1 mmol of 2a or 2q'. (a) XPhosAuCl (5 mol%), AgNTf2 (5 mol%), THF, rt. (b) (1) HCl (4 M in dioxane), 80 °C; (2) DMAP (0.2 equiv.), DIPEA (3.0 equiv.), p-nitrobenzoyl chloride (2.5 equiv.), DCM, 0 °C. (c) Bu₄NI (1.5 equiv.), BF₃·Et₂O (1.0 equiv.), DCM, rt. (d) Pd(OAc)₂ (0.1 equiv.), Bu₄NI (1.0 equiv.), PhI(OAc)₂ (3.0 equiv.), CH₃CN, 60 °C. (e) m-CPBA (1.2 equiv.), DCM, rt. (f) m-CPBA (1.2 equiv.), DCM, rt.



Scheme 7 Asymmetric versions of the two processes for the production of 2a and 2a'.

room temperature. Considering that product 2a contained an alkylidenecyclopropane subunit, treatment of 2a with Pd(OAc)2 and PhI(OAc)2 afforded the corresponding diacetoxylated product 5 in 42% yield through a palladium-catalyzed ringopening reaction according to our previous work.18 Furthermore, epoxidation of 2a and 2q' with m-CPBA in DCM successfully gave the desired epoxide 6 in 68% yield as a 3:1 diastereomeric mixture and the epoxide 7 in 94% yield as a single diastereomer (Scheme 6).

Asymmetric variants of VDCPs 1a and 1q were also reacted with the chiral gold catalyst (R)-DTBM-SegPhosAuCl combined with AgNTf₂ in DCM at room temperature (for details, see S8 in the ESI†). As shown in Scheme 7, we found that treatment of 1a with 5 mol% (R)-DTBM-SegPhosAuCl and 10 mol% AgNTf2 afforded 2a in 78% yield along with 55% ee value. For substrate 1q, adding 5 mol% (R)-DTBM-SegPhosAuCl and 5 mol% AgNTf2 gave the desired product 2q' in 88% yield along with 72% ee value.

In conclusion, we have developed an efficient and novel synthetic protocol for the construction of morpholine, piperazine and oxazepane derivatives bearing an alkylidenecyclopropane or a cyclobutene moiety in good yields under mild conditions through a gold(1)-catalyzed intramolecular cycloisomerization of alcohol and amine tethered vinylidenecyclopropanes via a carbene or non-carbene process with a broad substrate scope. The steric bulkiness of the substituent at the allene moiety of vinylidenecyclopropanes and the chain length played key roles in the reaction pathway. In general, sterically bulky substituents and the prolonged alkyl chain prefer the gold(1)-catalyzed cyclization via a carbene process. In addition, a variety of transformations of the obtained product 2 have been performed to demonstrate their synthetic utility. Further investigations on the mechanistic paradigm of this gold(1)-catalyzed cyclization and the preparation of biologically active heterocycles are underway in our laboratory.

Author contributions

Shi, M. and Tavakol, H. directed the project and revised the manuscript. Wei, J. S., Shamsaddinimotlagh, S. and Yang, S.

wrote the manuscript and carried out the reactions. Wei, Y. checked the spectroscopic data and revised the manuscript.

Conflicts of interest

There are no conflicts of interest to declare.

Acknowledgements

We are grateful for the financial support from the National Key R & D Program of China (2022YFC2303100), the National Natural Science Foundation of China (21372250, 21121062, 21302203, 20732008, 21772037, 21772226, 21861132014, 91956115, and 22171078), and the Fundamental Research Funds for the Central Universities 222201717003.

Notes and references

- 1 (a) F. Arshad, M. F. Khan, W. Akhtar, M. M. Alam, L. M. Nainwal, S. K. Kaushik, M. Akhter, S. Parvez, S. M. Hasan and M. Shaquiquzzaman, Revealing Quinquennial Anticancer Journey of Morpholine: A SAR Based Review, Eur. J. Med. Chem., 2019, 167, 324-356; (b) V. Hahn, A. Mikolasch, K. Wende, H. Bartrow, U. Lindequist and F. Schauer, Synthesis of Model Morpholine Derivatives with Biological Activities by Laccase-Catalysed Reactions, Biotechnol. Appl. Biochem., 2009, 54, 187-195; (c) A. P. Kourounakis, D. Xanthopoulos and A. Tzara, Morpholine as a Privileged Structure: A Review on the Medicinal Chemistry and Pharmacological Activity of Morpholine Containing Bioactive Molecules, Med. Res. Rev., 2020, 40, 709-752; (d) E. Lenci, L. Calugi and A. Trabocchi, Occurrence of Morpholine in Central Nervous System Drug Discovery, ACS Chem. Neurosci., 2021, **12**, 378–390; (e) M. Taha and A. Wadood, Synthesis and Molecular Docking Study of Piperazine Derivatives as Potent Urease Inhibitors, Bioorg. Chem., 2018, 78, 411-417; (f) E. Vitaku, D. T. Smith and J. T. Njardarson, Analysis of the Structural Diversity, Substitution Patterns, and Frequency of Nitrogen Heterocycles among U.S. FDA Approved Pharmaceuticals, J. Med. Chem., 2014, 57, 10257-10274.
- 2 S. Lemaire, F. Van Bambeke and P. M. Tulkens, Activity of Finafloxacin, a Novel Fluoroquinolone with Increased Activity at Acid PH, towards Extracellular and Intracellular Staphylococcus Aureus, Listeria Monocytogenes and Legionella Pneumophila, *Int. J. Antimicrob. Agents*, 2011, 38, 52–59.
- 3 T. Kalai, M. Khan, M. Balog, V. K. Kutala, P. Kuppusamy and K. Hideg, Structure–Activity Studies on the Protection of Trimetazidine Derivatives Modified with Nitroxides and their Precursors from Myocardial Ischemia-Reperfusion Injury, *Bioorg. Med. Chem.*, 2006, 14, 5510–5516.

- 4 H. R. Arias, N. B. Fedorov, L. C. Benson, P. M. Lippiello, G. J. Gatto, D. Feuerbach and M. O. Ortells, Functional and Structural Interaction of (–)-Reboxetine with the Human α4β2 Nicotinic Acetylcholine Receptor, *J. Pharmacol. Exp. Ther.*, 2013, 344, 113–123.
- 5 D. Dauzonne, J. M. Gillardin, F. Lepage, R. Pointet, S. Risse, G. Lamotte and P. Demerseman, Synthesis and Some CNS Activities of New Benzofuranylacryloylpiperazines, *Eur. J. Med. Chem.*, 1995, 30, 53–59.
- 6 K. Audouze, E. O. Nielsen and D. Peters, New Series of Morpholine and 1,4-Oxazepane Derivatives as Dopamine D₄ Receptor Ligands: Synthesis and 3D-QSAR Model, J. Med. Chem., 2004, 47, 3089–3104.
- 7 T. Yukawa, Y. Nakada, N. Sakauchi, T. Kamei, M. Yamada, Y. Ohba, I. Fujimori, H. Ueno, M. Takiguchi, M. Kuno, I. Kamo, H. Nakagawa, Y. Fujioka, T. Igari, Y. Ishichi and T. Tsukamoto, Design, Synthesis, and Biological Evaluation of a Novel Series of Peripheral-Selective Noradrenaline Reuptake Inhibitors Part 3, *Bioorg. Med. Chem.*, 2016, 24, 3716–3726.
- 8 (a) S. J. Gharpure, D. Anuradha, J. V. K. Prasad and P. Srinivasa Rao, Stereoselective Synthesis of cis-2,6-Disubstituted Morpholines and 1,4-Oxathianes Intramolecular Reductive Etherification of 1,5-Diketones, Eur. J. Org. Chem., 2015, 86-90; (b) J. V. Matlock, T. D. Svejstrup, P. Songara, S. Overington, E. M. McGarrigle and V. K. Aggarwal, Synthesis of 6- and 7-Membered N-Heterocycles Using α -Phenylvinylsulfonium Salts, *Org.* Lett., 2015, 17, 5044-5047; (c) J. Kishore Vandavasi, W.-P. Hu, G. Chandru Senadi, H.-T. Chen, H.-Y. Chen, K.-C. Hsieh and J.-J. Wang, Aryl λ^3 -Iodane-Mediated 6-exotrig Cyclization to Synthesize Highly Substituted Chiral Morpholines, Adv. Synth. Catal., 2015, 357, 2788-2794; (d) B. Ritzen, S. Hoekman, E. D. Verdasco, F. L. van Delft and F. P. Rutjes, Enantioselective Chemoenzymatic Synthesis of *cis*- and *trans*-2,5-Disubstituted Morpholines, J. Org. Chem., 2010, 75, 3461-3464; (e) P. Maity and B. Konig, Synthesis and Structure of 1,4-Dipiperazino Benzenes: Chiral Terphenyl-type Peptide Helix Mimetics, Org. Lett., 2008, 10, 1473-1476; (f) K. H. Rohde, H. A. Michaels and A. Nefzi, Synthesis and Antitubercular Activity of 1,2,4-Trisubstitued Piperazines, Bioorg. Med. Chem. Lett., 2016, 26, 2206–2209; (g) G. J. Mercer and M. S. Sigman, Diastereoselective Synthesis of Piperazines by Manganese-Mediated Reductive Cyclization, Org. Lett., 2003, 5, 1591–1594; (h) F. Hafeez, A. Mansha, A. F. Zahoor, K. G. Ali, S. G. Khan and S. A. R. Naqvi, Facile Green Approach towards the Synthesis of Some Phenyl Piperazine Based Dithiocarbamates as Potent Hemolytic and Thrombolytic Agents, Pak. J. Pharm. Sci., 2021, 34, 1885-1890; (i) M. Taha and A. Wadood, Synthesis and Molecular Docking Study of Piperazine Derivatives as Potent Urease Inhibitors, Bioorg. Chem., 2018, 78, 411-417.
- 9 (a) A. S. K. Hashmi, Gold-Catalyzed Organic Reactions, *Chem. Rev.*, 2007, **107**, 3180–3211; (b) N. D. Shapiro and F. D. Toste, A Reactivity-Driven Approach to the Discovery

and Development of Gold-Catalyzed Organic Reactions, Synlett, 2010, 675-691, DOI: 10.1055/s-0029-1219369; (c) A. S. K. Hashmi, Gold-Catalyzed Organic Reactions, in Inventing Reactions, ed. L. J. Gooßen, 2013, vol. 44, pp. 143-164; (d) H. Ohno, Gold-Catalyzed Cascade Reactions of Alkynes for Construction of Polycyclic Compounds, Isr. J. Chem., 2013, 53, 869-882; (e) C. Shu, L. Li, T.-D. Tan, D.-Q. Yuan and L.-W. Ye, Ring Strain Strategy for the Control of Regioselectivity. Gold-Catalyzed Anti-Tandem Markovnikov Cycloisomerization Initiated Reactions of Alkynes, Sci. Bull., 2017, 62, 352-357; (f) H. Xiong, H. Wang, L. He, Y. Zhang and Z. Zeng, Application of Gold Catalyst in Nucleophilic Addition of Allene, Chin. J. Org. Chem., 2011, 31, 466-479.

Research Article

- 10 (a) C. F. Bender and R. A. Widenhoefer, Gold(I)-Catalyzed Intramolecular Hydroamination of Unactivated C-C Bonds with Alkyl Ammonium Salt, Chem. Commun., 2008, 2741-2743, DOI: 10.1039/b804081h; (b) J. L. Zhang, C. G. Yang and C. He, Gold(I)-Catalyzed Intra- and Intermolecular Hydroamination of Unactivated Olefin, J. Am. Chem. Soc., 2006, 128, 1798-1799; (c) X.-Y. Liu, C.-H. Li and C.-M. Che, Phosphine Gold(I)-Catalyzed Hydroamination of Alkenes under Thermal and Microwave-Assisted Conditions, Org. 2006, **8**, 2707–2710; (d) R. L. LaLonde, W. E. Brenzovich, Jr., D. Benitez, E. Tkatchouk, K. Kelley, W. A. Goddard, III and F. D. Toste, Alkylgold Complexes by the Intramolecular Aminoauration of Unactivated Alkenes, Chem. Sci., 2010, 1, 226-233; (e) S. Zhu, L. Ye, W. Wu and N-Heterocyclic Carbene-Gold(I)-Catalyzed Carboheterofunctionalization of Alkenes with Arylboronic Tetrahedron, 10375-10383; Acids, 2013, 69, (f) A. S. K. Hashmi, L. Schwarz, J. H. Choi and T. M. Frost, A New Gold-Catalyzed C-C Bond Formation, Angew. Chem., Int. Ed., 2000, 39, 2285-2288; (g) A. Aponick, C.-Y. Li and B. Biannic, Au-Catalyzed Cyclization of Monoallylic Diols, Org. Lett., 2008, 10, 669-671.
- 11 (a) K. E. Soklou, H. Marzag, J. P. Bouillon, M. Marchivie, S. Routier and K. Plé, Gold(I)-Catalyzed Intramolecular Hydroamination and Hydroalkoxylation of Alkynes: Access to Original Heterospirocycles, Org. Lett., 2020, 22, 5973-5977; (b) M. C. Blanco Jaimes, F. Rominger, M. M. Pereira, R. M. Carrilho, S. A. Carabineiro and A. S. K. Hashmi, Active Phosphite Gold(I) Intramolecular Hydroalkoxylation, Enyne Cyclization and Furanyne Cyclization, Chem. Commun., 2014, 50, 4937-4940; (c) Z.-Y. Han, H. Xiao, X.-H. Chen and L.-Z. Gong, Consecutive Intramolecular Hydroamination/Asymmetric Transfer Hydrogenation under Relay Catalysis of an Achiral Gold Complex/Chiral Brønsted Acid Binary System, J. Am. Chem. Soc., 2009, 131, 9182-9183; (d) T. Enomoto, A.-L. Girard, Y. Yasui and Y. Takemoto, Gold(I)-Catalyzed Tandem Reactions Initiated by Hydroamination of Alkynyl Carbamates: Application to the Synthesis of Nitidine, J. Org. Chem., 2009, 74, 9158-9164; (e) H. Chiba, S. Oishi, N. Fujii and H. Ohno, Total Synthesis of (-)-Quinocarcin by Gold(I)-Catalyzed Regioselective Hydroamination, Angew.

- Chem., Int. Ed., 2012, 51, 9169-9172; (f) X. Zeng, G. D. Frey, S. Kousar and G. Bertrand, A Cationic Gold(I) Complex as a General Catalyst for the Intermolecular Hydroamination of Alkynes: Application to the One-Pot Synthesis of Allenes from Two Alkynes and a Sacrificial Amine, Chem. - Eur. J., 2009, 15, 3056-3060; (g) V. Belting and N. Krause, Gold-Catalyzed Tandem Cycloisomerization-Hydroalkoxylation of Homopropargylic Alcohols, Org. Lett., 2006, 8, 4489-4492; (h) R. Gauthier, M. Mamone and J.-F. Paquin, Gold-Catalyzed Hydrofluorination of Internal Alkynes Using Aqueous HF, Org. Lett., 2019, 21, 9024-9027; (i) A. Yamaguchi, S. Inuki, Y. Tokimizu, S. Oishi and H. Ohno, Gold(I)-Catalyzed Cascade Cyclization of Anilines with Diynes: Controllable Formation of Eight-Membered Ring-Fused Indoles and Propellane-Type Indolines, J. Org. Chem., 2020, 85, 2543-2559; (j) D. P. Zimin, D. V. Dar'in, V. A. Rassadin and V. Y. Kukushkin, Gold-Catalyzed Hydrohydrazidation of Terminal Alkynes, Org. Lett., 2018, 20, 4880-4884; (k) J. A. Goodwin and A. Aponick, Regioselectivity in the Au-Catalyzed Hydration and Hydroalkoxylation of Alkynes, Chem. Commun., 2015, 51, 8730-8741.
- 12 (a) Y. Yamamoto and N. Nishina, Gold-Catalyzed Intermolecular Hydroamination of Allenes: First Example of the Use of an Aliphatic Amine in Hydroamination, Synlett, 2007, 1767-1770; (b) D. Eom, D. Kang and P. H. Lee, Synthesis of 2-Alkyl- and Aryl-3-Ethoxycarbonyl-2,5-Dihydrofurans through Gold-Catalyzed Intramolecular Hydroalkoxylation, J. Org. Chem., 2010, 75, 7447-7450; (c) R. E. M. Brooner, T. J. Brown, M. A. Chee and R. A. Widenhoefer, Effect of Substitution, Ring Size, and Counterion on the Intermediates Generated in the Gold-Catalyzed Intramolecular Hydroalkoxylation of Allenes, Organometallics, 2016, 35, 2014-2021; (d) Menggenbateer, M. Narsireddy, G. Ferrara, N. Nishina, T. Jin and Y. Yamamoto, Gold-Catalyzed Regiospecific Intermolecular Hydrothiolation of Allenes, Tetrahedron Lett., 2010, 51, 4627-4629; (e) N. Nishina and Y. Yamamoto, Gold-Catalyzed Hydrofunctionalization of Allenes with Nitrogen and Oxygen Nucleophiles and its Mechanistic Insight, Tetrahedron, 2009, 65, 1799-1808; (f) Z. Zhang, C. Liu, R. E. Kinder, X. Han, H. Qian and R. A. Widenhoefer, Highly Active Au(I) Catalyst for the Intramolecular exo-Hydrofunctionalization of Allenes with Carbon, Nitrogen, and Oxygen Nucleophiles, J. Am. Chem. Soc., 2006, 128, 9066-9073; (g) C. Zhang, S. Q. Zhang, H. J. Cai and D. M. Cui, Gold-Catalyzed Intermolecular Hydroamination of Allenes with Sulfonamides, Beilstein J. Org. Chem., 2013, 9, 1045-1050; (h) Z. Wang, C. Nicolini, C. Hervieu, Y.-F. Wong, G. Zanoni and L. Zhang, Remote Cooperative Group Strategy Enables Ligands for Accelerative Asymmetric Gold Catalysis, J. Am. Chem. Soc., 2017, 139, 16064-16067.
- 13 (a) T. Wang and A. S. K. Hashmi, 1,2-Migrations onto Gold Carbene Centers, Chem. Rev., 2021, 121, 8948-8978; (b) L. W. Ye, X. Q. Zhu, R. L. Sahani, Y. Xu, P. C. Qian and

- R. S. Liu, Nitrene Transfer and Carbene Transfer in Gold Catalysis, Chem. Rev., 2021, 121, 9039-9112; (c) G.-Q. Gao, G. Ma, X.-L. Jiang, Q. Liu, C.-L. Fan, D.-C. Lv, H. Su, G.-X. Ru and W.-B. Shen, Gold-Catalyzed Cycloadditions of Allenes via Metal Carbenes, Org. Biomol. Chem., 2022, 20, 5035-5044; (d) R. J. Harris and R. A. Widenhoefer, Gold Carbenes, Gold-Stabilized Carbocations, and Cationic Intermediates Relevant to Gold-Catalysed Cycloaddition, Chem. Soc. Rev., 2016, 45, 4533-4551; (e) L. Zhang, A Non-Diazo Approach to α-Oxo Gold Carbenes via Gold-Catalyzed Alkyne Oxidation, Acc. Chem. Res., 2014, 47, 877-888.
- 14 D. Y. Li, Y. Wei, I. Marek, X. Y. Tang and M. Shi, Gold(I)-Catalyzed Cycloisomerization of Vinylidenecyclopropaneenes via Carbene or non-Carbene Processes, Chem. Sci., 2015, 6, 5519-5525.
- 15 (a) W. Li, W. Yuan, S. Pindi, M. Shi and G. Li, Au/Ag-Catalyzed Intramolecular Ring-Opening Vinylidenecyclopropanes (VDCPs): An Easy Access to Functional Tetrahydropyrans, Org. Lett., 2010, 12, 920-923; (b) B.-L. Lu, Y. Wei and M. Shi, Gold(I) and Brønsted Acid Catalyzed Intramolecular Rearrangements Vinylidenecyclopropanes, Chem. - Eur. J., 2010, 16, 10975-10979; (c) B.-L. Lu and M. Shi, Gold(I)-Catalyzed Tandem Oxidative Ring-Opening/C-C Bond Cleavage Reactions of Vinylidenecyclopropanes with Secondary Amines Under an Oxygen Atmosphere, Chem. - Eur. J., 2011, 17, 9070-9075; (d) W. Yuan, X. Tang, Y. Wei and M. Shi, Gold-Catalyzed Cycloisomerization of Yne-Vinylidenecyclopropanes: A Three-Carbon Synthon for [3+2] Cycloadditions, Chem. -Eur. J., 2014, 20, 3198-3204; (e) L. Wu and M. Shi, Yb or Au(I)-Catalyzed Domino Intramolecular Hydroamination and Ring-Opening of Sulfonamide-Substituted 1,1-Vinylidenecyclopropanediesters, Chem. -Eur. J., 2011, 17, 13160-13165; (f) W. Zang, Y. Wei and Gold(I) Catalyzed Cascade Cyclization: Intramolecular Two-Fold Nucleophilic Addition
- Vinylidenecyclopropanes (VDCPs), Org. Chem. Front., 2018, 5, 197-202; (g) W. Yuan, X. Dong, Y. Wei and M. Shi, An Efficient Method for the Synthesis Alkylidenecyclobutanones by Gold-Catalyzed Oxidative Ring Enlargement of Vinylidenecyclopropanes, Chem. -Eur. I., 2012, 18, 10501-10505; (h) D.-Y. Li, Y. Wei and M. Shi, Gold(I)-Catalyzed Intramolecular Carbon-Oxygen Bond Cleavage Reaction via Gold Carbenes Derived from Vinylidenecyclopropanes, Adv. Synth. Catal., 2016, 358, 3002-3009; (i) D.-Y. Li, W. Fang, Y. Wei and M. Shi, C(sp³)-H Functionalizations Promoted by the Gold Carbene Generated from Vinylidenecyclopropanes, Chem. - Eur. I., 2016, 22, 18080-18084.
- 16 (a) E. Solel, M. Ruth and P. R. Schreiner, London Dispersion Helps Refine Steric A-Values: The Halogens, J. Org. Chem., 2021, 86, 7701-7713; (b) F. R. Jensen, C. H. Bushwell and B. H. Beck, Conformational Preferences in Monosubstituted Cyclohexanes Determined by Nuclear Magnetic Resonance Spectroscopy, J. Am. Chem. Soc., 1969, 91, 344-351; (c) C. H. Marzabadi, J. E. Anderson, J. Gonzalez-Outeirino, P. R. J. Gaffney, C. G. H. White, D. A. Tocher and L. J. Todaro, Why Are Silyl Ethers Conformationally Different from Alkyl Ethers? Chair-Chair Conformational Equilibria in Silyloxycyclohexanes and Their Dependence on the Substituents on Silicon. The Wider Roles of Eclipsing, of 1,3-Repulsive Steric Interactions, and of Attractive Steric Interactions, J. Am. Chem. Soc., 2003, 125, 15163-15173; (d) S. Winstein and N. J. Holness, Neighboring Carbon and Hydrogen. XIX. t-Butylcyclohexyl Derivatives. Quantitative Conformational Analysis, J. Am. Chem. Soc., 1955, 77, 5562-5578.
- 17 W. Yang and A. S. K. Hashmi, Mechanistic Insights into the Gold Chemistry of Allenes, Chem. Soc. Rev., 2014, 43, 2941-2955.
- 18 M. Jiang and M. Shi, Palladium-Catalyzed Diacetoxylation of Methylenecyclopropanes via $C(sp^3)-C(sp^3)$ Breaking, Organometallics, 2009, 28, 5600-5602.