


RESEARCH ARTICLE

[View Article Online](#)
[View Journal](#) | [View Issue](#)



Cite this: *Org. Chem. Front.*, 2023, **10**, 4658

An unusual four-step cascade reaction for accessing furo[3,4-c]pyridine-1,4-diones via rhodium catalysis†

Yidi Li,^{‡,a,e} Huiying Xu, ^{‡,c} Zhi Zhou, ^{‡,d} Lin Huang, ^{b,d} Zhenhao Tang, ^d Wei Yi ^{‡,c} and Xiaowei Wu ^{‡,a,b,d,e}

The development of efficient cascade reactions is highly important and appealing because of their desirable step-economy and convenience in constructing multiple chemical bonds and complex molecules in one shot. Herein, we report a rare four-step tandem reaction between acrylamides and 4-hydroxy-2-alkynoates to prepare novel furo[3,4-c]pyridine-1,4-diones which are difficult to synthesize by traditional methods. This unique domino reaction includes C–H activation, Lossen rearrangement, annulation, and lactonization. Additionally, this protocol features good functional group tolerance, obtainment of products by simple filtration, room temperature, and air compatibility. DFT calculations were conducted to shed some light on the reaction mechanism.

Received 19th June 2023,
Accepted 5th August 2023

DOI: 10.1039/d3qo00909b
rsc.li/frontiers-organic

Introduction

The prevalence of heterocycles in natural products and pharmaceuticals has led to the importance and benefits of synthesizing such compounds efficiently. In the past decades, directing group (DG) assisted C–H activation/annulation reactions *via* transition-metal catalysis for the synthesis of heterocycles have achieved tremendous advances.^{1,2} DGs play crucial roles in controlling chemical reactivity and regioselectivity in C–H activation reactions. On the other hand, DGs simply adhere to primordial positions and function as auxiliary groups in these reactions without other transformations. Amide-type DGs are frequently applied to C–H activation/annulation reactions for the synthesis of prevalent isoquinolones

and 2-pyridones, where DGs remain static in most cases (Scheme 1a).^{1,2a–f,3} The rearrangement or migration of DGs in C–H activation reactions provides an intriguing strategy to construct C–C/C–N bonds and also contributes to the synthesis of novel heterocyclic scaffolds.^{4,5} Therefore, it is fascinating to apply this emerging strategy to organic synthesis. Nevertheless, the development of this emerging strategy involving DG rearrangement or migration still falls far behind when compared to the numerous reports of static DGs participating in C–H activation reactions.

Even though DGs could improve chemoselectivity and regioselectivity, they may be present as undesired moieties on products at times, and additional steps are required to remove unwanted DGs. If DGs undergo a rearrangement process, it will not only create a new and efficient avenue for synthesizing heterocycles, but also possibly generate novel heterocycles that traditional approaches can hardly furnish. By taking advantage of the strategy of directing group rearrangement, a series of intriguing spirooxindole pyrrolones were synthesized elegantly by Dai's group (Scheme 1b).⁶ Acrylamides involving the rearrangement of DGs undergo three processes: C–H activation, directing group rearrangement, and annulation. Recently, we developed an efficient rhodium-catalyzed domino C–H alkenylation, DG migration, and lactonization reaction between *N*-carbamoyl indoles and 4-hydroxy-2-alkynoates for the synthesis of the furan-2(5*H*)-one scaffold (Scheme 1c).⁷ Efficient cascade reactions are highly appealing due to their desirable step-economy and convenience in constructing complex molecules with multiple chemical bonds in one shot.^{8–11}

To our knowledge, the implementation of a multistep cascade reaction involving four distinct processes of C–H acti-

^aDrug Discovery and Development Center, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai 201203, China.

E-mail: zhouzhi@gzmu.edu.cn, wuxiaowei@simm.ac.cn

^bSchool of Pharmaceutical Sciences, Southern Medical University, Guangzhou 510515, China

^cGuangzhou Municipal and Guangdong Provincial Key Laboratory of Protein Modification and Degradation & Molecular Target and Clinical Pharmacology, State Key Laboratory of Respiratory Disease, School of Pharmaceutical Sciences & the Fifth Affiliated Hospital, Guangzhou Medical University, Guangzhou, Guangdong 511436, China. E-mail: yiwei@gzmu.edu.cn

^dZhongshan Institute for Drug Discovery, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Zhongshan 528400, China




^eUniversity of Chinese Academy of Sciences, No. 19A Yuquan Road, Beijing 100049, China

† Electronic supplementary information (ESI) available. CCDC 2281784. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d3qo00909b>


‡ These authors contributed equally.



Previous work:

a) C(sp²)-H activation/[4+2] annulation assisted by static DGs (well-established)b) C(sp²)-H activation/DG Lossen rearrangement/annulation with diazo oxindolesc) C(sp²)-H activation/DG migration/lactonization with 4-hydroxy-2-alkynoates

This work:

d) Unprecedented C(sp²)-H activation/DG Lossen rearrangement/annulation/lactonization cascade

Scheme 1 Transition-metal catalyzed C–H activation/annulation reactions.

vation, directing group rearrangement, annulation, and lactonization has yet to be documented. In this study, we report a rare four-step tandem reaction between acrylamides and 4-hydroxy-2-alkynoates¹² to prepare novel furo[3,4-*c*]pyridine-1,4-diones which are difficult to synthesize by traditional methods. The four-step tandem reaction involves C–H activation, DG Lossen rearrangement, [4 + 2] annulation, and lactonization. The method can tolerate a variety of functional groups. Furthermore, the products can be easily purified by filtration and the reaction is compatible with both room temperature and air. These features contribute to the practicality and versatility of the protocol.

Results and discussion

We embarked on our studies with the optimization of the reaction conditions for the cyclization between acrylamide **1a** and ethyl 4-hydroxy-4-phenylbut-2-ynoate **2a**. No desired product **3a** was obtained when catalyzed by $\text{MnBr}(\text{CO})_5$, $[\text{RuCl}_2(p\text{-cym})]_2$, $\text{Cp}^*\text{Co}(\text{CO})\text{I}_2$ and $\text{Pd}(\text{OAc})_2$, respectively (Table 1, entries 1–6). Switching to $[\text{Cp}^*\text{IrCl}_2]_2$ led to only 6% yield (entry 2). In contrast, the yield of **3a** was improved to 73%, and 43% of the product could be obtained by simple filtration when using $[\text{Cp}^*\text{RhCl}_2]_2$ (entry 5). The structure of **3a** was confirmed using 1D-NMR and NOESY experiments as well as mass spectra. Solvent screening revealed that TFE outperformed

Table 1 Optimization of reaction conditions^a


| Entry | Catalyst | Additive | Solvent | Yield ^{b,c} (%) |
|-----------------|---|--------------------------|------------------------|--------------------------|
| 1 | $\text{MnBr}(\text{CO})_5$ | NaOAc | TFE | 0 |
| 2 | $[\text{Cp}^*\text{IrCl}_2]_2$ | NaOAc | TFE | 6 |
| 3 | $\text{Cp}^*\text{Co}(\text{CO})\text{I}_2$ | NaOAc | TFE | 0 |
| 4 | $[\text{Ru}(p\text{-cym})\text{Cl}_2]_2$ | NaOAc | TFE | 0 |
| 5 | $[\text{Cp}^*\text{RhCl}_2]_2$ | NaOAc | TFE | 73 (47) |
| 6 | $\text{Pd}(\text{OAc})_2$ | NaOAc | TFE | 0 |
| 7 | $[\text{Cp}^*\text{RhCl}_2]_2$ | NaOAc | THF | Trace |
| 8 | $[\text{Cp}^*\text{RhCl}_2]_2$ | NaOAc | MeOH | 15 |
| 9 | $[\text{Cp}^*\text{RhCl}_2]_2$ | NaOAc | DCE | 32 |
| 10 | $[\text{Cp}^*\text{RhCl}_2]_2$ | NaOAc | CH_3CN | 0 |
| 11 | $[\text{Cp}^*\text{RhCl}_2]_2$ | NaOAc | EA | 20 |
| 12 | $[\text{Cp}^*\text{RhCl}_2]_2$ | Na_2CO_3 | EtOH | 0 |
| 13 | $[\text{Cp}^*\text{RhCl}_2]_2$ | Na_2CO_3 | TFE | 78 (53) |
| 14 | $[\text{Cp}^*\text{RhCl}_2]_2$ | K_2CO_3 | TFE | 19 |
| 15 | $[\text{Cp}^*\text{RhCl}_2]_2$ | CsOAc | TFE | 54 (37) |
| 16 | $[\text{Cp}^*\text{RhCl}_2]_2$ | KOAc | TFE | 46 (33) |
| 17 ^d | $[\text{Cp}^*\text{RhCl}_2]_2$ | — | TFE | 0 |
| 18 ^e | $[\text{Cp}^*\text{RhCl}_2]_2$ | Na_2CO_3 | TFE | 0 |

^a Reaction conditions: **1a** (0.2 mmol), **2a** (0.24 mmol), catalyst (3 mol%), additive (0.2 mmol), under air, solvent (1.0 mL), room temperature, 6 h. ^b Total isolated yield. ^c Isolated yields obtained by simple filtration are shown in parenthesis. ^d Na_2CO_3 was not used. ^e $[\text{Cp}^*\text{RhCl}_2]_2$ was not used.

other common solvents, such as THF (only trace), MeOH (15% yield), DCE (32% yield with 40% substrate not being consumed) and EA (20% yield) (entries 7–11). When using ethanol and adding Na_2CO_3 as a base, the yield of **3a** was 0% (entry 12). After extensive screening of additives, such as Na_2CO_3 , K_2CO_3 , CsOAc , and KOAc, Na_2CO_3 was identified as the optimal one (entries 13–16). The control experiments confirmed that no desired reaction occurred in the absence of an additive or rhodium (entries 17 and 18).

With the optimal conditions in hand, we next examined the scope of this cyclization reaction (Scheme 2). Aryl-substituted propynol bearing electron-donating groups are all tolerated in this system, such as CH_3 , *t*-Bu, OMe, and CH_2CH_3 , and the corresponding products can be obtained in high yields by direct filtration (**3a**–**e**). To our delight, electron-withdrawing groups including F, Cl, Br, and I were also tolerated well with the isolated yields ranging from 62% to 72% (**3f**–**i**). Although the yield of the product dropped dramatically when the group was changed to cyclopentyl (**3k**), the reaction was compatible with other different rings (naphthalene, furan, and thiophene) and afforded the products smoothly (**3j**, **3l**, and **3m**). We also explored tertiary alkynol and obtained the desired product in a moderate yield (**3n**). Next, we investigated R^1 of acrylamide **1** bearing electron-donating and electron-withdrawing groups at the *para* position of the phenyl group, and the reactions proceeded well to generate the corresponding products. In comparison, when the phenyl group was changed to the benzyl





Scheme 2 Substrate scope. Reaction conditions: **1** (0.2 mmol), **2** (0.24 mmol), $[\text{Cp}^*\text{RhCl}_2]_2$ (3 mmol%), Na_2CO_3 (0.2 mmol), under air, trifluoroethanol (1.0 mL), rt, 6–12 h. Total yields are shown, and yields obtained by simple filtration are shown in parentheses.

and methyl group, the yield dropped to 47% and 20%, respectively (**3o–3u**). Additionally, the structures of **3h**, **3n**, **3p**, and **3q** were further confirmed by NOESY experiments.


Several experiments were conducted to probe the reaction mechanism. In the presence of **1a**, Na_2CO_3 , $[\text{Cp}^*\text{RhCl}_2]_2$, and 1.0 mL methanol-*d*⁴, the experiment of H/D exchange occurring at the olefinic bond of acrylamide was performed (Scheme 3a). And no obvious deuterium incorporation was observed at this position (8% D). In addition, there was also almost no D at the olefinic bond of **3a** after adding **2a**. The result demonstrated that the cleavage of the C–H bond could be irreversible.

Additionally, two parallel reactions were performed giving a KIE value of 0.98 (Scheme 3b), which suggested that the C–H activation step was less likely to be involved in the rate-determining step.

To further obtain the mechanistic features of the unprecedented Rh(III)-catalyzed sequential C–H activation/Losken rearrangement/[4 + 2] annulation/lactonization cascade, we next performed DFT calculations on the key reaction steps: alkyne insertion, Losken rearrangement, [4 + 2] annulation and lactonization. As shown in Scheme 4, the five-membered rhodacycle **INT-0** derived from C–H activation was rationally




a) Deuterium incorporation experiments



b) Kinetic isotope effect study


parallel reactions



Scheme 3 Deuteration and KIE experiments.

selected as the starting point, which coordinated with propargyl alcohol 2a and produced the intermediate **INT-1** ($\Delta G^\ddagger = -11.3$ kcal mol⁻¹) involving hydrogen-bonding between the

hydroxyl group of **2a** and the pivalate group. Then the C–C unsaturated bond of **2a** was inserted into the Rh–C bond *via* **TS-1** ($\Delta G^\ddagger = -4.6$ kcal mol⁻¹) to deliver **INT-2** with a free energy of -35.8 kcal mol⁻¹. The subsequent facile coordination change *via* **TS-2** ($\Delta G^\ddagger = -28.1$ kcal mol⁻¹) produced the unstable intermediate **INT-3** which afforded different reaction paths. The concerted Lossen rearrangement/N–O bond cleavage from **INT-3** occurred *via* **TS-3** ($\Delta G^\ddagger = -26.9$ kcal mol⁻¹) which demanded an energy barrier of 8.9 kcal mol⁻¹ (from **INT-2** to **TS-3**) to give the intermediate **INT-4** along with a considerable quantity of heat. The relatively elongated N–O¹ bond (1.48 Å vs. 1.42 Å in **INT-0**) in the geometry of **INT-3** and the significantly elongated N–O¹ bond (2.41 Å) and shortened C¹–N bond (1.98 Å) in the transition state **TS-3** suggest that the Lossen rearrangement was assisted by the pivalate group and the rhodium center. Alternatively, classic C–N bond reductive elimination *via* **TS-3ii** ($\Delta G^\ddagger = -14.7$ kcal mol⁻¹) involved a higher energy barrier of 21.1 kcal mol⁻¹ (from **INT-2** to **TS-3ii**). Moreover, the Rh(III)–Rh(V)–Rh(III) reaction path involving an oxidative addition process from **INT-3** was also ruled out owing to the relatively high free energies of **INT-4i/TS-4i**. After low-barrier isomerization *via* **TS-4**, **INT-4** was transformed into the isocyanate intermediate **INT-5** from which the direct C–C bond formation/annulation occurred *via* **TS-5i** ($\Delta G^\ddagger = -50.2$ kcal mol⁻¹) and required a barrier of 22.4 kcal mol⁻¹ (from **INT-5** to **TS-5i**). However, the coordination change *via*



Scheme 4 DFT calculations of the reaction pathways.



TS-5/TS-6 led to a more favorable C-C bond formation reaction *via* **TS-7** ($\Delta G^\ddagger = -55.1$ kcal mol⁻¹) which only needs an activation energy of 17.5 kcal mol⁻¹. Afterwards, the protonolysis reaction *via* **INT-9/TS-8** was followed by lactonization *via* **TS-9** ($\Delta G^\ddagger = -81.0$ kcal mol⁻¹). Alternatively, the metal-free lactonization without the rhodium complex goes through **TS-9i** ($\Delta G^\ddagger = -60.2$ kcal mol⁻¹) which is of a much higher barrier and can be ruled out.

In order to shed light on the driving force of the Lossen rearrangement, we examined closely the geometry change along the intrinsic reaction coordinate (IRC) corresponding to **TS-3** (see the ESI for details†). The results revealed that at the beginning the N–O¹ bond distance increases which seems to drive the C¹ atom attack on the N atom. The reaction goes through **TS-3** and then the N–O¹ bond breaks which is followed by C¹–N bond formation and C¹–C² bond cleavage. This gives a chance for the O¹ atom to attack the positively charged C² atom. With the assistance of the pivalate group and the rhodium center, the Lossen rearrangement involves multiple σ-bond metathesis processes.

Conclusions

In summary, we have developed a rare four-step tandem reaction between acrylamides and 4-hydroxy-2-alkynoates to prepare novel furo[3,4-*c*]pyridine-1,4-diones. This unique tandem reaction consists of C–H activation, Lossen rearrangement, [4 + 2] annulation, and lactonization. The protocol features good functional group tolerance, obtainment of pure products by simple filtration, room temperature, and air compatibility. In addition, DFT calculations were conducted to shed some light on the reaction mechanism.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Financial support from the Basic and Applied Basic Research Foundation of Guangdong Province (2021A1515110468), Shanghai Pujiang Program (21PJ1415800), Natural Science Foundation of Guangdong Province (2019A1515010935), High-level New R&D Institute (2019B090904008), High-level Innovative Research Institute (2021B0909050003), and NSFC (21877020 and 22007020) is gratefully acknowledged. We also thank Wei Zhang from the Zhongshan Institute for Drug Discovery for X-ray crystallographic analysis.

References

1 (a) P. Gandeepan, T. Müller, D. Zell, G. Cera, S. Warratz and L. Ackermann, 3d Transition Metals for C–H Activation,

Chem. Rev., 2019, **119**, 2192–2452; (b) D. J. Abrams, P. A. Provencher and E. J. Sorensen, Recent applications of C–H functionalization in complex natural product synthesis, *Chem. Soc. Rev.*, 2018, **47**, 8925–8967; (c) C. Sambiagio, D. Schönbauer, R. Blieck, T. D. Huy, G. Pototschnig, P. Schaaf, T. Wiesinger, M. F. Zia, J. W. Delord, T. Basset, B. U. W. Maes and M. Schnürch, A comprehensive overview of directing groups applied in metal-catalysed C–H functionalisation chemistry, *Chem. Soc. Rev.*, 2018, **47**, 6603–6743; (d) Y. Yang, J. Lan and J. You, Oxidative C–H/C–H Coupling Reactions between Two (Hetero)arenes, *Chem. Rev.*, 2017, **117**, 8787–8863; (e) T. Gensch, M. N. Hopkinson, F. Glorius and J. W. Delord, Mild metal-catalyzed C–H activation: examples and concepts, *Chem. Soc. Rev.*, 2016, **45**, 2900–2936; (f) P. Augustin, G. Catherine and C. Laurent, Recent Advances in [Cp^{*}M^{III}] (M = Co, Rh, Ir)-Catalyzed Intramolecular Annulation Through C–H Activation, *Eur. J. Org. Chem.*, 2018, 5777–5794; (g) D. Mandal, S. Roychowdhury, J. P. Biswas, S. Maiti and D. Maiti, Transition-metal-catalyzed C–H bond alkylation using olefins: recent advances and mechanistic aspects, *Chem. Soc. Rev.*, 2022, **51**, 7358–7426; (h) L. Ackermann, Metalla-electrocatalyzed C–H Activation by Earth-Abundant 3d Metals and Beyond, *Acc. Chem. Res.*, 2020, **53**, 84–104.

2 (a) S. Mochida, N. Umeda, K. Hirano, T. Satoh and M. Miura, Rhodium-catalyzed Oxidative Coupling/Cyclization of Benzamides with Alkynes via C–H Bond Cleavage, *Chem. Lett.*, 2010, **39**, 744–746; (b) T. K. Hyster and T. Rovis, Rhodium-Catalyzed Oxidative Cycloaddition of Benzamides and Alkynes via C–H/N–H Activation, *J. Am. Chem. Soc.*, 2010, **132**, 10565–10569; (c) N. Guimond, C. Gouliaras and K. Fagnou, Rhodium(III)-catalyzed isoquinolone synthesis: the N–O bond as a handle for C–N bond formation and catalyst turnover, *J. Am. Chem. Soc.*, 2010, **132**, 6908–6909; (d) Y. Su, M. Zhao, K. Han, G. Song and X. Li, Synthesis of 2-pyridones and iminoesters via Rh(III)-catalyzed oxidative coupling between acrylamides and alkynes, *Org. Lett.*, 2010, **12**, 5462–5465; (e) X. Xu, Y. Liu and C.-M. Park, Rhodium(III)-catalyzed intramolecular annulation through C–H activation: total synthesis of (±)-antofine, (±)-septicine, (±)-tylophorine, and rosetacin, *Angew. Chem., Int. Ed.*, 2012, **51**, 9372–9376; (f) N. Quiñones, A. Seoane, R. G. Fandiño, J. L. Mascareñas and M. Gulías, Rhodium(III)-catalyzed intramolecular annulations involving amide-directed C–H activations: synthetic scope and mechanistic studies, *Chem. Sci.*, 2013, **4**, 2874–2879; (g) X. Wu, B. Wang, Y. Zhou and H. Liu, Propargyl Alcohols as One-Carbon Synthons: Redox-Neutral Rhodium (III)-Catalyzed C–H Bond Activation for the Synthesis of Isoindolinones Bearing a Quaternary Carbon, *Org. Lett.*, 2017, **19**, 1294–1297; (h) X. Wu, B. Wang, S. Zhou, Y. Zhou and H. Liu, Ruthenium-Catalyzed Redox-Neutral [4 + 1] Annulation of Benzamides and Propargyl Alcohols via C–H Bond Activation, *ACS Catal.*, 2017, **4**, 2494–2499; (i) L. Grigorjeva and O. Daugulis, Cobalt-catalyzed, amino-



quinoline-directed C(sp²)-H bond alkenylation by alkynes, *Angew. Chem., Int. Ed.*, 2014, **53**, 10209–10212; (j) A. Lerchen, T. Knecht, M. Koy, C. G. Daniliuc and F. Glorius, A General Cp^{*}Co^{III}-Catalyzed Intramolecular C–H Activation Approach for the Efficient Total Syntheses of Aromathecin, Protoberberine, and Tylophora Alkaloids, *Chem. – Eur. J.*, 2017, **23**, 12149–12152; (k) X. Wu and H. Ji, Rhodium-Catalyzed [4 + 1] Cyclization via C–H Activation for the Synthesis of Divergent Heterocycles Bearing a Quaternary Carbon, *J. Org. Chem.*, 2018, **83**, 4650–4656; (l) X. Wu, P. Li, Y. Lu, J. Qiao, J. Zhao, X. Jia, H. Ni, L. Kong, X. Zhang and F. Zhao, Rhodium-Catalyzed Cascade Reactions of Indoles with 4-Hydroxy-2-Alkynoates for the Synthesis of Indole-Fused Polyheterocycles, *Adv. Synth. Catal.*, 2020, **362**, 2953–2960; (m) Y. Yu, L. Huang, W. Wu and H. Jiang, Palladium-catalyzed oxidative annulation of acrylic acid and amide with alkynes: a practical route to synthesize α -pyrones and pyridines, *Org. Lett.*, 2014, **16**, 2146–2149.

3 (a) H. Meng, H. Xu, Z. Zhou, Z. Tang, Y. Li, Y. Zhou, W. Yi and X. Wu, Recyclable rhodium-catalyzed C–H activation/[4 + 2] annulation with unconventional regioselectivity at ambient temperature: experimental development and mechanistic insight, *Green Chem.*, 2022, **24**, 7012–7021; (b) T. von Münchow, S. Dana, Y. Xu, B. Yuan and L. Ackermann, Enantioselective electrochemical cobalt-catalyzed aryl C–H activation reactions, *Science*, 2023, **379**, 1036–1042; (c) P. Wang, H. Wu, X.-P. Zhang, G. Huang, R. H. Crabtree and X. Li, Sigma-Bond Metathesis as an Unusual Asymmetric Induction Step in Rhodium-Catalyzed Enantiodivergent Synthesis of C–N Axially Chiral Biaryls, *J. Am. Chem. Soc.*, 2023, **145**, 8417–8429; (d) Y. Chao, L. Shi, F. Wang, Y. Su, J.-B. Xia and F. Li, Rhodium-Catalyzed Asymmetric (3 + 2 + 2) Annulation via N–H/C–H Dual Activation and Internal Alkyne Insertion toward N-Fused 5/7 Bicycles, *ACS Catal.*, 2022, **12**, 14194–14208; (e) T. Kurihara, M. Kojima, T. Yoshino and S. Matsunaga, Cp^{*}Rh^(III)/Chiral Lewis Base Cooperative Catalysis for Enantioselective Cyclization via C–H Activation, *J. Am. Chem. Soc.*, 2022, **144**, 7058–7065; (f) R. K. Shukla, A. M. Nair and C. M. R. Volla, Pd(ii)-catalyzed β - and γ -C(sp³)-H dienylation with allenyl acetates, *Chem. Sci.*, 2022, **14**, 955–962.

4 (a) C.-Q. Wang, Y. Zhang and C. Feng, Fluorine Effects on Group Migration via a Rhodium(v) Nitrenoid Intermediate, *Angew. Chem., Int. Ed.*, 2017, **56**, 14918–14922; (b) T. Yamada, Y. Shibata and K. Tanaka, Formal Lossen Rearrangement/Alkenylation or Annulation Cascade of Heterole Carboxamides with Alkynes Catalyzed by CpRh^{III} Complexes with Pendant Amides, *Chem. – Eur. J.*, 2019, **25**, 16022–16031; (c) T. Yamada, Y. Shibata, S. Kawauchi, S. Yoshizaki and K. Tanaka, Formal Lossen Rearrangement/[3 + 2] Annulation Cascade Catalyzed by a Modified Cyclopentadienyl Rh^{III} Complex, *Chem. – Eur. J.*, 2018, **24**, 5723–5727; (d) M. Bian, H. Mawjudha, H. Gao, H. Xu, Z. Zhou and W. Yi, Lossen Rearrangement vs C–N Reductive Elimination Enabled by Rh^(III)-Catalyzed C–H Activation/Selective Lactone Ring-Opening: Chemodivergent Synthesis of Quinolinones and Dihydroisoquinolinones, *Org. Lett.*, 2020, **22**, 9677–9682; (e) J.-F. Tan, C. T. Bormann, K. Severin and N. Cramer, Alkynyl Triazenes as Fluoroalkyne Surrogates: Regioselective Access to 4-Fluoro-2-pyridones by a Rh^(III)-Catalyzed C–H Activation–Lossen Rearrangement–Wallach Reaction, *ACS Catal.*, 2020, **10**, 3790–3796; (f) M. Peng, C.-S. Wang, P.-P. Chen, T. Roisnel, H. Doucet, K. N. Houk and J. F. Soulé, Merging C–H Bond Activation, Alkyne Insertion, and Rearrangements by Rh^(III)-Catalysis: Oxindole Synthesis from Nitroarenes and Alkynes, *J. Am. Chem. Soc.*, 2023, **145**, 4508–4516; (g) Y. Li, H. Xu, L. Huang, Z. Zhou, Z. Tang, H. Meng, W. Zhang, W. Yi and X. Wu, Rhodium-catalyzed Regioselective C–H Activation/ Lossen Rearrangement/Annulation for the Green Synthesis of Trisubstituted 2-Pyridones, *Org. Chem. Front.*, 2023, **10**, 3000–3009; (h) R. B. Dateer and S. Chang, Selective Cyclization of Arylnitrones to Indolines under External Oxidant-Free Conditions: Dual Role of Rh^(III) Catalyst in the C–H Activation and Oxygen Atom Transfer, *J. Am. Chem. Soc.*, 2015, **137**, 4908–4911; (i) J.-L. Pan, P. Xie, C. Chen, Y. Hao, C. Liu, H.-Y. Bai, J. Ding, L.-R. Wang, Y. Xia and S.-Y. Zhang, Rhodium^(III)-Catalyzed Redox-Neutral Cascade [3 + 2] Annulation of N-Phenoxyacetamides with Propiolates via C–H Functionalization/Isomerization/ Lactonization, *Org. Lett.*, 2018, **20**, 7131–7136.

5 (a) Y. Wu, C. Pi, Y. Wu and X. Cui, Directing group migration strategy in transition-metal-catalysed direct C–H functionalization, *Chem. Soc. Rev.*, 2021, **50**, 3677–3689; (b) X. Zhou, Z. Fan, Z. Zhang, P. Lu and Y. Wang, Construction of Pyrrolo [1,2-a]indoles via Cobalt^(III)-Catalyzed Enaminylation of 1-(Pyrimidin-2-yl)-1H-indoles with Ketenimines and Subsequent Base-Promoted Cyclization, *Org. Lett.*, 2016, **18**, 4706–4709; (c) S.-Y. Chen, X.-L. Han, J.-Q. Wu, Q. Li, Y. Chen and H. Wang, Manganese(i)-Catalyzed Regio- and Stereoselective 1,2-Diheteroarylation of Allenes: Combination of C–H Activation and Smiles Rearrangement, *Angew. Chem., Int. Ed.*, 2017, **56**, 9939–9943; (d) C. Wang, A. Wang and M. Rueping, Manganese Catalyzed C–H Functionalizations: Hydroarylations and Alkenylations Involving an Unexpected Heteroaryl Shift, *Angew. Chem., Int. Ed.*, 2017, **56**, 9935–9938; (e) C. Zhu, R. Kuniyil, B. B. Jei and L. Ackermann, Domino C–H Activation/Directing Group Migration/Alkyne Annulation: Unique Selectivity by d⁶-Cobalt^(III) Catalysts, *ACS Catal.*, 2020, **10**, 4444–4450; (f) X. Wang, T. Gensch, A. Lerchen, C. G. Daniliuc and F. Glorius, Cp^{*}Rh^(III)/Bicyclic Olefin Cocatalyzed C–H Bond Amidation by Intramolecular Amide Transfer, *J. Am. Chem. Soc.*, 2017, **139**, 6506–6512; (g) Y. Wu, Z. Chen, Y. Yang, W. Zhu and B. Zhou, Rh^(III)-Catalyzed Redox Neutral Unsymmetrical C–H Alkylation and Amidation Reactions of N-Phenoxyacetamides, *J. Am. Chem. Soc.*, 2018, **140**, 42–45.

6 B. Ma, P. Wu, X. Wang, Z. Wang, H.-X. Lin and H.-X. Dai, Efficient Synthesis of Spirooxindole Pyrrolones by a



Rhodium(III)-Catalyzed C–H Activation/Carbene Insertion/Lossen Rearrangement Sequence, *Angew. Chem., Int. Ed.*, 2019, **58**, 13335–13339.

7 F. Zhao, Z. Zhou, Y. Lu, J. Qiao, X. Zhang, X. Gong, S. Liu, S. Lin, X. Wu and W. Yi, Chemo, Regio, and Stereoselective Assembly of Polysubstituted Furan-2(5H) ones Enabled by Rh(III)-Catalyzed Domino C–H Alkenylation/Directing Group Migration/Lactonization: A Combined Experimental and Computational Study, *ACS Catal.*, 2021, **11**, 13921–13934.

8 A review on cascade reactions: A. Baccalini, G. Faita, G. Zanoni and D. Maiti, Transition Metal Promoted Cascade Heterocycle Synthesis through C–H Functionalization, *Chem. – Eur. J.*, 2020, **26**, 9749–9783.

9 A review on cascade reactions: L. Song and E. V. Van der Eycken, Transition Metal-Catalyzed Intermolecular Cascade C–H Activation/Annulation Processes for the Synthesis of Polycycles, *Chem. – Eur. J.*, 2021, **27**, 121–144.

10 K. K. Gollapelli, S. Kallepu, N. Govindappa, J. B. Nanubolu and R. Chegondi, Carbonyl-assisted reverse regioselective cascade annulation of 2-acetylenic ketones triggered by Ru-catalyzed C–H activation, *Chem. Sci.*, 2016, **7**, 4748–4753.

11 Y. Fukui, P. Liu, Q. Liu, Z.-T. He, N.-Y. Wu, P. Tian and G.-Q. Lin, Tunable arylative cyclization of 1,6-enynes triggered by rhodium(III)-catalyzed C–H activation, *J. Am. Chem. Soc.*, 2014, **136**, 15607–15614.

12 (a) G. Liao, H. Song, X. Yin and B. Shi, Expeditious synthesis of pyrano[2,3,4-de]quinolines via Rh(III)-catalyzed cascade C–H activation/annulation/lactonization of quinolin-4-ol with alkynes, *Chem. Commun.*, 2017, **53**, 7824–7827; (b) Y. Xu, B. Li, X. Zhang and X. Fan, One-Pot Synthesis of Fused N,O-Heterocycles through Rh(III)-Catalyzed Cascade Reactions of Aromatic/Vinylic N-Alkoxy-Amides with 4-Hydroxy-2-Alkynoates, *Adv. Synth. Catal.*, 2018, **360**, 2613–2620; (c) A. Kumar, V. Hanchate and K. R. Prabhu, Rhodium(III)-Catalyzed Cascade Reactions of Imines/Imidates with 4-Hydroxy-2-alkynoates to Synthesize Regioselective Furanone-Fused Isoquinoline Scaffolds, *J. Org. Chem.*, 2021, **86**, 17965–17974; (d) Y.-Y. Wang, M. Liu and L. Dong, Rh(III)-Catalyzed multi-site-selective C–H bond functionalization: condition-controlled synthesis of diverse fused polycyclic benzimidazole derivatives, *Org. Chem. Front.*, 2021, **8**, 2487–2493; (e) N. Muniraj, A. Kumar and K. R. Prabhu, Cobalt-Catalyzed Regioselective [4 + 2] Annulation/Lactonization of Benzamides with 4-Hydroxy-2-Alkynoates under Aerobic Conditions, *Adv. Synth. Catal.*, 2020, **362**, 152–159; (f) A. Kumar and K. R. Prabhu, Rhodium(III)-Catalyzed C–H Activation: A Cascade Approach for the Regioselective Synthesis of Fused Heterocyclic Lactone Scaffolds, *J. Org. Chem.*, 2020, **85**, 3548–3559; (g) Y. Liu, Z. Yang, R. Chauvin, W. Fu, Z. Yao, L. Wang and X. Cui, One-Pot Synthesis of Furo[3,4-c]indolo[2,1-a]isoquinolines through Rh(III)-Catalyzed Cascade Reactions of 2-Phenylindoles with 4-Hydroxy-2-alkynoates, *Org. Lett.*, 2020, **22**, 5140–5144.