



Cite this: *RSC Adv.*, 2023, 13, 13511

# Stereoselective synthesis of (*E*)- $\alpha,\beta$ -unsaturated esters: triethylamine-catalyzed allylic rearrangement of enol phosphates†

Yulong Zhang, Huichuang Guo, Qian Wu, Xiaojing Bi, \* Enxue Shi \* and Junhua Xiao \*

$\alpha,\beta$ -Unsaturated esters are key structural motifs widely distributed in various biologically active molecules, and their *Z/E*-stereoselective synthesis has always been considered highly attractive in organic synthesis. Herein, we present a >99% (*E*)-stereoselective one-pot synthetic approach towards  $\beta$ -phosphoroxylated  $\alpha,\beta$ -unsaturated esters via a mild trimethylamine-catalyzed 1,3-hydrogen migration of the corresponding unconjugated intermediates derived from the solvent-free Perkow reaction between low-cost 4-chloroacetoacetates and phosphites. Versatile  $\beta,\beta$ -disubstituted (*E*)- $\alpha,\beta$ -unsaturated esters were thus afforded with full (*E*)-stereoretentivity by cleavage of the phosphoenol linkage via Negishi cross-coupling. Moreover, a stereoretentive (*E*)-rich mixture of a  $\alpha,\beta$ -unsaturated ester derived from 2-chloroacetoacetate was obtained and both isomers were easily afforded in one operation.

Received 12th April 2023  
Accepted 27th April 2023

DOI: 10.1039/d3ra02430j

rsc.li/rsc-advances

$\alpha,\beta$ -Unsaturated carbonyl motifs, such as the relevant esters, amides, and aldehydes, are widely distributed in biologically active molecules as key structural components (Fig. 1).<sup>1–4</sup> Generally, the (*Z*) and (*E*)-isomers of those molecules possess very different living activities.<sup>5</sup> Moreover, ubiquitous  $\alpha,\beta$ -unsaturated esters are also widely employed as useful intermediates for enantioselective hydrogenation,<sup>6</sup> allylic substitution,<sup>7</sup> conjugate addition,<sup>8</sup> and especially for the stereoselective generation of acyclic substituted alkenes in either (*Z*) or (*E*)-isomeric forms.<sup>9</sup>

Whilst numerous methods have been developed towards  $\alpha,\beta$ -unsaturated esters,<sup>10–13</sup> configuration-retentive transition-metal catalyzed (TMC) cross-coupling of alkenyl (pseudo)halides is universally recognized as one of the most practical methodologies.<sup>14</sup> Among the known non-classical pseudohalides,<sup>15</sup> diethylphosphoroxyl (DEP) functionality has been proved as a good leaving group in many organic reactions and the corresponding enol phosphates (EPs), possessing high stability and accessibility, were found to participate in various organic transformations.<sup>16</sup> Particularly, EPs have been utilized in many types of TMC coupling reactions including Suzuki–Miyaura, Stille, Negishi, and Heck reactions by cleavage of the enol-linkage affording highly substituted alkenes.<sup>17</sup> However, the EPs-involved (*Z*) and (*E*)-stereocomplementary synthetic method towards  $\alpha,\beta$ -unsaturated esters with sufficient substrate

generality is still quite limited at present. The latest impressive approach was reported by Tanabe group, which employed *N*-methylimidazole (NMI)-promoted phosphorylation of  $\beta$ -ketoesters to obtain (*Z*) and (*E*)- $\alpha,\beta$ -unsaturated esters, but which suffers from pre-activation of the unstable diphenyl phosphorochloridate (DPPCl) and usage of strong metallic *tert*-butoxide bases.<sup>18</sup> Based on our recent progress in regioselective solvent-free synthesis of EPs,<sup>19</sup> we envisioned that phosphoroxylated (*Z*) and/or (*E*)- $\alpha,\beta$ -unsaturated esters may act as the universal synthon of  $\alpha,\beta$ -unsaturated esters and should be facily obtained from the commercially available and low-cost chloroacetoacetates and phosphites via a simple metal-free Perkow reaction. Herein, we wish to present a stereoselective one-pot synthetic approach towards  $\beta$ -phosphoroxylated (*E*)- $\alpha,\beta$ -unsaturated esters, which are subsequently converted into the corresponding disubstituted  $\alpha,\beta$ -unsaturated esters by Negishi cross-coupling (Scheme 1).





Fig. 1 Selected bioactive  $\alpha,\beta$ -unsaturated carbonyl motifs.

State Key Laboratory of NBC Protection for Civilian, Beijing 102205, P. R. China.  
E-mail: junhua@pku.edu.cn; exshi@sina.com; xiaojingbimail@yeah.net



† Electronic supplementary information (ESI) available: Full experimental details and analytical data. CCDC 2250165. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d3ra02430j>



## Tanabe's work



## This work


Scheme 1 *E*-Stereoselective synthesis of  $\alpha,\beta$ -unsaturated esters from enol phosphates.

Scheme 2 Perkow reaction of phosphite with chloroacetoacetate.

Since both 2-chloroacetoacetates and 4-chloroacetoacetates are capable of undergoing Perkow reaction with phosphites, we then took them together for comparison. Solvent-free Perkow reaction conditions were initially selected in view of high

regioselectivity.<sup>19</sup> As shown in Scheme 2, reaction between  $(\text{EtO})_3\text{P}$  and 2-chloroacetoacetate **2a** gave a mixture of (*E*) and (*Z*)-isomers of  $\beta$ -phosphoroxylated  $\alpha,\beta$ -unsaturated ester **4a** in ratio of 2.6 : 1, whereas reaction between  $(\text{EtO})_3\text{P}$  and 4-chloroacetoacetate **3a** gave the  $\beta$ -phosphoroxylated allylic ester **5a** as the only product. In other words, only moderate *E/Z*-stereoselectivity can be achieved if using 2-chloroacetoacetate, while no conjugated EP product can be obtained if using 4-chloroacetoacetate. However, according to Seeman's report that bases, such as NaH, are supposed to be able to promote 1,3-hydrogen relocation of allyl compounds, we then suspect that the unconjugated EP product **5a** may be able to be transformed into the conjugated one in a stereoselective way.<sup>20</sup>

Inspired by the above idea, we then turned to examine the possibility of the base-promoted 1,3-hydrogen rearrangement of **5a**. As shown in Table 1, among the eight kinds of bases examined, including inorganic *t*-BuOK,  $\text{CH}_3\text{ONa}$ , NaOH, NaH,

Table 1 Optimization of base-promoted 1,3-hydrogen rearrangement of unconjugated  $\beta$ -phosphoroxylated allylic ester **5a**<sup>a</sup>

Entry	Base	Load (x eq.)	Solvent	<i>T</i> (°C)	Time (h)	Yield <sup>b</sup> (%)	<i>E/Z</i> ( <b>4a</b> ) <sup>c</sup>
1	<i>t</i> -BuOK	1.2	THF	rt	24	0	—
2	$\text{CH}_3\text{ONa}$	1.2	THF	rt	24	0	—
3	NaOH	1.2	THF	rt	24	0	—
4	NaH	1.2	THF	rt	24	0	—
5	$\text{K}_2\text{CO}_3$	1.2	THF	rt	24	0	—
6	$\text{Et}_3\text{N}$	1.2	THF	rt	24	90	>99 : 1
7	Pyridine	1.2	THF	rt	24	0	—
8	$(i\text{-Pr})_2\text{NEt}$	1.2	THF	rt	24	20	>99 : 1
9	$\text{Et}_3\text{N}$	1.2	$\text{CH}_3\text{CN}$	rt	4	92	>99 : 1
10	$\text{Et}_3\text{N}$	1.2	DCM	rt	20	90	>99 : 1
11	$\text{Et}_3\text{N}$	1.2	$\text{CH}_3\text{OH}$	rt	22	83	>99 : 1
12	$\text{Et}_3\text{N}$	1.2	DMF	rt	24	75	>99 : 1
13	$\text{Et}_3\text{N}$	0.5	$\text{CH}_3\text{CN}$	rt	7	92	>99 : 1
14	$\text{Et}_3\text{N}$	0.1	$\text{CH}_3\text{CN}$	rt	12	92	>99 : 1
15	$\text{Et}_3\text{N}$	0.05	$\text{CH}_3\text{CN}$	rt	20	93	>99 : 1
16	$\text{Et}_3\text{N}$	0.1	$\text{CH}_3\text{CN}$	0	24	95	>99 : 1
17	$\text{Et}_3\text{N}$	0.1	$\text{CH}_3\text{CN}$	80	4	90	>99 : 1


<sup>a</sup> Reaction conditions: **5a** (1.0 equiv.), base (*x* equiv.), solvent (3 ml). <sup>b</sup> Isolated yields. <sup>c</sup> Determined by NMR.



Having identified the optimal reaction conditions, we next set out to examine the scope of this new mild one-pot enol phosphorylation procedure (Table 2). As for the different *O*-alkyl

$$\text{R}^4\text{O}-\text{P}(\text{R}^2)(\text{R}^3) \xrightarrow[\text{one-pot}]{\begin{array}{l} 1. \text{ 3 (1 eq) / 40 }^\circ\text{C, neat;} \\ 2. \text{ Et}_3\text{N (0.1 eq) / rt, CH}_3\text{CN} \end{array}} \text{R}^2\text{P}(\text{R}^3)(\text{OCH}=\text{C}(\text{CH}_3)\text{C}(=\text{O})\text{OR}^1) \quad \mathbf{4}$$

*E*-stereospecific  
 High yields  
 Mild conditions  
 Metal-free  
 Gram scale  
 Group tolerance

EtO-P(=O)(EtO)-O-CH=C(CH<sub>3</sub>)-C(=O)OEt **4a**, 92% yield  
*i*-PrO-P(=O)(*i*-PrO)-O-CH=C(CH<sub>3</sub>)-C(=O)OEt **4b**, 96% yield  
*n*-BuO-P(=O)(*n*-BuO)-O-CH=C(CH<sub>3</sub>)-C(=O)OEt **4c**, 90% yield  
 Et<sub>2</sub>N-P(=O)(*i*-PrO)-O-CH=C(CH<sub>3</sub>)-C(=O)OEt **4d**, 90% yield  
 Ph-P(=O)(MeO)-O-CH=C(CH<sub>3</sub>)-C(=O)OEt **4e**, 96% yield  
 Ph-P(=O)(Ph)-O-CH=C(CH<sub>3</sub>)-C(=O)OEt **4f**, 90% yield  
 MeO-P(=O)(MeO)-O-CH=C(CH<sub>3</sub>)-C(=O)OEt **4g**, 90% yield  
 EtO-P(=O)(EtO)-O-CH=C(CH<sub>3</sub>)-C(=O)OMe **4h**, 92% yield  
 **4f**  
 EtO-P(=O)(EtO)-O-CH=C(CH<sub>3</sub>)-C(=O)OP*i*-Pr **4i**, 95% yield  
 EtO-P(=O)(EtO)-O-CH=C(CH<sub>3</sub>)-C(=O)OBu-*t* **4j**, 93% yield  
 EtO-P(=O)(EtO)-O-CH=C(CH<sub>3</sub>)-C(=O)OBn **4k**, 94% yield

With the *E*-stereospecific  $\beta$ -phosphoroxyolated  $\alpha,\beta$ -unsaturated esters in hand, we then investigated their stereoretentive Negishi cross-coupling to prepare the corresponding *E*-stereo-defined disubstituted  $\alpha,\beta$ -unsaturated esters. Among the typical catalysts screened including  $\text{Pd}(\text{PPh}_3)_4$ ,  $\text{Ni}(\text{acac})_2$  and  $\text{Pd}(\text{dppb})\text{Cl}_2$ , the latter demonstrated the best performance in this Negishi reaction with only 0.02 equivalent loading by refluxing in acetonitrile. Various aromatic  $\text{ArZnCl}$  nucleophiles containing electron-donating and/or electron-withdrawing substituents at *ortho*, *meta*, and/or *para* positions were all tolerated well, affording the desired products in good to excellent yields (80–

CC(=O)OCCOP(=O)(OCC)OCC + ArMgBr >> CC(=O)OCC=C(Ar)C
  
 ZnCl<sub>2</sub>, Pd(dppb)Cl<sub>2</sub>, reflux in MeCN

92% yield; 50 mmol scale
 100% *E*-stereorentative

---

**6a**, 84%
 **6b**, 80%
**6c**, 93%

**6d**, 88%
 **6e**, 94%
**6f**, 92%

**6g**, 88%
 **6h**, 95%
**6i**, 90%

**6j**, 96%
 **6k**, 82%
**6l**, 87%

**6m**, 94%
 **6n**, 85%
**6o**, 90%

**6p**, 92%
 **6q**, 89%
**6r**, 85%

<sup>a</sup> Reaction conditions: **4a** (1.0 mmol), ArMgBr (1.5 mmol), ZnCl<sub>2</sub> (1.5 mmol), Pd(dppb)Cl<sub>2</sub> (0.02 mmol), CH<sub>3</sub>CN (5.0 mL), reflux about 3 h.  
<sup>b</sup> Isolated yields.


96%) without generating any stereochemical integrity (Table 3, **6a–6m**). Disubstituted, condensed and hetero aromatic organometallic substrates also gave 85–92% yields of the products (Table 3, **6n–6r**). However, it's regrettable that alkyl organozinc reagents was found unreactive under such conditions.

Furthermore, under the above optimal Negishi cross-coupling reaction conditions, both (*Z*) and (*E*) isomers of  $\alpha,\beta$ -unsaturated esters **6a** could be easily achieved, just by one operation, directly from the (*Z*) and (*E*) mixture of **4a** in 22% and 70% yields respectively (Scheme 3).


According to the Cram's mechanistic interpretation for the allylic rearrangements, an intra-molecular pathway of the  $\text{Et}_3\text{N}$ -promoted stereoselective 1,3-hydrogen rearrangement of the EPs **5a** was proposed because that the degree of the observed intramolecularity depended strongly on the base and solvent used.<sup>21</sup> As shown in Scheme 4, triethylamine firstly removes the proton from the  $\alpha$ -carbon position of ester **5a**, resulting in a coplanar anionic allylic system by three carbon atoms. The hydrogen atom of the  $\text{H-Et}_3\text{N}$  ammonium then bonds to both terminal carbon atoms to form the intermediate **Int**, collapse of which would then give the thermodynamically favourable conjugated  $\alpha,\beta$ -unsaturated ester product *E*-**4a**.

In summary, a mild and environmental trimethylamine-catalyzed *E*-stereoselective 1,3-hydrogen allylic rearrangement of enol phosphates was firstly developed to afford versatile  $\beta$ -phosphoroxylated (*E*)- $\alpha,\beta$ -unsaturated esters which can be then efficiently converted into the corresponding  $\beta,\beta$ -disubstituted (*E*)- $\alpha,\beta$ -unsaturated esters in high yields by a 100% stereoretentive Negishi cross-coupling reaction. Moreover, both (*Z*) and (*E*)- $\alpha,\beta$ -unsaturated esters were able to be achieved in one manipulation when just employing 2-chloroacetoacetate instead of 4-chloroacetoacetate for the solvent and metal-free Perkow reaction.

It is interesting to note that more structure-diverse  $\alpha,\beta$ -unsaturated esters should be easily obtained by derivation reactions at the allylic position of  $\alpha,\beta$ -unsaturated esters and/or by utilizing 2-substituted 4-chloroacetoacetates as the starting materials.



Scheme 3 Preparation of (*Z*) and (*E*) isomers of **6a** in one operation.



Scheme 4 Proposed (*E*)-stereospecific allylic rearrangement mechanism.

## Conflicts of interest

There are no conflicts to declare.

## Notes and references

- (a) C. X. Zhuo and A. Furstner, *J. Am. Chem. Soc.*, 2018, **140**, 10514; (b) D. Romo, N. S. Choi, S. Li, I. Buchler, Z. Shi and J. O. Liu, *J. Am. Chem. Soc.*, 2004, **126**, 10582; (c) S. D. Marco, A. Cammas, J. Pelletier and I. E. Gallouzi, *Nat. Commun.*, 2012, **3**, 8963; (d) S. K. Naineni, J. Liang, B. Nagar and J. Pelletier, *Cell Chem. Biol.*, 2021, **28**, 825.
- (a) J. K. Nunnery, N. Engene, T. Byrum, T. F. Murray and W. H. Gerwick, *J. Org. Chem.*, 2012, **77**, 4198; (b) S. Sekharan and K. Morokuma, *J. Am. Chem. Soc.*, 2011, **133**, 19052; (c) E. A. Zhukovsky, P. R. Robinson and D. D. Oprian, *Science*, 1990, **251**, 558.
- M. Kiser and K. Golczak, *Chem. Rev.*, 2014, **114**, 194.
- (a) P. Guo, Y. Zhang, L. Zhang and Q. Xia, *J. Biol. Chem.*, 2021, **297**, 101234; (b) Y. Ando, K. Matsumoto, K. Misaki, G. Mano, T. Shinada and S. G. Goto, *Gen. Comp. Endocrinol.*, 2020, **289**, 113394; (c) M. Nouzova, C. Rivera-Pérez and F. G. Noriega, *Curr. Opin. Insect. Sci.*, 2018, **29**, 49; (d) K. Li, Q. Jia and S. Li, *Insect Sci.*, 2019, **26**, 600.
- (a) D. A. Evans, P. J. Coleman, L. C. Dias and A. N. Tyler, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 2744; (b) D. A. Evans, D. M. Fitch, T. E. Smith and V. Cee, *J. Am. Chem. Soc.*, 2000, **122**, 10033; (c) D. A. Evans, P. H. Carter, E. M. Carreira, A. B. Charette, J. A. Prunet and M. Lautens, *J. Am. Chem. Soc.*, 1999, **121**, 7540; (d) I. Fleming, A. Barbero and D. Walter, *Chem. Rev.*, 1997, **97**, 2063.
- (a) E. Negishi, Q. Hu, Z. Huang, M. Qian and G. Wang, *Aldrichimica Acta*, 2005, **38**, 71; (b) J. Li, A. S. Grillo and M. D. Burke, *Acc. Chem. Res.*, 2015, **48**, 2297; (c) V. Hornillos, M. Giannerini, C. Vila, M. F. Mastral and B. L. Feringa, *Chem. Sci.*, 2015, **6**, 1394.
- (a) K. Murakami and H. Yorimitsu, *Beilstein J. Org. Chem.*, 2013, **9**, 278; (b) M. G. Suero, E. D. Bayle, B. S. L. Collins and M. J. Gaunt, *J. Am. Chem. Soc.*, 2013, **135**, 5332; (c) F. Xue, J. Zhao, T. S. A. Hor and T. Hayashi, *J. Am. Chem. Soc.*, 2015, **137**, 3189; (d) S. Wang and C. Xi, *Org. Lett.*, 2018, **20**, 4131.
- (a) F. Guibe, *Tetrahedron*, 1998, **54**, 2967; (b) F. Guibe, *Tetrahedron*, 1997, **53**, 13509; (c) S. Escoubet, S. Gastaldi and M. Bertrand, *Eur. J. Org. Chem.*, 2005, 3855.
- (a) F. W. Sum and L. Weiler, *Can. J. Chem.*, 1979, **57**, 1431; (b) M. Ide and M. Nakata, *Synlett*, 2001, 1511.
- (a) A. B. Flynn and W. W. Ogilvie, *Chem. Rev.*, 2007, **107**, 4698; (b) P. Polak, H. Vanova, D. Dvorak and T. Tobrman, *Tetrahedron Lett.*, 2016, **57**, 3684; (c) B. E. Maryanoff and A. B. Reitz, *Chem. Rev.*, 1989, **89**, 863; (d) Y. Ashida and Y. Tanabe, *Chem. Rec.*, 2020, **20**, 1.
- (a) C. Gürtler and S. L. Buckwald, *Chem. - Eur. J.*, 1999, **5**, 3107; (b) M. Shindo, Y. Sato, T. Yoshikawa, R. Koretsune and K. Shishido, *J. Org. Chem.*, 2004, **69**, 3912.
- (a) A. B. Lemay, K. S. Vulic and W. W. Ogilvie, *J. Org. Chem.*, 2006, **71**, 3615; (b) J. S. Mercier, A. B. Flynn and W. W. Ogilvie, *Tetrahedron*, 2008, **64**, 5472.



- 13 (a) Y. Yamamoto, N. Kirai and Y. Harada, *Chem. Commun.*, 2008, 2010; (b) N. Kirai and Y. Yamamoto, *Org. Synth.*, 2010, **87**, 53; (c) Z. He, S. Kirchberg, R. Froehlich and A. Studer, *Angew. Chem., Int. Ed.*, 2012, **124**, 3759.
- 14 (a) J. M. Baxter, D. Steinhuebel, M. Palucki and I. W. Davies, *Org. Lett.*, 2005, **7**, 215; (b) A. Klapars, K. R. Campos, C. Chen and R. P. Volante, *Org. Lett.*, 2005, **7**, 1185; (c) H. Nakatsuji, K. Ueno, T. Misaki and Y. Tanabe, *Org. Lett.*, 2008, **10**, 2131; (d) H. Nakatsuji, H. Nishikado, K. Ueno and Y. Tanabe, *Org. Lett.*, 2009, **11**, 4258.
- 15 Y. Ashida, K. Nakata, D. Yoshitake, Y. Sato, Y. Miyazaki and Y. Tanabe, *Asian J. Org. Chem.*, 2020, **9**, 604.
- 16 (a) A. L. Hansen, J. P. Ebran, M. Ahlquist, P. O. Norrby and T. Skrydstrup, *Angew. Chem., Int. Ed.*, 2006, **45**, 3349; (b) T. Hayashi, T. Fujiwa, Y. Okamoto, Y. Katsuro and M. Kumada, *Synthesis*, 1981, 1001; (c) K. C. Nicolaou, G. Q. Shi, G. P. Grtner and Z. Yang, *J. Am. Chem. Soc.*, 1997, **119**, 5467; (d) A. S. E. Karlström, K. Itami and J. E. Bckvall, *J. Org. Chem.*, 1999, **64**, 1745; (e) J. Wu and Z. Yang, *J. Org. Chem.*, 2001, **66**, 7875; (f) J. P. Ebran, A. L. Hansen, T. M. Gøgsig and T. Skrydstrup, *J. Am. Chem. Soc.*, 2007, **129**, 6931; (g) A. L. Hansen, J.-P. Ebran, T. M. Gøgsig and T. Skrydstrup, *J. Org. Chem.*, 2007, **72**, 6464; (h) W. You, Y. Li and M. K. Brown, *Org. Lett.*, 2013, **15**, 1610.
- 17 (a) T. M. Gøgsig, A. T. Lindhardt and T. Skrydstrup, *Org. Lett.*, 2009, **11**, 4886; (b) J. Jiang, R. DeVita, G. Doss, M. Goulet and M. Wyvratt, *J. Am. Chem. Soc.*, 1999, **121**, 593; (c) A. L. Hansen, J. P. Ebran, T. M. Gøgsig and T. Skrydstrup, *Chem. Commun.*, 2006, **39**, 4137; (d) J. W. Coe, *Org. Lett.*, 2000, **2**, 4205.
- 18 H. Nakatsuji, Y. Ashida, H. Hori, Y. Sato, M. Taira and Y. Tanabe, *Org. Biomol. Chem.*, 2015, **13**, 8205.
- 19 (a) Y. Cao, Z. Gao, J. Li, X. Bi, L. Yuan, C. Pei, Y. Guo and E. Shi, *RSC Adv.*, 2020, **10**, 29493; (b) H. Guo, Y. Zhang, Z. Li, P. Zhao, N. Li and E. Shi, *RSC Adv.*, 2022, **12**, 14844.
- 20 S. G. Alcock, J. E. Baldwin, R. Bohlmann, L. M. Harwood and J. Seeman, *J. Org. Chem.*, 1985, **50**, 3526.
- 21 (a) M. Hassam, A. Taher, G. E. Arnott, I. R. Green and W. A. L. van Otterlo, *Chem. Rev.*, 2015, **115**, 5462; (b) D. J. Cram and R. T. Uyeda, *J. Am. Chem. Soc.*, 1964, **86**, 5466.

