## **RSC** Advances



View Article Online

View Journal | View Issue

## PAPER

Check for updates

Cite this: RSC Adv., 2023, 13, 19091

Received 27th April 2023 Accepted 7th June 2023 DOI: 10.1039/d3ra02791k

rsc.li/rsc-advances

## 1 Introduction

Mercury has attracted attention from researchers owing to its strong toxicity and bioaccumulation.<sup>1-4</sup> With the development of industry, especially gold mining, the burning of fossil and oil refining, mercury ion pollution is widely distributed in the environment.<sup>5-7</sup> Through biological enrichment, mercury ions in water can be transformed into organic mercury ions with stronger toxicity and then enter the human body *via* the food chain.<sup>8-10</sup> Even very small amounts of mercury ions have a severe impact on the human body, including the digestive system and kidneys, cognitive disorders, and even central nervous system damage.<sup>11-16</sup> Consequently, the development of methods for the detection of mercury ions with simple synthesis and high selectivity and sensitivity is significant.

Recently, many methods have been reported for the detection of mercury ions (Hg<sup>2+</sup>), including gas chromatography, inductive coupled plasma mass spectroscopy, and atomic absorption spectrometry.<sup>17–21</sup> However, most of the above-mentioned methods have some disadvantages, such as complex pretreatment, use of expensive instruments, and difficulty in realizing real-time and on-site monitoring.<sup>22,23</sup> In contrast, the use of fluorescent probes has attracted significant attention due to their simple operation and high selectivity and sensitivity.<sup>24–30</sup> Thus, an increasing number of fluorescent probes has been used to detect

# Highly selective fluorescent probe for detecting mercury ions in water<sup>†</sup>

Yanfeng Shi, <sup>(b)</sup> <sup>a</sup> Bingxu Li,<sup>a</sup> Zhifeng Wang, <sup>(b)</sup> <sup>a</sup> Yanhao Zhang,<sup>a</sup> Zhibin Zhang,<sup>a</sup> Xu Zhang<sup>\*b</sup> and Fulin Li<sup>\*c</sup>

Mercury ion  $(Hg^{2+})$  is a well-known toxic heavy metal. It has become one of the most significant environmental pollutants in the world because of its serious physiological toxicity, persistence, easy migration, and high bioconcentration. Thus, the development of methods for monitoring  $Hg^{2+}$  is indispensable. Herein, we have designed and synthesized a new fluorescent probe, TPH, for the detection of  $Hg^{2+}$  in the water environment. The TPH probe could quantitatively detect  $Hg^{2+}$  between 0 and 5  $\mu$ M (LOD = 16 nM), with a linear range of 0–2.5  $\mu$ M. In addition, the TPH probe was used to monitor  $Hg^{2+}$  in water samples successfully. Thus, this probe is suitable for monitoring  $Hg^{2+}$  in the actual water environment.

environmental heavy metal pollutants including Hg<sup>2+</sup>. However, the reported probes for monitoring Hg<sup>2+</sup> still have some short-comings, such as poor selectivity and water solubility and high detection limits (LODs).<sup>31–35</sup> Therefore, new fluorescent probes need to be developed for monitoring Hg<sup>2+</sup> in the environment with excellent selectivity and sensitivity and good water solubility.

Accordingly, herein, we synthesized the **TPH** probe, which was based on the TPC-OH dye as the fluorophore<sup>36</sup> and phenyl thiochloroformate as the recognition receptor of  $Hg^{2^+}$ .<sup>37,38</sup> Phenyl thiochloroformate possesses high selectivity for the detection of  $Hg^{2^+}$ , and thus the **TPH** probe could also achieve the specific and sensitive detection of  $Hg^{2^+}$ . The **TPH** probe exhibited the following excellent properties: (1) good water solubility, (2) excellent sensitivity (LOD = 16 nM), (3) high selectivity, and (4) excellent application in the environment. Thus, this probe will have a wide application prospect for monitoring  $Hg^{2^+}$  in the environment.

## 2 Experimental

#### 2.1 Materials and instruments

All chemical reagents were obtained from commercial sources and used without further purification. Absorption and fluorescence spectra were recorded on a UV-3101PC spectro-photometer and Horiba FluoroMax-4 spectrophotometer, respectively.

#### 2.2 Synthesis of TPH probe

TPC-OH dye (236 mg, 1 mmol) was dissolved in dry  $CH_2Cl_2$  (15 mL), and then phenyl thiochloroformate (259 mg, 1.5 mmol) and *N*,*N*-diisopropylethylamine (DIPEA) (194 mg, 1.5 mmol) were added (Scheme 1). The mixed solution was stirred at 25 °C for 2 h.

<sup>&</sup>quot;School of Municipal and Environmental Engineering, Shandong Jianzhu University, Jinan, China

<sup>&</sup>lt;sup>b</sup>School of Architecture and Urban Planning, Shandong Jianzhu University, Jinan, China. E-mail: 15668303582@163.com; Tel: +86 156 6830 3582

<sup>&</sup>lt;sup>c</sup>Water Resources Research Institute of Shandong Province, Jinan, China. E-mail: fulinli@126.com

<sup>†</sup> Electronic supplementary information (ESI) available. See DOI: https://doi.org/10.1039/d3ra02791k



Scheme 1 Synthesis of TPH probe.



Fig. 1 (a) Fluorescence and (b) absorption spectrum changes of TPH probe (5  $\mu$ M and 20  $\mu$ M, respectively) before and after the addition of Hg<sup>2+</sup> (20  $\mu$ M).  $\lambda_{ex} = 400$  nm and  $\lambda_{em} = 505$  nm. Conditions: HEPES (5 mM, pH 7.4).

The crude product was purified by column silica chromatography over silica gel using dichloromethane/petroleum ether (14:5) as the eluent to provide a faint-yellow pure solid product. <sup>1</sup>H NMR (400 MHz, DMSO)  $\delta$  (ppm): 7.94 (d, J = 5.6 Hz, 1H), 7.81 (d, J = 8.0 Hz, 2H), 7.68 (s, 1H), 7.58 (s, 1H), 7.55–5.52 (m, 4H), 7.48 (t, J = 8.0 Hz, 2H), 7.39 (t, J = 8.0 Hz, 3H), 4.21 (s, 2H). <sup>13</sup>C NMR (100 MHz, DMSO)  $\delta$  (ppm): 193.95, 192.55, 158.01, 153.62, 152.68, 136.23, 135.34, 135.21, 133.66, 131.32, 130.46, 130.42, 129.52, 127.57, 125.82, 122.74, 122.20, 120.54, 32.44.

## 3 Results and discussion

#### 3.1 Spectral response of TPH probe

All the reactions were carried out in aqueous solution (HEPES 5 mM, pH = 7.4). The fluorescence spectra of the **TPH** probe for monitoring  $Hg^{2+}$  was investigated. When  $Hg^{2+}$  (20  $\mu$ M) was added, the fluorescence intensity displayed a significant enhancement at 505 nm (Fig. 1a). The quantum yield of the **TPH** probe was calculated to be 0.07. Then, its absorption spectrum was also studied. According to the results, the absorption peak changed from 325 nm to 350 nm (Fig. 1b), implying that  $Hg^{2+}$  could promote the splitting of the carbonothioate moiety (Scheme 2). Furthermore, we conducted HRMS and NMR to explore the reaction mechanism of the **TPH** probe and  $Hg^{2+}$  (Fig. S1–S4 in the ESI†).

#### 3.2 Quantification of Hg<sup>2+</sup>

The **TPH** probe showed good water solubility, and thus the influence of the concentration of  $Hg^{2+}$  on its fluorescence intensity in pure water was investigated. With an increase in  $Hg^{2+}$  concentration (0–5 µM), the fluorescence intensity of the **TPH** probe at 505 nm increased accordingly (Fig. 2a). In addition, when the concentration of  $Hg^{2+}$  was 0–2.5 µM, it was linearly correlated with the fluorescence intensity (y = 201 921 [ $Hg^{2+}$ ] (µM) + 85 603,  $R^2 = 0.9828$ ) (Fig. 2b), and the LOD was 16 nM ( $3\sigma/k$ ). Thus, all the above-mentioned results indicate



Scheme 2 Recognition mechanism of TPH probe for Hg<sup>2+</sup>



Fig. 2 (a) Fluorescence spectra of TPH probe (5  $\mu$ M) for Hg<sup>2+</sup> (0–5  $\mu$ M). (b) Linear plot of fluorescence intensity (505 nm) to Hg<sup>2+</sup> (0–2.5  $\mu$ M).  $\lambda_{ex}=400$  nm and  $\lambda_{em}=505$  nm. Conditions: in HEPES (5 mM, pH 7.4).

that the **TPH** probe can provide a sensitive detection tool for  $Hg^{2+}$  in the actual water environment.

#### 3.3 Specificity for Hg<sup>2+</sup>

The specificity of the **TPH** probe toward  $Hg^{2+}$  and other various relevant analytes including  $Cd^{2+}$ ,  $Cu^{2+}$ ,  $Fe^{2+}$ ,  $Ca^{2+}$ ,  $K^+$ ,  $Mg^{2+}$ ,  $Na^+$ ,  $Ni^{2+}$ ,  $Pb^{2+}$ ,  $Sn^{2+}$ ,  $Zn^{2+}$ ,  $Co^{2+}$ ,  $Fe^{3+}$ ,  $NO^{3-}$ ,  $SO_4^{-2-}$  and  $Cl^-$  was analyzed. The concentration of  $Hg^{2+}$  and the other



Fig. 3 (a) Fluorescence response of TPH probe (5  $\mu$ M) toward Hg<sup>2+</sup> (5  $\mu$ M) and other ions (50  $\mu$ M). (b) Fluorescence response of TPH probe (5  $\mu$ M) toward Hg<sup>2+</sup> (5  $\mu$ M) in the presence of other ions (50  $\mu$ M).  $\lambda_{ex} =$  400 nm and  $\lambda_{em} =$  505 nm. Conditions: in HEPES (5 mM, pH 7.4).

analytes was 5  $\mu$ M and 50  $\mu$ M. Only Hg<sup>2+</sup> caused a fluorescence response at 505 nm, while the other relevant analytes did not cause obvious fluorescence changes (Fig. 3a). Besides, interference experiments were also conducted. The fluorescence intensity response values exhibited a slight change at 505 nm (Fig. 3b). Thus, all these results strongly suggest that the **TPH** probe can specifically recognize Hg<sup>2+</sup>.

#### 3.4 Analytical applications in real water samples

Then, the analytical application of the **TPH** probe in three water samples (lake water, underground water and river water) for the

Table 1         Application of TPH probe in three water samples"					
Real water samples	Found Hg <sup>2+</sup>	Addition $Hg^{2+}(\mu M)$	Found (µM)	Recovery (%)	RSD $(n = 3)$ (%)
Sample A	ND	1	$0.97\pm0.03$	96.93	3.08
		2	$2.01\pm0.13$	100.38	6.40
Sample B	ND	1	$1.09\pm0.06$	109.20	6.42
		2	$1.68\pm0.13$	84.07	6.64
Sample C	ND	1	$0.96\pm0.08$	95.97	8.41
		2	$1.72\pm0.11$	85.81	5.40

<sup>a</sup> ND: not detected. Sample A from JiaZi Lake, University of Jinan and samples B and C from Jinyun River and Jinyang River in Jinan, China.

detection of Hg<sup>2+</sup> was investigated. Firstly, no Hg<sup>2+</sup> was found in the samples. Then, after 5  $\mu$ M **TPH** probe was added to the test water samples, 1 and 2  $\mu$ M Hg<sup>2+</sup> were also respectively added. Each sample was tested three times. As can be seen in Table 1, the recoveries of the three water samples were 84.07–109.20%, further confirming that this newly synthesized probe could effectively detect Hg<sup>2+</sup> in the real water environment.

## 4 Conclusions

The fluorescent **TPH** probe with phenyl thiochloroformate as the  $Hg^{2+}$  recognition site was synthesized in this study. This probe could specifically recognize  $Hg^{2+}$  and quantitatively detect  $Hg^{2+}$  in aqueous solution. According to the experimental results, we calculated that its detection limit is 16 nM. Meanwhile, the **TPH** probe has excellent water solubility, which is conducive for its application in the actual environment.

## Data availability

The datasets generated and analyzed during the current study are available from the corresponding author on reasonable request.

## Conflicts of interest

The authors have no conflicts of interest to declare.

## Acknowledgements

We gratefully acknowledge financial support from the NSFC (42007176, 52070122), and Natural Science Foundation of Shandong Province (ZR2020QD125, ZR2020ME222), Key Scientific and Technological Innovation Foundation of Shandong Province (No. 2020CXGC011404).

## References

- 1 S. Y. Chen, Z. Li, K. Li and X. Q. Yu, *Coord. Chem. Rev.*, 2021, **429**, 213691.
- 2 M. A. Peckham, M. S. Gustin and P. J. Weisberg, *Environ. Sci. Technol.*, 2019, **53**, 3663.
- 3 L. Wang, Y. Zhao and J. Zhang, *Ind. Eng. Chem. Res.*, 2017, 56, 3804.
- 4 S. X. Wang, L. Zhang, L. Wang, Q. R. Wu, F. Y. Wang and J. M. Hao, *Front. Environ. Sci. Eng.*, 2014, **8**, 631.
- 5 Y. M. Yu, C. Y. Liu, B. Tian, X. Y. Cai, H. C. Zhu, P. Jia, Z. L. Li, X. Zhang, W. L. Sheng and B. C. Zhu, *Dyes Pigm.*, 2020, 177, 108290.
- 6 X. Yuan, T. H. Leng, Z. Q. Guo, C. Y. Wang, J. Z. Li, W. W. Yang and W. H. Zhu, *Dyes Pigm.*, 2019, **161**, 403.
- 7 Q. X. Duan, M. Zhang, C. X. Sheng, C. Y. Liu, L. Wu, Z. M. Ma, Q. Zhao, Z. P. Wang and B. C. Zhu, *Anal. Sci.*, 2017, 33, 1169.
- 8 F. Song, C. Yang, X. T. Shao, L. Du, J. Zhu and C. Kan, *Dyes Pigm.*, 2019, **165**, 444.

- 9 C. Y. Qin, B. Y. Du, R. S. Yin, B. Meng, X. W. Fu, P. Li, L. M. Zhang and X. B. Feng, *Environ. Sci. Technol.*, 2020, 54, 14334.
- 10 Y. Zhang, C. Y. Liu, M. J. Su, X. D. Rong, X. Wang, K. Wang, X. W. Li, H. C. Zhu, M. H. Yu, W. L. Sheng and B. C. Zhu, *J. Photochem. Photobiol.*, A, 2022, 425, 113706.
- 11 X. Wang and W. X. Wang, Environ. Sci. Technol., 2015, 49, 10173.
- 12 J. Xu, Z. K. Wang, C. Y. Liu, Z. H. Xu, N. Wang, X. Cong and B. C. Zhu, *J. Lumin.*, 2018, **6**, 1122.
- 13 R. Roy, S. Rakshit, S. Bhar and S. Bhattacharya, *RSC Adv.*, 2015, **5**, 67833.
- 14 M. Jing, D. Lin, P. Wu, M. J. Kainz, K. Bishop, H. Yan, Q. Li and X. Feng, *Sci. Total Environ.*, 2021, **782**, 146410.
- 15 P. Srivastava, S. Razi, R. Ali, R. Gupta, S. Yadav, G. Narayan and A. Misra, *Anal. Chem.*, 2014, **86**, 8693.
- 16 B. Yuan, D. X. Wang, L. N. Zhu, Y. L. Lan, M. Cheng, L. M. Zhang, J. Q. Chu and X. Z. Li, *Chem. Sci.*, 2019, **10**, 4220.
- 17 Z. Mermer, O. Yavuz, S. K. Atasen, Y. Alcay and I. Yilmaz, *J. Hazard. Mater.*, 2021, **410**, 124597.
- 18 A. Thongsaw, R. Sananmuang, Y. Udnan, G. M. Ross and W. C. Chaiyasith, Spectrochim. Acta, Part B, 2019, 160, 105685.
- 19 Q. X. Zhou, Y. L. Liu, Y. L. Wu, Z. Li, Y. H. Li, M. H. Liu, T. X. Qu and C. M. Chen, *Chemosphere*, 2021, 274, 129959.
- 20 S. E. Long, J. E. Norris, J. Carney and J. V. Ryan, *Pollut. Res.*, 2020, **11**, 909.
- 21 I. Narin, M. Soylak, L. Elci and M. Dogan, *Talanta*, 2000, **52**, 1041.
- 22 S. Huang, X. Cheng, Q. Lei, B. Feng, X. H. Liu, J. P. Ding,
  C. Zhong, J. H. Liang and W. B. Zeng, *Dyes Pigm.*, 2021,
  187, 109125.
- 23 Y. N. Zhang, S. Guo, Z. R. Jiang, G. B. Mao, X. H. Ji and Z. K. He, *Anal. Chem.*, 2018, **90**, 9796.
- 24 T. H. Chen, S. Zhang, M. Jaishi, R. Adhikari, J. Bi, M. Fang, S. Xia, Y. Zhang, R. L. Luck, R. Pati, H. M. Lee, F. T. Luo, A. Tiwari and H. Liu, *ACS Appl. Bio Mater.*, 2018, 1, 549.
- 25 S. Y. Ma, L. Y. Li, M. Y. She, Y. Mo, S. Y. Zhang, P. Liu and J. L. Li, *Chin. Chem. Lett.*, 2017, 28, 2014.
- 26 Q. X. Duan, H. C. Zhu, C. Y. Liu, R. F. Yuan, Z. T. Fang, Z. K. Wang, P. Jia, Z. L. Li, W. L. Sheng and B. C. Zhu, *Analyst*, 2019, **144**, 1426.
- 27 W. Shu, L. G. Yan, J. Liu, Z. K. Wang, S. Zhang, C. C. Tang,
  C. Y. Liu, B. C. Zhu and B. Du, *Ind. Eng. Chem. Res.*, 2015,
  54, 8056.
- 28 Y. Z. Liu, H. Z. Zhao, N. Gao, C. L. Yang, R. B. Zhang and X. L. Zhang, Sens. Actuators, B, 2021, 344, 130201.
- 29 B. Yuan, D. X. Wang, L. N. Zhu, Y. L. Lan, M. Cheng, L. M. Zhang, J. Q. Chu, X. Z. Li and D. M. Kong, *Chem. Sci.*, 2019, **10**, 4220.
- 30 Q. X. Duan, X. Y. Lv, C. Y. Liu, Z. F. Geng, F. F. Zhang,
  W. L. Sheng, Z. K. Wang, P. Jia, Z. L. Li, H. C. Zhu and
  B. C. Zhu, *Ind. Eng. Chem. Res.*, 2019, 58, 11.
- 31 M. Zhao, G. K. Shao, Y. S. Guo, Y. L. Tang, J. B. Liu and D. S. Guo, *New J. Chem.*, 2020, 44, 12538.
- 32 S. Maity, S. Banerjee, K. Sunwoo, J. Kim and P. Bharadwaj, *Inorg. Chem.*, 2015, **54**, 3929.

#### Paper

- 33 H. Un, C. Huang, C. Huang, T. Jia, X. Zhao, C. Wang, L. Xu and H. Yang, *Org. Chem. Front.*, 2014, **1**, 1083.
- 34 T. Zhang, B. Wu, Z. Zou, Y. Wu, J. Zheng, W. Wong and K. Wong, *J. Lumin.*, 2016, **170**, 187.
- 35 S. Y. Qin, B. Chen, J. Huang and Y. F. Han, *New J. Chem.*, 2018, 42, 12766.
- 36 C. Y. Liu, P. Jia, Z. H. Zhuang, Z. K. Wang, Q. X. Duan, Z. R. Li, H. C Zhu, X. Zhang, B. C. Zhu and W. L. Sheng, *Sens. Actuators, B*, 2019, 291, 243.
- 37 J. Xu, Z. H. Xu, Z. K. Wang, C. Y. Liu, B. C. Zhu, X. R. Wang, K. Wang, J. T. Wang and G. Q. Sang, *J. Lumin.*, 2018, 33, 219.
- 38 W. Shu, Y. W. Wang, L. Wu, Z. K. Wang, Q. X. Duan, Y. B. Gao, C. Y. Liu, B. C. Zhu and L. G. Yan, *Ind. Eng. Chem. Res.*, 2016, 55, 8713.