



Cite this: *RSC Adv.*, 2023, **13**, 30665

Metabolomics and molecular networking approach for exploring the anti-diabetic activity of medicinal plants†

Arjun Prasad Timilsina,^{‡,a} Bimal Kumar Raut,^{‡,a} Chen Huo,^{‡,b} Karan Khadayat,^a Prakriti Budhathoki,^a Mandira Ghimire,^a Rabin Budhathoki,^a Niraj Aryal,^c Ki Hyun Kim ^{*,b} and Niranjan Parajuli ^{*,a}

Metabolomics and molecular networking approaches have expanded rapidly in the field of biological sciences and involve the systematic identification, visualization, and high-throughput characterization of bioactive metabolites in natural products using sophisticated mass spectrometry-based techniques. The popularity of natural products in pharmaceutical therapies has been influenced by medicinal plants with a long history of ethnobotany and a vast collection of bioactive compounds. Here, we selected four medicinal plants *Cleistocalyx operculatus*, *Terminalia chebula*, *Ficus lacor*, and *Ficus semicordata*, the biochemical characteristics of which remain unclear owing to the inherent complexity of their plant metabolites. In this study, we aimed to evaluate the potential of these aforementioned plant extracts in inhibiting the enzymatic activity of α -amylase and α -glucosidase, respectively, followed by the annotation of secondary metabolites. The methanol extract of *Ficus semicordata* exhibited the highest α -amylase inhibition with an IC_{50} of $46.8 \pm 1.8 \mu\text{g mL}^{-1}$, whereas the water fraction of *Terminalia chebula* fruits demonstrated the most significant α -glucosidase inhibition with an IC_{50} value of $1.07 \pm 0.01 \mu\text{g mL}^{-1}$. The metabolic profiling of plant extracts was analyzed through Liquid Chromatography-Mass Spectrometry (LC-HRMS) of the active fractions, resulting in the annotation of 32 secondary metabolites. Furthermore, we applied the Global Natural Product Social Molecular Networking (GNPS) platform to evaluate the MS/MS data of *Terminalia chebula* (bark), revealing that there were 205 and 160 individual ion species observed as nodes in the methanol and ethyl acetate fractions, respectively. Twenty-two metabolites were tentatively identified from the network map, of which 11 compounds were unidentified during manual annotation.

Received 15th June 2023

Accepted 4th October 2023

DOI: 10.1039/d3ra04037b

rsc.li/rsc-advances

^aBiological Chemistry Lab, Central Department of Chemistry, Tribhuvan University, Kirtipur, Kathmandu, 44618, Nepal. E-mail: niranjan.parajuli@cdc.tu.edu.np; Tel: +977-1-4332034

^bSchool of Pharmacy, Sungkyunkwan University, Suwon 16419, Republic of Korea. E-mail: khkim83@skku.edu; Tel: +82-31-290-7700

^cDepartment of Biology, University of Florida, Gainesville, FL 32611, USA

† Electronic supplementary information (ESI) available: Fig. S1: mass spectrum of chebulic acid from *Terminalia chebula*; Fig. S2: mass spectrum of 2-O-caffeyl hydroxyacrylic acid from *Terminalia chebula*; Fig. S3: mass spectrum of ellagic acid from *Terminalia chebula*; Fig. S4: mass spectrum of Trigalloylveoglucosan IX from *Terminalia chebula*; Fig. S5: mass spectrum of 2-O-feruloylhydroxyacrylic acid from *Terminalia chebula*; Fig. S6: mass spectrum of arjunglucoside I from *Terminalia chebula*; Fig. S7: mass spectrum of 23-galloylterminolic acid from *Terminalia chebula*; Fig. S8: mass spectrum of flavogallonic acid from *Terminalia chebula*; Fig. S9: mass spectrum of Leucodelphinidin from *Terminalia chebula*; Fig. S10: mass spectrum of arjunolic acid from *Terminalia chebula*; Fig. S11: mass spectrum of bartogenic acid from *Terminalia chebula*; Fig. S12: mass spectrum of arjungenin from *Terminalia chebula*; Fig. S13: mass spectrum of (–)-epicatechin-3-O-gallate from *Terminalia chebula*; Fig. S14: mass spectrum of gambiriin B1 from

Terminalia chebula; Fig. S15: mass spectrum of rotundioic acid from *Terminalia chebula*; Fig. S16: mass spectrum of butin from *Terminalia chebula*; Fig. S17: mass spectrum of gallocatechin or epigallocatechin from *Cleistocalyx operculatus*; Fig. S18: mass spectrum of 4,4'-di-O-methyllellagic acid from *Cleistocalyx operculatus*; Fig. S19: mass spectrum of isoquercitrin from *Ficus lacor*; Fig. S20: mass spectrum of catechin from *Ficus semicordata*; Fig. S21: mass spectrum of procyanidin B2 from *Ficus semicordata*; Fig. S22: mass spectrum of ellagic acid-O-pentoside from *Cleistocalyx operculatus*; Fig. S23: mass spectrum of quinic acid from *Ficus lacor*; Fig. S24: mass spectrum of hydroxyl-6-gingerol-O- β -D-glucuronide from *Ficus semicordata*; Fig. S25: mass spectrum of 3-O-galloylnorbergenin from *Cleistocalyx operculatus*; Fig. S26: mass spectrum of 3,4,5-tri-O-galloylquinic acid from *Cleistocalyx operculatus*; Fig. S27: mass spectrum of tri-O-methoxyellagic acid from *Cleistocalyx operculatus*; Fig. S28: mass spectrum of butyl isobutyl phthalate from *Cleistocalyx operculatus*; Fig. S29: mass spectrum of eudesmin from *Ficus lacor*; Fig. S30 mass spectrum of lasiodiplodin from *Ficus lacor*; Fig. S31 mass spectrum of 1-monopalmitin from *Ficus lacor*; Fig. S32 mass spectrum of α -tocospiro B from *Ficus lacor*. See DOI: <https://doi.org/10.1039/d3ra04037b>

‡ These authors contributed equally to this work.

Introduction

Despite considerable advancements in synthetic chemistry, the search for new and innovative pharmaceuticals is perpetual, and nature continues to be a crucial source of potential new medicines. Recent instances of natural product drug development have highlighted the use of genomic and metabolomics methodologies to supplement conventional methods of natural product research. Medicinal plants contain a range of bioactive metabolites with therapeutic properties. Therefore, robust techniques and technological advancements have been used to explore phytochemistry.

The field of metabolomics has been rapidly growing over the past two decades owing to its ability to provide information on dozens to hundreds of metabolites in a single experiment. It includes the identification and quantification of endogenous and exogenous metabolites present in biological samples using highly sensitive analytical tools and bioinformatics.¹ Significant advancements in metabolomics have enhanced its applicability to biomarker investigations, toxicological analysis, drug discovery, and natural product chemistry.² The metabolic data acquired in plant metabolomics using single analytical tools are often insufficient and incomplete, which has led the scientific community to utilize MS-based techniques such as GC-MS and LC-MS.³ Complex MS/MS data acquired in metabolomics experiments can be visualized and analyzed using a computer-based approach, such as molecular networking. The Global Natural Product Social Molecular Networking (GNPS) is an online bioinformatics platform currently being utilized in research to perform molecular networking. It tends to detect possible similarities among all MS/MS spectra in the dataset and extend annotation to unknown but closely related metabolites.^{4,5}

Diabetes is an unassailable burden in low and high-income countries.⁶ Managing diabetes remains a significant challenge for the scientific community since individuals with this condition are vulnerable to various health complications resulting in the annual mortality of 1.5 million people.⁷ Consequently, promoting research on medicinal plants for the identification of new alternative therapeutics for diabetes is currently relevant due to favorable outcomes associated with plant-based therapeutics in terms of long-term safety, mode of action, and metabolic activity.

Based on their ethnobotanical pharmaceutical properties, four medicinal plants, namely *C. operculatus*, *T. chebula*, *F. lacor*, and *F. semicordata*, were selected in this study. Although these plants have a wide range of pharmacological applications, this study intended to investigate the applicability of these species and their related metabolites as anti-diabetic drugs. The efficacy and toxicity of herbal formulations must be thoroughly explored even though they are prepared and consumed locally. Finding viable drug candidates from crude extracts is more challenging due to the complexity of biological samples.⁸

Various classes of drugs like insulin, α -glucosidase inhibitors (acarbose, miglitol, voglibose), biguanides (metformin, phenformin), thiazolidinediones, sulfonylureas, dipeptidyl-

peptidase-4 inhibitors, meglitinides, dopamine agonists have currently been used to treat diabetes. However, these are reported to have several side effects such as allergic reactions, increased risk of bladder cancer, hypoglycemia, bloating, flatulence, etc.⁹ As a result, the issue demands the development of new potent drugs with no or minimal human side effects.

The purpose of this research was to explore secondary metabolites using a mass spectrometry-based metabolomics approach with the GNPS platform and to evaluate the possible enzyme inhibitory activity of *C. operculatus*, *T. chebula*, *F. semicordata*, and *F. lacor* in targeting diabetes and its complications.

Materials and methods

Chemicals

Methanol, ethanol, ethyl acetate, dichloromethane, and hexane were purchased from Thermo Fisher Scientific (India). Gallic acid and quercetin were procured from HiMedia, an Indian supplier. α -Glucosidase (*Saccharomyces cerevisiae*), 4-nitrophenyl- α -D-glucopyranoside (pNPG), pancreatic porcine α -amylase (PPA), 2-chloro-4-nitrophenyl- α -D-maltotriose (CNPG3), and acarbose were ordered from Sigma-Aldrich (Germany).

Medicinal plants

Medicinal plants were collected from four distinct locations in Nepal, and the herbarium of each plant was submitted to the Central Herbarium of the Central Botany Department at Tribhuvan University, where voucher specimens were subsequently registered (Table 1). The entire collection of plant samples was shade-dried and pulverized into powder.

Methanolic extracts and fractionation

Methanolic extracts of four plants (*Cleistocalyx operculatus*, *Terminalia chebula*, *Ficus lacor*, and *Ficus semicordata*) were prepared using the cold percolation method, which involves the absorption of the powder for a day in methanol before percolation. The same procedure was repeated continuously for 3 days, and the collected solvent was dried at temperatures below 50 °C in the rotary evaporator. After 50 g of methanolic extracts were dissolved in distilled water for the fractionation process, the extracts were separated using the following solvents: hexane, DCM, and ethyl acetate in order of polarity to obtain four fractions from each plant. This procedure was performed three times for each solvent.²⁶

Determination of polyphenolic and flavonoid contents

Total polyphenol and flavonoid contents (TPC and TFC) were quantified using standard Folin-Ciocalteu and aluminum trichloride methods.²⁷⁻²⁹ Both reactions were performed in 200 μ L final volume, and absorbance was measured using a microplate reader (SynergyLX, BioTek, Instruments, Inc., USA). Calibration curves were generated by using various concentrations of gallic acid and quercetin. The extract concentrations were calculated and expressed as gallic acid and quercetin equivalents (mg GAE per g and mg QE per g, respectively).

Table 1 Medicinal plants used in the study and their ethnobotanical uses and reported chemical constituents

Voucher specimen	Medicinal plant	Family	Ethnobotanical uses	Chemical constituents
TUCH 210018	<i>Terminalia chebula</i>	Combretaceae	It is used for antidiabetic, anticancer, cardioprotective, neuroprotective, anti-inflammatory, and antiarthritic activities. ¹⁰ It is also used to treat dyspepsia, piles, hepatosplenomegaly, irritable bowel syndrome, and heart failure. ¹¹	Gallic acid, digallic acid, ellagic acid, caffeic acid, pyragallol, rutin, quercetin, isoquercetin, chebulagic acid, chebulinic acid, chebulic acid, eugenol, terflavin A, terchebulin ¹⁰
TUCH 210017	<i>Cleistocalyx operculatus</i>	Myrtaceae	It is used for its cytotoxic, ¹² antitumor, ¹³ antihyperglycemic, ¹⁴ cardiotonic, ¹⁵ and anti-inflammatory ¹⁶ activities	2',4'-Dihydroxy-6'-methoxy-3',5'-dimethylchalcone and ursolic acid, 5,7,8,4'-tetrahydroxy-3'-5'-dimethoxyflavone-3-O-β-D-galactopyranoside, gossypetin-8, 3'-dimethyl ether-3-O-β-D-galactoside, myricetin-3'-methyl ether-3-O-β-D-galactopyranoside, myricetin-3'-methyl ether, quercetin, kaempferol ¹⁶ β-Sitosterol, lupeol, α-amarin, β-amarin, stigmasterol, campesterol, scutellarein glucoside, scutellarein, infection, sorbifolin, bergaptol, and bergapten ^{22,23}
TUCH 21022	<i>Ficus lacor</i>	Moraceae	The powdered ripe fruit is utilized as an anti-diabetic. ¹⁷ It is also used for wound healing ¹⁸ ulcers, ¹⁹ hay fever, dysentery, and stomach disorder. ¹⁸⁻²¹	α-amarin, β-amarin, stigmasterol, campesterol, scutellarein glucoside, scutellarein, infection, sorbifolin, bergaptol, and bergapten ^{22,23}
TUCH 21020	<i>Ficus semicordata</i>	Moraceae	For headaches and diarrhea, raw fruit juice and fruits are employed. Fruits are also used for visceral ulcers, jaundice, colic pain, and hepatitis. Externally, the juice of the leaves is useful to treat scabies. It has been shown to have antioxidant, antibacterial, and anticancer properties. ²⁴	Catechin, rutin, quercetin, dodecane, indole, linalool, gallic acid and germacrene ²⁵

Antioxidant assay

The antioxidant assay was performed as described previously in literature.³⁰ The reaction was performed using an equal volume of plant extracts with 0.1 mM DPPH to maintain a final volume of 200 μL. After 15 min of incubation at room temperature in the dark, the absorbance was recorded at 517 nm. The percentage scavenging activity was determined using the following formula:

$$\% \text{ scavenging} = \frac{A_o - A_t}{A_o} \times 100$$

where A_o = optical density of the negative control (50% DMSO), and A_t = optical density of the test or reference sample.

Assay of α-glucosidase inhibition

The α-glucosidase inhibition was performed by dissolving the enzyme and substrate in a 50 mM phosphate buffer (pH 6.8). The process was initiated by mixing 20 μL of plant extracts of

varying concentrations with 20 μL enzyme (0.2 units) along with 120 μL phosphate buffer followed by 10 min incubation at 37 °C. After incubation, pNPG substrate was added (0.7 mM) and incubated for 15 min at the same temperature.³¹ The absorbance of the released *p*-nitrophenol was measured at 405 nm using Synergy LX (BioTek Instruments, Inc., USA). Each test was performed in triplicate. The following formula was used to calculate the % α-glucosidase inhibitory activity

$$\% \text{ inhibition} = \frac{A_o - A_t}{A_o} \times 100$$

where A_o is the optical density of the negative control with 30% DMSO and A_t is the optical density of the test/positive control.

Assay of α-amylase inhibition

The reaction was performed by dissolving the enzyme and substrate in 50 mM phosphate-buffered saline (pH; 0.9% NaCl, pH 7.0). The assay was inducted by mixing 20 μL plant extracts at different concentrations and adding 80 μL PPA (1.5 units per



mL) followed by 10 min incubation at 37 °C. After incubation, CNPG3 (0.5 mM) was added and the cells were incubated for 15 min.²⁶ The absorbance was measured at 405 nm using a microplate reader (Synergy LX, BioTek Instruments, Inc., USA). The percentage of inhibition was determined using the following formula:

$$\% \text{ inhibition} = \frac{A_o - A_t}{A_o} \times 100$$

where A_o is the optical density of the negative control with 30% DMSO and A_t is the optical density of the test/positive control.

Liquid chromatography-mass spectrometry analysis

LC-HRMS of the ethyl acetate and water fractions of *T. chebula* fruits, *C. operculatus*, and *F. lacor*, *F. semicordata* were analyzed at the Sophisticated Analytical Instrument Facility (SAIF), CSIR-Central Drug Research Institute, Lucknow (Lot No. 1607642300 and 1606747677). Compounds were identified using an Agilent 1200 series HPLC system (Agilent Technologies, Santa Clara, CA, USA) equipped with an Agilent 6520 I Accurate-Mass Q-TOF LC/MS system (Agilent Technologies, Santa Clara, CA, USA). An Agilent 6520 Accurate-Mass Q-TOF Mass Spectrometer equipped with a G1311A quaternary pump, G1329A autosampler, and G1315D diode array detector (DAD) was used for further analysis. The elution solvent comprised a mixture of acetonitrile, 5 mM acetate buffer, and water flowing at 1.5 mL min⁻¹, following a gradient from 5% to 30% acetonitrile over 10 minutes, reaching 80% acetonitrile over 32 minutes, and returning to the initial conditions, with the column temperature consistently held at 30 °C throughout the process. For positive mode analysis of electrospray ionization, the source gas temperature was maintained at 30 °C, with a gas flow rate of 11.01 L min⁻¹, and a nebulizer pressure of 40 psi. VCap voltage was set to 3500 V, fragmentor at 175 V, skimmer at 65.0 V, and Octopole RF peak at 750 V. The MS data acquisition was done in the range of 100–2000 Da at a scan rate of 1.03 for *T. chebula* (fruit), *C. operculatus*, *F. lacor*, and *F. semichordata*.³² Moreover, the molecular annotation of the methanolic extract (AT1) and ethyl acetate fraction (AT2) of *T. chebula* bark was completed in SIRIUS 5.6.3, using the CSI: FingerID interface, where the efficiency of the predicted molecular formula, functional group, and fragmentation pattern was evaluated in terms of the Sirius score.³³

HRMS and GNPS-based metabolomic analysis

Samples of AT1 and AT2 were prepared for MS/MS analysis by dissolving them in HPLC-grade (1 mg mL⁻¹), and then 150 µL

volume of each sample was transferred to an HPLC autosampler vial. Metabolomic profiling of the two samples was performed using an Agilent G6545B quadrupole time-of-flight (Q-TOF) mass spectrometer (Agilent Technologies, Santa Clara, CA, USA) equipped with a heated electrospray ion source (HESI). For chromatographic separation, an Acquity® UPLC BEH reverse-phase column C18 (150 mm × 2.1 mm, 1.7 µM) was used. The mobile phase was acidified with 0.1% formic acid and comprised of H₂O (A) and acetonitrile (B). The composition of organic solvent was 5% at 0.00–2.00 min, 20% at 5.00 min, 100% at 20.00 min, and back again to 5% at 23.00–25.00 min. The injection volume of two samples was kept at 3 µL and a flow rate of 0.5 mL min⁻¹ was maintained. MS/MS analysis was performed using electrospray ionization (ESI) in positive ion mode. Spectral hits were performed using a modified version of the method reported by Bashir *et al.* (2021)³⁴ with an *m/z* range of 50–1200, collision energies of 15 eV and 40 eV, and a full width at half maximum (FWHM) of 3000. The RAW files were converted to open-source '.mzXML' file format using the MSConvert tool of ProteoWizard MSConvert Version 3 software. The files were uploaded to the GNPS platform using WinSCP, the recommended FTP client. The acquired MS² data were visualized using a GNPS-based visualization following an established procedure (accessed on January 17, 2023). The molecular networks generated were further exported from GNPS to Cytoscape in '.graphml' format to enable customized visualization.

Data analysis

The collected data were processed using Gen5 Microplate Software, followed by MS Excel. GraphPad Prism Software version 8 was used for IC₅₀ calculation and data were represented as a mean ± standard error of the mean of triplicate. The Mnova software ver. 12.0 (Mestrelab Research, Santiago de Compostela, Spain) was used to process the raw data files obtained from the HPLC-QTOF-MS.

Results

Estimation of polyphenol contents and antioxidant potential

Phytochemical and antioxidant assays of the medicinal plants were performed in triplicate using methanolic extracts. The total phenol, flavonoid, and antioxidant activities of the plants are shown in Table 2. *T. chebula* (Bark and Fruit) showed the highest TPC (164.534 ± 3.70 mg GAE per g) and TFC values (158.67 ± 14.68 mg QE per g), respectively. Similarly, *T. chebula* (bark) showed the lowest IC₅₀ value of 5.144 ± 0.06 µg mL⁻¹ for

Table 2 Phytoconstituents and antioxidant activity of the methanol extracts of medicinal plants

Sample	TPC (mg GAE per g)	TFC (mg QE per g)	Antioxidant IC ₅₀ values (µg mL ⁻¹)
<i>Terminalia chebula</i> (fruit)	161.56 ± 0.39	158.67 ± 14.68	34.82 ± 1.12
<i>Terminalia chebula</i> (bark)	164.53 ± 3.70	115.34 ± 4.07	5.144 ± 0.06
<i>Cleistocalyx operculatus</i>	154.26 ± 1.01	69.17 ± 0.73	30.34 ± 1.12
<i>Ficus lacor</i>	108.67 ± 6.37	131.16 ± 7.75	130 ± 1.02
<i>Ficus semicordata</i>	107.76 ± 3.21	77.18 ± 0.73	22.76 ± 0.19
Quercetin (control)	—	—	6.3 ± 1.00



DPPH radical scavenging activity as compared with the control candidate, quercetin ($6.3 \pm 1.0 \mu\text{g mL}^{-1}$).

Inhibition of diabetic targets α -amylase and α -glucosidase

Preliminary screening was performed using $500 \mu\text{g mL}^{-1}$ of the extracts to determine the percentage inhibition of α -glucosidase and α -amylase, respectively. Only fractions with more than 50% inhibition were used to determine the IC_{50} values. As compared to the positive control, acarbose ($6.1 \pm 0.10 \mu\text{g mL}^{-1}$), and the lowest IC_{50} value towards α -amylase inhibition was found for *F. semicordata* in methanolic extract ($46.8 \pm 1.8 \mu\text{g mL}^{-1}$). Similarly, the lowest IC_{50} value for α -glucosidase inhibition was observed for *T. chebula* (fruits) in aqueous fraction ($1.07 \pm 0.01 \mu\text{g mL}^{-1}$) and for *C. operculatus* in methanolic extract ($1.26 \pm 0.03 \mu\text{g mL}^{-1}$). The results of α -glucosidase and α -amylase inhibition by different plant fractions are shown in Table 3.

Liquid chromatography and molecular annotation

Compound identification and characterization were based on the comparison of molecular masses, retention times, and MS spectra *via* authentic databases (PubChem, and Dictionary of Natural Products), as well as relevant literature. Because of the potent anti-diabetic activity exhibited by the ethyl acetate and water fractions of the four plants, the ethyl acetate and water fractions were selected for further molecular annotation analysis. Thirty-two compounds were identified from the ethyl acetate and water fraction of plant species, and the mass spectra of these compounds showed peaks at m/z 357.0455 (ESI Fig. S1†), 371.0619 (Fig. S2), 303.0140 (Fig. S3), 619.0970 (Fig. S4), 385.0788 (Fig. S5), 667.4054 (Fig. S6), 657.3627 (Fig. S7), 471.0197 (Fig. S8), 323.0762 (Fig. S9), 489.3570 (Fig. S10), 519.3313 (Fig. S11), 505.3529 (Fig. S12), 443.0978 (Fig. S13), 563.1549 (Fig. S14), 503.3363 (Fig. S15), 273.0759 (Fig. S16), 307.0815 (Fig. S17), 331.0463 (Fig. S18), 465.1023 (Fig. S19), 291.0859 (Fig. S20), 579.1504 (Fig. S21), 435.0578 (Fig. S22), 193.0705 (Fig. S23), 487.22 (Fig. S24), 467.0872 (Fig. S25), 649.1082 (Fig. S26), 393.0435 (Fig. S27), 279.1601 (Fig. S28), 387.1817 (Fig. S29), 293.1736 (Fig. S30), 331.2842 (Fig. S31), and 463.3794 (Fig. S32†) as protonated molecular ions $[\text{M} + \text{H}]^+$ as shown in Table 4. In ESI figures, ESI (+) indicates data collected in the positive mode of electrospray ionization mass spectrometry, centroid TIC and centroid MS indicate the data are presented in centroid mode only where in centroid mode each ion's mass and intensity are presented as discrete values, while profile mode indicates the entire distribution of intensity of ions recorded at each m/z value. Therefore, centroid MS data offers simpler spectra and helps in the interpretation of spectra. The annotated compounds and their fragment's peak were compared with previous findings as listed in Table 4. The structures of annotated molecules were drawn by using <http://www.chemspider.com/>. The structures of the annotated secondary metabolites are shown in Fig. 1.

GNPS-based metabolomic profiling

Based on the anti-diabetic activity of the methanol extract and ethyl acetate fraction of the bark of *T. chebula*, a comprehensive

Table 3 α -Amylase (A) and α -glucosidase (G) activities of various solvent fractions from medicinal plants^a

Medicinal plants	Minimum inhibitory concentration, IC_{50} ($\mu\text{g mL}^{-1}$)						Water fr.		
	Methanol ext.			Hexane fr.			Dichloromethane fr.	Ethyl acetate fr.	
	A	G	A	G	A	G	A	G	
<i>Terminalia chebula</i> (fruit)	153.2 \pm 1.91	95.37 \pm 0.30	<50%	236.06 \pm 4.78	<50%	95.5 \pm 1.5	12.72 \pm 0.40	309.1 \pm 0.9	1.07 \pm 0.01
<i>Terminalia chebula</i> (bark)	180.4 \pm 10.22	8.251 \pm 1.08	167.5 \pm 9.47	<50%	141.2 \pm 1.43	<50%	140.6 \pm 2.61	6.633 \pm 1.82	219.2 \pm 3.26
<i>Cleistocalyx operculatus</i>	62.49 \pm 1.89	1.26 \pm 0.03	414.9 \pm 7.8	80.31 \pm 1.22	429.4 \pm 5.2	95.19 \pm 0.75	100.8 \pm 1.9	8.67 \pm 0.55	31.96 \pm 0.74
<i>Ficus lacor</i>	485.6 \pm 1.8	33.78 \pm 3.6	<50%	76.57 \pm 0.5	355.2 \pm 11.2	<50%	240.1 \pm 3.4	94.2 \pm 8.1	4.91 \pm 0.02
<i>Ficus semicordata</i>	46.8 \pm 1.8	46.3 \pm 3.07	<50%	<50%	<50%	<50%	241.7 \pm 4.3	34.7 \pm 1.1	<50%
Acarbose	6.1 \pm 0.10 (α -amylase)						344.23 \pm 1.03 (α -glucosidase)		5.0 \pm 0.2

^a Extract: ext. Fraction: fr.

Table 4 Details of the annotated compounds from ethyl acetate and water fractions^a

Purposed compound	Calculated mass	Observed mass	Formula	RDBE	Absolute error (ppm)	(Rt) Min	Fragment peak	Solvent fraction and plant species	CSI: Finger ID (score%)	
Chebulic acid	356.0374	357.0455	C ₁₄ H ₁₂ O ₁₁	9	0.77	2.84	339, 321, 293, 203	EA <i>T. chebula</i> (fruits)	—	
2-O-Caffeoyl hydroxy citric acid	370.0530	371.0619	C ₁₅ H ₁₄ O ₁₁	9	2.88	2.93	191, 163, 145	EA <i>T. chebula</i> (fruits)	—	
Ellagic acid	302.0057	303.0140	C ₁₄ H ₆ O ₈	12	1.52	8.98	193	EA <i>T. chebula</i> (fruits)	—	
	302.0057	303.0143	C ₁₄ H ₆ O ₈	12	2.60	8.70	—	Water <i>C. operculatus</i>	—	
	302.0057	303.0134	C ₁₄ H ₆ O ₈	12	0.31	8.62	201.01, 229.01, 247.02, 201.01, 173.02, ^b	<i>T. chebula</i> (bark)	98.05	
Trigallyllevoglucosan IX	618.0910	619.0970	C ₂₀ H ₂₆ O ₂₂	8	2.83	10.62	153	EA <i>T. chebula</i> (fruits)	—	
2-O-Feruloylhydroxy citric acid	384.0690	385.0788	C ₁₆ H ₁₆ O ₁₁	9	4.39	4.58	385, 209, 195	EA <i>T. chebula</i> (fruits)	—	
Aijunglucoside I	666.3973	667.4054	C ₃₆ H ₅₈ O ₁₁	8	0.46	10.06	527.33, 483.34	<i>T. chebula</i> (bark)	99.41	
23-Galloylterminolic acid	656.3555	657.3627	C ₃₇ H ₅₂ O ₁₀	12	0.94	10.33	639.35, 469.32, 451.31, 405.31, ^b	<i>T. chebula</i> (bark)	55.46	
Flavogallonic acid	470.0115	471.0197	C ₂₁ H ₁₀ O ₁₃	17	0.73	5.68	187.14	453.09, 379.00, 363.01	72.99	
Leucodelphidiin	322.0683	323.0762	C ₁₅ H ₁₄ O ₈	9	0.24	6.85	305.06, 139.03, 217.04, 263.05, ^b	<i>T. chebula</i> (bark)	54.41	
Arjunolic acid	488.3496	489.3570	C ₃₀ H ₄₈ O ₅	7	0.92	11.50	163.03	163.03	92.54	
Bartogenic acid	518.3238	519.3313	C ₃₀ H ₄₈ O ₇	8	0.47	11.18	471.34, 453.33, 205.15, 135.04	<i>T. chebula</i> (bark)	90.33	
Aijungenin	504.3445	505.3529	C ₃₀ H ₄₈ O ₆	7	1.10	10.06	501.32, 215.14, 119.08, 187.14, ^b	<i>T. chebula</i> (bark)	95.0	
(—)Epicatechin-3-O-gallate	442.0894	443.0978	C ₂₂ H ₁₈ O ₁₀	14	1.41	8.95	437.30	469.33, 451.32, 297.17, 187.14, ^b	<i>T. chebula</i> (bark)	95.0
Gambiriin B1	562.1469	563.1549	C ₃₀ H ₂₆ O ₁₁	18	0.29	9.02	173.13, 405.31, 423.32	173.13, 405.31, 423.32	99.31	
Rotundioic acid	502.3288	503.3363	C ₃₀ H ₄₆ O ₆	8	0.77	13.40	11.04	139.03, 123.04, 153.01, 273.07, ^b	<i>T. chebula</i> (bark)	93.01
Butin	272.0679	273.0759	C ₁₂ H ₂₂ O ₄	10	0.77	8.95	437.12, 411.10, 139.03, 123.04, ^b	437.12, 411.10, 139.03, 123.04, ^b	93.01	
Gallocatechin or epigallocatechin	306.0734	307.0815	C ₁₅ H ₁₄ O ₇	9	0.89	7.22	273.07	449.30, 421.31, 485.32, 173.13, ^b	449.30, 421.31, 485.32, 173.13, ^b	82.97
4,4'-Di-O-methylellagic acid	330.0370	331.0463	C ₁₆ H ₁₀ O ₈	12	4.61	17.48	119.08, ^b	119.08, ^b	95.05	
Isoquercitrin	464.0949	465.1023	C ₂₁ H ₂₀ O ₁₂	12	3.31	8.95	95.05	123.04, 77.03, 51.02	83.65	
Catechin	290.0784	291.0859	C ₁₅ H ₁₄ O ₆	9	4.30	8.80	289.07, 139.03, 163.03, 181.04, ^b	289.07, 139.03, 163.03, 181.04, ^b	—	
	290.0784	291.0866	C ₁₅ H ₁₄ O ₆	9	0.29	6.66	65.03	225, 245, 270, 299, 300.9, 316	EA <i>C. operculatus</i>	—
Procyanidin B2	578.1418	579.1515	C ₃₀ H ₂₆ O ₁₂	18	3.84	10.79	7.77	303.04, 137.05	EA <i>C. operculatus</i>	—
	578.1418	579.1504	C ₃₀ H ₂₆ O ₁₂	12	1.24	7.44	7.77	123.04, 163.03, 247.06	EA <i>F. lacor</i>	—
Ellagic acid-O-pentoside	434.0479	435.0578	C ₁₉ H ₁₄ O ₁₂	14	4.67	7.27	303.01	123.04, 163.03, 247.06	Water <i>C. operculatus</i>	—
Quinic acid	192.0628	193.07	C ₇ H ₁₂ O ₆	2	3.95	3.02	—	175.06, 157.05, 113.02	EA <i>F. semicordata</i>	—
	192.0628	193.0705	C ₇ H ₁₂ O ₆	—	0.53	1.05	—	311.15, 295.11	Water <i>F. lacor</i>	—
	486.20	487.22	C ₂₃ H ₃₄ O ₁₁	7	4.90	12.60	100	311.15, 295.11	Water <i>F. semicordata</i>	—



Table 4 (Contd.)

Purposed compound	Calculated mass	Observed mass	Formula	RDBe	Absolute error (ppm)	(Rt) Min	Fragment peak	Solvent fraction and plant species	CSI: Finger ID (score%)
Hydroxy- β -gingerol- O - β -D-glucuronide	466.0700	467.0872	$C_{20}H_{18}O_{13}$	1.37	11.18	171.03, 53.01		EA <i>C. operculatus</i>	—
3-O-Galloylnorbergenin	648.0915	649.1082	$C_{28}H_{24}O_{18}$	1.76	10.17	495, 343		EA <i>C. operculatus</i>	—
3,4,5-Tri-O-galloylquinic acid								EA <i>C. operculatus</i>	—
Tri-O-methoxyellagic acid	392.0374	393.0435	$C_{17}H_{12}O_{11}$	4.29	3.21	—		Water <i>C. operculatus</i>	—
Butyl isobutyl phthalate	278.1512	279.1601	$C_{16}H_{22}O_4$	6	3.66	30.01		Water <i>T. chebula</i> (fruits)	—
	278.1512	279.1603	$C_{16}H_{22}O_4$	6	4.72	29.94		Water, EA <i>F. lacor</i>	—
	278.1512	279.1590	$C_{16}H_{22}O_4$	6	0.30	22.65		<i>T. chebula</i> (bark)	84.29
	278.1512	279.1592	$C_{16}H_{22}O_4$	6	0.40	18.64		Water <i>F. lacor</i>	—
Eudesmin	386.1723	387.1817	$C_{22}H_{26}O_6$	10	3.98	17.45		Water, EA <i>F. lacor</i>	—
Lasiodiplodin	292.1669	293.1736	$C_{17}H_{22}O_4$	6	3.00, 3.22	24.28, 24.01		Water, EA <i>F. lacor</i>	—
1-Monopalmitin	330.2764	331.2842	$C_{19}H_{38}O_4$	1	3.54, 0.72	27.76, 27.78	125.09	Water, EA <i>F. lacor</i>	—
α -Tocospiro B	462.3703	463.3794	$C_{29}H_{50}O_4$	5	2.75	36.62	—	Water <i>F. lacor</i>	—

^a EA: ethyl acetate, RDBe: ring double bond equivalents.

and detailed phytochemical profile was investigated using MS/MS and a GNPS-based metabolomics platform. A total of 205 and 160 individual ion species (MS/MS spectra) were present as nodes in the network map for the methanol extract and the EA fraction, respectively. Moreover, 22 compounds were tentatively identified from *T. chebula* bark, belonging to various groups of phytochemicals including phenolic compounds, diterpenoids-*O*-glycosides, triterpenoids, triterpenoid-*O*-glycosides, flavonoids, flavonoid-*O*-glycosides, fatty acids, and ceramides (Fig. 2, Table 5).

Molecular networking analysis involves representing each metabolite as a node labeled with its *m/z* value. The metabolites were then grouped into several clusters based on the similarity of their fragmentation patterns, indicating similarities in their core chemical structures.³⁵ From molecular networking and chart of categorized compounds (Fig. 2A), the major clusters included diterpenoid-*O*-glycosides, triterpenoids (Fig. 2B), flavonoid derivatives (Fig. 2C), and triterpenoid-*O*-glycosides (Fig. 2D).

The terpenoid clusters, characterized by precursor ions at *m/z* 471.3449, 485.3109, and 487.3403, were identified as arjunic acid, β -D-galactopyranoside, and madecassic acid, respectively. The flavonoids were annotated for *m/z* 291.0869 and 307.0736 clusters of epicatechin and epigallocatechin, respectively. Moreover, the neutral loss of 152 Da from both precursor ions at *m/z* 443.0947 and 459.0964 indicated the presence of gallate substitution, suggesting the presence of catechin gallate and epigallocatechin gallate. Finally, the networking of glycosylated triterpenoids, namely arjunglucoside I and arjunetin, was attributed to the presence of intense product ion signals at *m/z* 527.3334 and 511.3390, respectively, originating from the aglycone moiety. Additionally, the fragmentation patterns exhibited prominent neutral losses of 162 Da owing to the elimination of glucose. The various constituents observed in the two fractions from *T. chebula* bark are listed in Table 5 along with their *m/z* values in positive ion mode, retention times, MS/MS fragmentation patterns, and molecular formulas.

Discussion

This study aimed to explore the secondary metabolites, TPC, TFC, and antioxidant activities of extracts of *C. operculatus*, *T. chebula*, *F. semicordata*, and *F. lacor*. In addition, the enzyme inhibitory activities of the different fractions against α -glucosidase and α -amylase were determined and are presented in Tables 2 and 3. The results suggest that methanolic extracts, ethyl acetate, and water fractions showed potent activity against diabetic enzymes, which might be due to the synergistic effects of different chemical constituents. The ethyl acetate fractions exhibit a high flavonoid content, thereby attributing to their role in the inhibition of enzymes.^{36,37} Previous studies have also suggested that the methanolic, ethyl acetate, and water fractions significantly inhibit the activities of both α -glucosidase and α -amylase, respectively.³²

High blood glucose levels lead to the formation of free radicals that damage macromolecules within the cell and increase the risk of complications related to hyperglycemia. Thus, compounds with antioxidant properties can aid in the




Fig. 1 Structure of annotated compounds from *T. chebula*, *F. lacor*, *C. operculatus*, and *F. semicordata* based on LC-HRMS.

mitigation of diabetes-related problems.³⁸ Phenolic compounds are an important class of secondary metabolites with antioxidants as well as antidiabetic activity. Polyphenols with several hydroxyl groups can act as a source of hydrogen and electron donor to radicals to stabilize them and hence reduce oxidative stress which plays an important role in the mitigation of diabetes.³⁹ The introduction of hydroxyl and methoxyl groups to different positions of flavonoid structures is the most common

form in natural products. The patterns of hydroxyl and methoxyl groups on aryl rings A and B of flavonoids were reported to be closely associated with their biological activity towards enzymes. The hydroxylation of flavonoids at positions C3, C6, C3', and C4' all exhibited better inhibitory activity on α -glucosidase than the unhydroxylated forms at the corresponding positions.³⁹ The antidiabetic effect of flavonoids also can be increased by replacing the hydroxyl groups at the C3 position






Fig. 2 Molecular networking and identification of natural compounds from the extract of *T. chebula* bark. (A) Molecular networking and chart of categorized compounds in this study. (B) Zoomed-in molecular networking of triterpenoids and diterpenoids. (C) Zoomed-in molecular networking of flavonoids and flavonoid-O-glycosides. (D) Zoomed-in molecular networking of triterpenoid-O-glycosides.

with various functional group moieties, such as methoxy, sugar-like rhamnose, galloyl group, and chlorine atom. However, the glycosyl and geranyl moieties at C7 can suppress the antidiabetic effect. The presence of a C2=C3 double bond, along with the C4-oxo group and aryl ring A, is essential for maintaining the planar structure of flavonoids. Hydrogenation of this double bond weakens the enzyme's inhibitory activity likely because the planar molecular structure significantly influences the binding conformation within the enzyme's active site (Fig. 3).^{40,41}


Among four selected medicinal plants, *T. chebula* (bark) has shown excellent free radical scavenging activity with IC_{50} values of $5.144 \pm 0.06 \mu\text{g mL}^{-1}$ corresponding to control quercetin ($6.3 \pm 1.00 \mu\text{g mL}^{-1}$). This may be due to the presence of high TPC and TFC values in *T. chebula*. Such promising antioxidant potential could be ascribed to ellagic acid ($IC_{50} = 2.012 \pm 0.10 \mu\text{g mL}^{-1}$), catechin ($IC_{50} = 6.7 \mu\text{M}$), epicatechin ($IC_{50} = 6.8$

μM),⁴² arjunglucoside I, quercitrin, flavogallonic acid,⁴³ bartogenic acid and arjunic acid identified in the *T. chebula* bark extracts.

Since α -glucosidase and α -amylase enzymes are thought to be promising targets for the treatment of diabetes mellitus,⁴⁴ we assessed the antidiabetic potential of four medicinal plants. The promising inhibitory activity towards α -glucosidase was observed in the water fraction of *T. chebula* (fruits) with an IC_{50} value of $1.07 \pm 0.01 \mu\text{g mL}^{-1}$ followed by methanolic and water fraction of *C. operculatus* ($IC_{50} = 1.26 \pm 0.03 \mu\text{g mL}^{-1}$ and $IC_{50} = 4.91 \pm 0.02 \mu\text{g mL}^{-1}$ respectively). Moreover, the water fraction of *F. semicordata* ($IC_{50} = 5.0 \pm 0.2 \mu\text{g mL}^{-1}$) and ethyl acetate fraction of the *T. chebula* bark ($IC_{50} = 6.633 \pm 1.82 \mu\text{g mL}^{-1}$) displayed excellent activity as compared to acarbose (positive control, $IC_{50} = 344.23 \pm 1.03 \mu\text{g mL}^{-1}$). The high polyphenol and flavonoid contents in these plant extracts may be responsible for their promising α -glucosidase-inhibitory activity.

Table 5 Details of the identified constituents in *T. chebula* bark fractions using MS/MS and GNPS analysisMethanol (AT1) and EA (AT2) extract of the bark of *T. chebula*

Comp. name	RT	<i>m/z</i>	MS ² fragmentation pattern	Molecular formula	Exact mass	Fractions
Epigallocatechin	4.87	307.07	181.04, 139.03	C ₁₅ H ₁₄ O ₇	306.07	AT2
Epicatechin	7.42	291.09	165.05, 139.04, 123.04	C ₁₅ H ₁₄ O ₆	290.08	AT1, 2
Catechin	7.44	291.09	162.05, 139.04, 123.04	C ₁₅ H ₁₄ O ₆	290.08	AT1, 2
Procyanidin B2	7.50	579.15	409.09, 287.05, 247.06, 163.04, 127.04	C ₃₀ H ₂₆ O ₁₂	578.14	AT1, 2
Arjunglucoside I	10.47	689.39	527.33, 185.04, 89.06	C ₃₆ H ₅₈ O ₁₁	666.40	AT1, 2
Arjunetin	11.66	673.39	511.34, 185.04	C ₃₆ H ₅₈ O ₁₀	650.40	AT1, 2
Sesamin	15.60	337.11	319.09, 289.08, 267.06, 203.08, 185.06, 135.04	C ₂₀ H ₁₈ O ₆	354.11	AT1
Epigallocatechin gallate	7.93	459.09	139.04	C ₂₂ H ₁₈ O ₁₁	458.08	AT2
Quercitrin	8.34	449.09	303.04, 85.02	C ₂₁ H ₂₀ O ₁₁	448.10	AT1, 2
Ellagic acid	8.53	303.01	291.09, 273.07, 153.02	C ₁₄ H ₆ O ₈	302.01	AT1
(–)-Catechin 3-gallate	8.93	443.09	273.08, 153.02, 123.04	C ₂₂ H ₁₈ O ₁₀	442.09	AT1, 2
Epicatechin gallate	8.99	443.10	273.07, 139.04, 123.04	C ₂₂ H ₁₈ O ₁₀	442.09	AT1, 2
Arjunic acid	11.66	471.35	407.33, 201.16, 107.08	C ₃₀ H ₄₈ O ₅	488.35	AT2
Madecassic acid	12.19	487.342	451.32, 187.15, 145.10, 119.08, 95.08	C ₃₀ H ₄₈ O ₆	504.35	AT1, 2
β-D-Galactopyranoside	12.25	485.31	199.15, 171.12, 145.10, 105.07, 81.07	C ₂₆ H ₄₄ O ₈	484.30	AT2
Phytosphingosine	15.38	318.30	282.28, 60.04	C ₁₈ H ₃₉ NO ₃	317.29	AT1
Palmitic acid	17.38	402.36	283.26	C ₂₂ H ₄₂ O ₆	402.30	AT1
Dibutyl phthalate	18.64	279.16	149.02, 121.03	C ₁₆ H ₂₂ O ₄	278.15	AT1, 2
Linolenic acid	20.53	279.23	109.10, 95.05, 81.07, 67.05	C ₁₈ H ₃₀ O ₂	278.22	AT1, 2
Oleanolic acid	20.74	439.36	249.18, 217.19, 191.18, 95.08	C ₃₀ H ₄₈ O ₃	456.36	AT1
Diethyl phthalate	22.52	391.28	167.03, 149.02, 71.08, 57.07	C ₂₄ H ₃₈ O ₄	390.28	AT1
Erucamide	23.80	675.67	338.34, 321.31	C ₂₂ H ₄₃ NO	337.33	AT1, 2

Fig. 3 Structure–activity relationship of α -glucosidase inhibitory activity of flavonoids.

Previously, α -glucosidase inhibitory activity was observed with an IC_{50} value of $38.2 \mu\text{g mL}^{-1}$ (methanolic extract), and $9.0 \mu\text{g mL}^{-1}$ (water extract) for *T. chebula*, $68.2 \pm 3.4 \mu\text{g mL}^{-1}$ for *C. operculatus*,¹⁴ $4.226 \pm 0.43 \mu\text{g mL}^{-1}$ (ethyl acetate) for *F. semicordata*.⁴⁵

Similarly, the significant α -amylase inhibitory activity was shown by the methanolic extract of *F. semicordata* ($IC_{50} = 46.8 \pm 1.8 \mu\text{g mL}^{-1}$) followed by water and methanolic extract of *C. operculatus* ($53.2 \pm 1.0 \mu\text{g mL}^{-1}$ and $62.49 \pm 1.89 \mu\text{g mL}^{-1}$ respectively) corresponding positive control acarbose ($6.1 \pm 0.10 \mu\text{g mL}^{-1}$). In previous work, α -amylase inhibitory activity was observed in ethyl acetate fraction of *F. semicordata* and *T. chebula* extract with IC_{50} values of 4.861 ± 0.41 and $15.1 \pm 1.4 \mu\text{g mL}^{-1}$ respectively.^{45,46} The variation between our results and the literature data might be attributed to various factors, including the degree of ripeness at the time of harvesting, environmental factors, processing, and storage.⁴⁷

The MS-based metabolomics approach was employed in our study to putatively identify all the secondary metabolites

present in the plant extracts responsible for the anti-diabetic activity. Most of the compounds annotated in our study were similar to those in the literature; the ethyl acetate extract of *T. chebula* showed a base peak at $357.0455 [\text{M} + \text{H}]^+$ and fragment peaks at 339, 321, 293, and 203 corresponding to chebulic acid.⁴⁸ It has been reported as a potent compound for preventing the vascular complications associated with diabetes.⁴⁹ The base peak at 371.0619 , with fragment peaks at 191, 163, and 145, was predicted to be 2-O-caffeooyl hydroxy citric acid, which correlates with the fragmentation pattern reported in the literature.⁵⁰ The molecular formula $C_{14}H_{6}O_8$, with a base peak of 303.0140, has been proposed to be ellagic acid based on a comparative analysis of the spectral findings of Wu *et al.*⁵¹ As per the existing literature, ellagic acid demonstrates anti-diabetic effects by targeting the β -cells in the pancreas.⁵² Similarly, $[\text{M} + \text{H}]^+$ at $m/z 619.0970$ along with a fragment peak at 153 was annotated as trigalloevaloglucosan IX based on the results of Abu-Reidah *et al.*⁵³ Likewise, molecular ion $[\text{M} + \text{H}]^+$ at $m/z 385.07$ was annotated as 2-O-feruloyl hydroxy citric acid with fragment peaks at 385, 209, and 195. The spectral data were consistent with the literature.⁵⁰

A molecular ion at $m/z 667.4054 [\text{M} + \text{H}]^+$ detected at retention time 10.33 min was identified as arjunglucoside I in *T. chebula* bark, isolated previously from *T. arjuna*.⁵⁴ Previous study indicates that arjunglucoside I exhibits potential anti-diabetic effects against the α -glucosidase enzyme ($IC_{50} = 1074 \pm 32 \mu\text{M}$).⁵⁵ Similarly, a protonated ion detected at $m/z 657.3627 [\text{M} + \text{H}]^+$ was tentatively identified as galloylterminolic acid which was reported already in *T. albida*.⁵⁶ Another molecular ion at $m/z [\text{M} + \text{H}]^+$ was identified as flavogallonic acid, and was consistent with the literature found in the negative ESI-mode of



T. chebula.⁵⁷ The results of HPLC-MS analysis of *T. chebula* (fruits) showed a protonated ion with m/z 323.0762 [$M + H$]⁺ and sodiated ion as adduct at m/z 345.0580 [$M + Na$]⁺. Based on the literature, this compound was putatively identified as leuodelphinidin.⁵⁸ Arjunolic acid, previously identified in the *T. arjuna*,⁵⁹ was also identified in (+)-ESI-MS/MS analysis at m/z 489.3570 [$M + H$]⁺. Against α -glucosidase isolated from *S. cerevisiae*, arjunolic acid demonstrated anti-diabetic activity ($IC_{50} = 18.63 \pm 0.32 \text{ g mL}^{-1}$).⁶⁰ In addition, bartogenic acid was annotated for a molecular ion at m/z 519.3313 [$M + H$]⁺. This compound has demonstrated anti-diabetic properties against intestinal-glucosidase ($IC_{50} = 168.09 \text{ } \mu\text{g mL}^{-1}$).⁶¹ Another molecular ion at m/z 505.3529 [$M + H$]⁺ was identified as arjungenin, previously reported by Honda and coworkers.⁵⁴ It has been reported to show moderate antibacterial activity and beta-glucuronidase inhibitor activity.^{62,63} From *T. chebula* bark extract, a molecular ion with m/z 443.0978 [$M + H$]⁺ was tentatively annotated as (–)-epicatechin-3-O-gallate which was already reported by Singh *et al.*⁶⁴ Based on a literature survey, gambiriin B1 was identified as a molecular ion with m/z 563.1549 [$M + H$]⁺ in the *T. chebula* bark.⁶⁵ Similarly, rotundioic acid was isolated and identified.^{66,67} These data are consistent with our results, molecular ion peaks at 503.3363 [$M + H$]⁺ for rotundioic acid and 273.0759 [$M + H$]⁺ for butin. From the data obtained from the extract of *C. operculatus* and *T. chebula* bark, a protonated ion was detected at m/z 307.0815 and putatively identified as epigallocatechin or gallicatechin, which is similar to the study carried out by Lee and colleagues.⁶⁸ Its efficacy for the treatment of prostate cancer is being currently studied and completed in phase 2 trial.⁶⁹ It has been reported that epigallocatechin modifies glucose and lipid metabolism in rat hepatoma cell line H4IIIE and notably enhances glucose tolerance in diabetic rodents.⁷⁰

The compound exhibiting a base peak of 465.1023 and fragment peaks at 303 and 137 was annotated as isoquercitrin sourced from *F. lacor*.⁷¹ Isoquercitrin is currently being studied in a phase 2 clinical trial for its efficacy on Coronavirus Disease 2019.⁷² Moreover, it has been reported to demonstrate a prophylactic impact on diabetes mellitus.⁷³ Similarly, from *F. semicordata* and *T. chebula* (bark) molecular ions at m/z 579.1504 and at m/z 291.0866 were annotated as procyanidin B2 and catechin based on the result analysis.^{74,75} *In vitro*-glucosidase activity showed that procyanidin B2 could be used to reduce blood glucose levels.⁷⁶ Whereas, in the α -glucosidase inhibitory assay, it has been found that catechin has possessed significant antidiabetic activity ($IC_{50} = 87.55 \pm 2.23 \text{ } \mu\text{g mL}^{-1}$).⁷⁷ We annotated quinic acid from *F. semicordata* and *F. lacor* with a protonated ion at m/z 193.0705 which was similar to those reported in the literature.⁷⁸ A previous study showed that quinic acid potentially stimulates insulin secretion and enhances glucose tolerance.⁷⁹ Hydroxyl-6-gingerol- O - β -D-glucuronide was annotated for a protonated ion with m/z 487.22 from the extract of *F. semicordata* with comparison to the spectral results from Zeng *et al.*⁸⁰ The compound with molecular ion at m/z 435.0578 [$M + H$]⁺ was tentatively identified as ellagic acid- O -pentoside from *C. operculatus* based on the results obtained by Di Stefano and co-workers.⁸¹ Similarly, a compound with protonated ion at

m/z 467.0872 [$M + H$]⁺ was identified as 3-O-galloylnorbergenin, which was previously reported.⁵³ Moreover, molecular ions at m/z 649.1082 and 393.0435 were annotated as 3,4,5-tri- O -galloylquinic acid and tri- O -methoxyellagic acid, respectively from *C. operculatus*, as reported by Bindra *et al.* and Maldini *et al.*^{82,83} A molecular ion at m/z 279.1601 [$M + H$]⁺ and sodiated ion as adduct at m/z 301.1 [$M + Na$]⁺ was tentatively annotated as butyl isobutyl phthalate from *C. operculatus*, *T. chebula*, and *F. lacor*. This was previously reported by Bu *et al.* through LC/MS analysis, which was further supported by ESI-MS and NMR analysis.⁸⁴ It was observed that butyl isobutyl phthalate exhibits a hypoglycemic effect when tested in living organisms, and therefore, acts as a non-competitive inhibitor of α -glucosidase ($IC_{50} = 10.6 \pm 1.1 \text{ } \mu\text{g mL}^{-1}$).⁸⁴ Likewise, molecular ion at m/z 387.1817 with retention time at 17.45 min was tentatively annotated as eudesmin from the water fraction of *F. lacor* and was already reported.⁸⁵ Furthermore, based on literature data, a molecular ion at m/z of 293.1736 [$M + H$]⁺ was annotated as lasiodiplodin,⁸⁶ while the compound analyzed in the ESI-MS (+) mode molecular ion at m/z 331.2842 was tentatively identified as 1-monopalmitin.⁸⁷ Similarly, molecular ion at m/z 463.3794 was putatively identified as α -tocospiro B from the water fraction of *F. lacor*, as per the analysis performed by Chiang *et al.*⁸⁸

GNPS-based molecular networking can be performed to validate manually annotated compounds and aid in the exploration of secondary metabolite derivatives according to similarities in the core chemical structure.⁸⁹ Hence, this approach improves the visualization of the chemical constituents present in biological samples, forming clusters in molecular networking. In this study, the methanolic extract and EA fraction of the bark of *T. chebula* was subjected to comprehensive phytochemical profiling using MS/MS and a GNPS-based metabolomics platform, resulting in the identification of 22 compounds based on GNPS analysis and previous reports. Among these, 11 compounds, including linolenic acid, phytosphingosine, sesamin, diethyl phthalate, palmitic acid, oleanolic acid, quercitrin, epigallocatechin gallate, arjunic acid, β -D-galactopyranoside, madecassic acid, and arjunetin, were annotated based on GNPS that were not identified by manual annotation. These compounds were categorized into three distinct molecular network clusters: diterpenoid- O -glycosides, triterpenoids, flavonoid derivatives, and triterpenoid- O -glycosides clusters (Fig. 2B–D). All compounds were previously reported to be isolated from *T. chebula*, indicating that further investigation is necessary for the unknown nodes and edges of *T. chebula*.

Conclusion

Since ancient times, many traditional plants have been used by various ethnic groups to manage diabetes. Consequently, their scientific exploration is necessary. Four ethnically selected plants *C. operculatus*, *T. chebula*, *F. lacor*, and *F. semicordata*, exhibited significant enzyme inhibitory activity. In addition, LC/MS-based metabolomics and GNPS-based molecular networking were employed to explore and correlate secondary metabolites with the observed activities in these medicinal plants. Metabolic profiling



revealed the presence of different classes of bioactive metabolites, including phenolic compounds, diterpenoids-O-glycosides, triterpenoids, triterpenoid-O-glycosides, flavonoids, flavonoid-O-glycosides, fatty acids, and ceramides. Additional investigation is recommended to isolate and identify potential inhibitors from the active fractions of plant extracts to develop a therapeutic candidate for diabetes and to explore the underlying mechanisms responsible for the observed anti-diabetic action.

Author contributions

Niranjan Parajuli and Ki Hyun Kim designed and supervised the research; Arjun Prasad Timilsina, Bimal Kumar Raut, Chen Huo, and Rabin Budhathoki conducted computational metabolomics work; Bimal Kumar Raut, Arjun Prasad Timilsina, Karan Khadayat, and Niranjan Parajuli wrote the manuscript; Arjun Prasad Timilsina Prakriti Budhathoki, Mandira Ghimire and Karan Khadayat conducted *in vitro* assays; Niraj Aryal, Ki Hyun Kim, and Niranjan Parajuli edited the manuscript.

Conflicts of interest

There are no conflicts of interest among authors.

Acknowledgements

This project was supported by the University Grants Commission, Nepal (Award No. CRIG-78/79-S&T-01) to Niranjan Parajuli. We are thankful to the Sophisticated Analytical Instrument Facility (SAIF), CSIR-Lucknow, India, for the LC-HRMS data acquisition. We are also thankful to Kusum Basnet for her research assistance.

References

- 1 S. C. Booth, A. M. Weljie and R. J. Turner, Computational Tools For The Secondary Analysis of Metabolomics Experiments, *Comput. Struct. Biotechnol. J.*, 2013, **4**, e201301003.
- 2 Q. Yang, *et al.*, Metabolomics biotechnology, applications, and future trends: a systematic review, *RSC Adv.*, 2019, **9**, 37245–37257.
- 3 M. K. Patel, *et al.*, Plants Metabolome Study: Emerging Tools and Techniques, *Plants*, 2021, **10**, 2409.
- 4 L.-F. Nothias, *et al.*, Feature-based molecular networking in the GNPS analysis environment, *Nat. Methods*, 2020, **17**, 905–908.
- 5 F. Vincenti, *et al.*, Molecular Networking: A Useful Tool for the Identification of New Psychoactive Substances in Seizures by LC-HRMS, *Front. Chem.*, 2020, **8**, 572952.
- 6 N. Shrestha, S. R. Mishra, S. Ghimire, B. Gyawali and S. Mehata, Burden of Diabetes and Prediabetes in Nepal: A Systematic Review and Meta-Analysis, *Diabetes Ther.*, 2020, **11**, 1935–1946.
- 7 WHO, *Diabetes*, 2023, <https://www.who.int/news-room/fact-sheets/detail/diabetes>.
- 8 N. E. Thomford, *et al.*, Natural Products for Drug Discovery in the 21st Century: Innovations for Novel Drug Discovery, *Int. J. Mol. Sci.*, 2018, **19**, 1578.
- 9 Md. A. Hossain and R. Pervin, Chapter 34 – Current Antidiabetic Drugs: Review of Their Efficacy and Safety, in *Nutritional and Therapeutic Interventions for Diabetes and Metabolic Syndrome*, ed. D. Bagchi and S. Nair, Academic Press, 2nd edn, 2018, pp. 455–473, DOI: [10.1016/B978-0-12-812019-4.00034-9](https://doi.org/10.1016/B978-0-12-812019-4.00034-9).
- 10 M. Nigam, *et al.*, Fruits of *Terminalia chebula* Retz.: a review on traditional uses, bioactive chemical constituents and pharmacological activities, *Phytother. Res.*, 2020, **34**, 2518–2533.
- 11 N. N. Rege, U. M. Thatte and S. A. Dahanukar, Adaptogenic properties of six rasayana herbs used in Ayurvedic medicine, *Phytother. Res.*, 1999, **13**, 275–291.
- 12 C.-L. Ye, Y.-H. Lu, X.-D. Li, D.-Z. Wei and C. S. Whitehead, HPLC analysis of a bioactive chalcone and triterpene in the buds of *Cleistocalyx operculatus*, *S. Afr. J. Bot.*, 2005, **71**, 312–315.
- 13 C.-L. Ye, J.-W. Liu, D.-Z. Wei, Y.-H. Lu and F. Qian, In vivo antitumor activity by 2',4'-dihydroxy-6'-methoxy-3',5'-dimethylchalcone in a solid human carcinoma xenograft model, *Cancer Chemother. Pharmacol.*, 2005, **56**, 70–74.
- 14 T. T. Mai and N. V. Chuyen, Anti-Hyperglycemic Activity of an Aqueous Extract from Flower Buds of *Cleistocalyx operculatus* (Roxb.) Merr and Perry, *Biosci., Biotechnol., Biochem.*, 2007, **71**, 69–76.
- 15 A. Y. H. Woo, M. M. Y. Waye, H. S. Kwan, M. C. Y. Chan, C. F. Chau and C. H. K. Cheng, Inhibition of ATPases by *Cleistocalyx operculatus*. A possible mechanism for the cardiotonic actions of the herb, *Vasc. Pharmacol.*, 2002, **38**, 163–168.
- 16 B.-S. Min, T. D. Cuong, J.-S. Lee, M.-H. Woo and T. M. Hung, Flavonoids from *Cleistocalyx operculatus* buds and their cytotoxic activity, *Bull. Korean Chem. Soc.*, 2010, **31**, 2392–2394.
- 17 K. Y. Khan, *et al.*, Ethno-medicinal Species of genus *Ficus L.* Used to Treat Diabetes in Pakistan, *J. Appl. Pharm. Sci.*, 2011, **3**, 209–211.
- 18 P. K. Mukherjee, K. Mukherjee, M. Rajesh Kumar, M. Pal and B. P. Saha, Evaluation of wound healing activity of some herbal formulations, *Phytother. Res.*, 2003, **17**, 265–268.
- 19 S. K. Rai, S. Subedi and S. Mishra, Utilization pattern of medicinal plants in Thumpakhar, Village Development Committee, Sindhupalchok, Nepal, *Botanica Orientalis: Journal of Plant Science*, 2004, **4**, 75–78.
- 20 D. Bajracharya, S. J. B. Rana and A. K. Shrestha, A general survey and biochemical analysis of fodder plants found in Nagarjun hill forest of Kathmandu Valley [Nepal], *J. Nat. Hist. Mus.*, 1978, **2**, 105–116.
- 21 Y. Uprety, *et al.*, Diversity of use and local knowledge of wild edible plant resources in Nepal, *J. Ethnobiol. Ethnomedicine*, 2012, **8**, 16.
- 22 P. Ghimire, S. Pandey and A. Shrestha, A comprehensive review on phytochemical, pharmacognostical properties

and pharmacological activities of *Ficus lacor* L. (Moraceae), *Int. J. Tradit. Herb. Med.*, 2020, **8**, 96–102.

23 R. K. Sindhu and S. Arora, Phytochemical and Pharmacognostical Investigations on Aerial roots of *Ficus lacor* Buch. Ham, *Int. J. Phytomed.*, 2013, **5**, 267.

24 S. Gupta, Ethnomedicinal claims of *Ficus semicordata* Buch. Ham. ex Sm.: a review, *Int. J. Green Pharm.*, 2018, **12**, 206–213.

25 P. Deepa, K. Sowndhararajan, S. Kim and S. J. Park, A role of *Ficus* species in the management of diabetes mellitus: A review, *J. Ethnopharmacol.*, 2018, **215**, 210–232.

26 K. Khadayat, B. P. Marasini, H. Gautam, S. Ghaju and N. Parajuli, Evaluation of the alpha-amylase inhibitory activity of Nepalese medicinal plants used in the treatment of diabetes mellitus, *Clin. Phytosci.*, 2020, **6**, 34.

27 E. A. Ainsworth and K. M. Gillespie, Estimation of total phenolic content and other oxidation substrates in plant tissues using Folin–Ciocalteu reagent, *Nat. Protoc.*, 2007, **2**, 875–877.

28 C.-C. Chang, M.-H. Yang, H.-M. Wen and J.-C. Chern, Estimation of total flavonoid content in propolis by two complementary colometric methods, *J. Food Drug Anal.*, 2020, **10**, 178–182.

29 X. Lu, *et al.*, Determination of total phenolic content and antioxidant capacity of onion (*Allium cepa*) and shallot (*Allium oschaninii*) using infrared spectroscopy, *Food Chem.*, 2011, **129**, 637–644.

30 T. Sabudak, O. Demirkiran, M. Ozturk and G. Topcu, Phenolic compounds from *Trifolium echinatum* Bieb. and investigation of their tyrosinase inhibitory and antioxidant activities, *Phytochemistry*, 2013, **96**, 305–311.

31 H. Fouotsa, *et al.*, Xanthones inhibitors of α -glucosidase and glycation from *Garcinia nobilis*, *Phytochem. Lett.*, 2012, **5**, 236–239.

32 B. Aryal, *et al.*, LC-HRMS Profiling and Antidiabetic, Antioxidant, and Antibacterial Activities of *Acacia catechu* (L.f.) Willd, *BioMed Res. Int.*, 2021, **2021**, e7588711.

33 K. Dührkop, K. Scheubert and S. Böcker, Molecular Formula Identification with SIRIUS, *Metabolites*, 2013, **3**, 506–516.

34 K. Bashir, *et al.*, Assessing the ethnobotanical potential of *Carissa opaca* berries by merging outcomes from metabolomics profiling, enzyme assays, and in silico docking studies, *Food Chem.*, 2021, **363**, 130259.

35 R. A. Quinn, *et al.*, Molecular Networking As a Drug Discovery, Drug Metabolism, and Precision Medicine Strategy, *Trends Pharmacol. Sci.*, 2017, **38**, 143–154.

36 G. Pistia-Brueggeman and R. I. Hollingsworth, A preparation and screening strategy for glycosidase inhibitors, *Tetrahedron*, 2001, **57**, 8773–8778.

37 F. Sobreira, *et al.*, Gastroprotective activity of the hydroethanolic extract and ethyl acetate fraction from *Kalanchoe pinnata* (Lam.) Pers, *Braz. J. Pharm. Sci.*, 2017, **53**, e16027.

38 Y.-Z. Cai, M. Sun, J. Xing, Q. Luo and H. Corke, Structure–radical scavenging activity relationships of phenolic compounds from traditional Chinese medicinal plants, *Life Sci.*, 2006, **78**, 2872–2888.

39 K. E. Heim, A. R. Tagliaferro and D. J. Bobilya, Flavonoid antioxidants: chemistry, metabolism and structure-activity relationships, *J. Nutr. Biochem.*, 2002, **13**, 572–584.

40 J. Xiao, G. Kai, K. Yamamoto and X. Chen, Advance in Dietary Polyphenols as α -Glucosidases Inhibitors: A Review on Structure-Activity Relationship Aspect, *Crit. Rev. Food Sci. Nutr.*, 2013, **53**, 818–836.

41 N. F. Shamsudin, *et al.*, Flavonoids as Antidiabetic and Anti-Inflammatory Agents: A Review on Structural Activity Relationship-Based Studies and Meta-Analysis, *Int. J. Mol. Sci.*, 2022, **23**, 12605.

42 J. E. Hayes, P. Allen, N. Brunton, M. N. O’Grady and J. P. Kerry, Phenolic composition and in vitro antioxidant capacity of four commercial phytochemical products: Olive leaf extract (*Olea europaea* L.), lutein, sesamol and ellagic acid, *Food Chem.*, 2011, **126**, 948–955.

43 J. Chen, *et al.*, Structural elucidation and antioxidant activity evaluation of key phenolic compounds isolated from longan (*Dimocarpus longan* Lour.) seeds, *J. Funct. Foods*, 2015, **17**, 872–880.

44 A. Mocan, *et al.*, Biological effects and chemical characterization of *Iris schachtii* Markgr. extracts: A new source of bioactive constituents, *Food Chem. Toxicol.*, 2018, **112**, 448–457.

45 V. Kaur, K. Upadhyaya and M. Pande, Bioassay-Guided Evaluation of *Ficus Semicordata* For Antidiabetic Activity, *Int. J. Pharm. Pharm. Sci.*, 2017, 71–77, DOI: [10.22159/ijpps.2017v9i3.16441](https://doi.org/10.22159/ijpps.2017v9i3.16441).

46 D. Y. Lee, H. Yang, H. W. Kim and S. H. Sung, New polyhydroxytriterpenoid derivatives from fruits of *Terminalia chebula* Retz. and their α -glucosidase and α -amylase inhibitory activity, *Bioorg. Med. Chem. Lett.*, 2017, **27**, 34–39.

47 C. Manach, A. Scalbert, C. Morand, C. Rémésy and L. Jiménez, Polyphenols: food sources and bioavailability, *Am. J. Clin. Nutr.*, 2004, **79**, 727–747.

48 J. Yin, *et al.*, Characterization of chemical composition and prebiotic effect of a dietary medicinal plant *Penthorum chinense* Pursh, *Food Chem.*, 2020, **319**, 126568.

49 H.-S. Lee, Y.-C. Koo, H. J. Suh, K.-Y. Kim and K.-W. Lee, Preventive effects of chebulic acid isolated from *Terminalia chebula* on advanced glycation endproduct-induced endothelial cell dysfunction, *J. Ethnopharmacol.*, 2010, **131**, 567–574.

50 I. Parveen, A. Winters, M. D. Threadgill, B. Hauck and P. Morris, Extraction, structural characterisation and evaluation of hydroxycinnamate esters of orchard grass (*Dactylis glomerata*) as substrates for polyphenol oxidase, *Phytochemistry*, 2008, **69**, 2799–2806.

51 L. Wu, Y. Liu, Y. Qin, L. Wang and Z. Wu, HPLC-ESI-qTOF-MS/MS Characterization, Antioxidant Activities and Inhibitory Ability of Digestive Enzymes with Molecular Docking Analysis of Various Parts of Raspberry (*Rubus idaeus* L.), *Antioxidants*, 2019, **8**, 274.

52 N. Fatima, *et al.*, Ellagic acid in *Embllica officinalis* exerts anti-diabetic activity through the action on β -cells of pancreas, *Eur. J. Nutr.*, 2017, **56**, 591–601.



53 I. M. Abu-Reidah, M. S. Ali-Shtayeh, R. M. Jamous, D. Arráez-Román and A. Segura-Carretero, HPLC-DAD-ESI-MS/MS screening of bioactive components from *Rhus coriaria* L. (Sumac) fruits, *Food Chem.*, 2015, **166**, 179–191.

54 T. Honda, T. Murae, T. Tsuyuki, T. Takahashi and M. Sawai, Arjungenin, Arjunglucoside I, and Arjunglucoside II. A New Triterpene and New Triterpene Glucosides from *Terminalia arjuna*, *Bull. Chem. Soc. Jpn.*, 1976, **49**, 3213–3218.

55 H. Yuca, *et al.*, α -Glucosidase and α -amylase inhibitory potential of main compounds and drug candidates from *Elaeagnus rhamnoides* (L.) A. Nelson, *Chem. Pap.*, 2022, **76**, 913–922.

56 A. Camara, *et al.*, *Terminalia albida* treatment improves survival in experimental cerebral malaria through reactive oxygen species scavenging and anti-inflammatory properties, *Malar. J.*, 2019, **18**, 431.

57 Y. Wang, *et al.*, Hydrolyzable tannins from *Balanophora polyandra*, *Acta Pharm. Sin. B*, 2013, **3**, 46–50.

58 M. B. T. Tali, *et al.*, *In vitro* antiplasmodial activity-directed investigation and UPLC-MS fingerprint of promising extracts and fractions from *Terminalia ivorensis* and *Terminalia brownii* Fresen, *J. Ethnopharmacol.*, 2022, **296**, 115512.

59 T. M. Manu, T. Anand, M. D. Pandareesh, P. B. Kumar and F. Khanum, *Terminalia arjuna* extract and arjunic acid mitigate cobalt chloride-induced hypoxia stress-mediated apoptosis in H9c2 cells, *Naunyn-Schmiedeberg's Arch. Pharmacol.*, 2019, **392**, 1107–1119.

60 W. Hou, *et al.*, Triterpene acids isolated from *Lagerstroemia speciosa* leaves as α -glucosidase inhibitors, *Phytother. Res.*, 2009, **23**, 614–618.

61 P. M. Gowri, A. K. Tiwari, A. Z. Ali and J. M. Rao, Inhibition of α -glucosidase and amylase by bartogenic acid isolated from *Barringtonia racemosa* Roxb. seeds, *Phytother. Res.*, 2007, **21**, 796–799.

62 D. P. A. Gossan, *et al.*, Antibacterial and cytotoxic triterpenoids from the roots of *Combretum racemosum*, *Fitoterapia*, 2016, **110**, 89–95.

63 A. Rahman, *et al.*, Some Chemical Constituents Of *Terminalia Glaucescens* And Their Enzymes Inhibition Activity, *Z. Naturforsch., B: J. Chem. Sci.*, 2005, **60**, 347–350.

64 A. Singh, S. Kumar and B. Kumar, LC-MS Identification of Proanthocyanidins in Bark and Fruit of six *Terminalia* species, *Nat. Prod. Commun.*, 2018, **13**, 1934578X1801300.

65 S. Taniguchi, *et al.*, Revised Structures of Gambiriins A1, A2, B1, and B2, Chalcone-Flavan Dimers from Gambir (*Uncaria gambir* Extract), *Chem. Pharm. Bull.*, 2007, **55**, 268–272.

66 B. Yang, *et al.*, A facile and selective approach to the qualitative and quantitative analysis of triterpenoids and phenylpropanoids by UPLC/Q-TOF-MS/MS for the quality control of *Ilex rotunda*, *J. Pharm. Biomed. Anal.*, 2018, **157**, 44–58.

67 R. Chokchaisiri, *et al.*, Bioactive Flavonoids of the Flowers of *Butea monosperma*, *Chem. Pharm. Bull.*, 2009, **57**, 428–432.

68 J.-H. Lee, J. V. Johnson and S. T. Talcott, Identification of Ellagic Acid Conjugates and Other Polyphenolics in *Muscadine Grapes* by HPLC-ESI-MS, *J. Agric. Food Chem.*, 2005, **53**, 6003–6010.

69 University of Southern California, *A Phase II Trial of a Combination Herbal Therapy for Men With Biochemical Recurrence of Prostate Cancer After Initial Local Therapy*, 2017, <https://clinicaltrials.gov/study/NCT00669656>.

70 S. Wolfram, *et al.*, Epigallocatechin Gallate Supplementation Alleviates Diabetes in Rodents1, *J. Nutr.*, 2006, **136**, 2512–2518.

71 Y. Cai, L. Wu, X. Lin, X. Hu and L. Wang, Phenolic profiles and screening of potential α -glucosidase inhibitors from *Polygonum aviculare* L. leaves using ultra-filtration combined with HPLC-ESI-qTOF-MS/MS and molecular docking analysis, *Ind. Crops Prod.*, 2020, **154**, 112673.

72 AB Science, *Randomized, Phase 2 Clinical Trial to Evaluate the Safety and Efficacy of Masitinib Combined With Isoquercetin, and Best Supportive Care in Hospitalized Patients With Moderate and Severe COVID-19*, 2023, <https://clinicaltrials.gov/study/NCT04622865>.

73 S. Panda and A. Kar, Antidiabetic and antioxidative effects of *Annona squamosa* leaves are possibly mediated through quercetin-3-O-glucoside, *BioFactors*, 2007, **31**, 201–210.

74 A. Gahlaut, A. Sharma, A. Shirolkar and R. Dabur, Non-targeted identification of compounds from regenerated bark of *Terminalia tomentosa* by HPLC-(+) ESI-QTOFMS, *J. Pharm. Res.*, 2013, **6**, 415–418.

75 T. Li, *et al.*, Analysis of Chemical Constituents of *Guizhi Shaoyao Zhimu Granules* by UPLC-Q-Orbitrap HRMS, 2020, <https://www.researchsquare.com/article/rs-73167/v1>.

76 C. H. J. Yu, Z. Migicovsky, J. Song and H. P. V. Rupasinghe, (Poly)phenols of apples contribute to in vitro antidiabetic properties: Assessment of Canada's Apple Biodiversity Collection, *Plants People Planet*, 2023, **5**, 225–240.

77 H. N. Mrabti, *et al.*, Separation, Identification, and Antidiabetic Activity of Catechin Isolated from *Arbutus unedo* L. Plant Roots, *Plants*, 2018, **7**, 31.

78 K. Masike, *et al.*, Highlighting mass spectrometric fragmentation differences and similarities between hydroxycinnamoyl-quinic acids and hydroxycinnamoyl-isocitric acids, *Chem. Cent. J.*, 2017, **11**, 29.

79 E. Heikkilä, *et al.*, The plant product quinic acid activates Ca^{2+} -dependent mitochondrial function and promotes insulin secretion from pancreatic beta cells, *Br. J. Pharmacol.*, 2019, **176**, 3250–3263.

80 S.-L. Zeng, P. Li and E.-H. Liu, Metabolic profile of Guge Fengtong tablet in rat urine and bile after oral administration, using high-performance liquid chromatography coupled with electrospray ionization quadrupole time-of-flight mass spectrometry, *Chin. J. Nat. Med.*, 2015, **13**, 954–960.

81 V. Di Stefano, S. Scandurra, A. Pagliaro, V. Di Martino and M. G. Melilli, Effect of Sunlight Exposure on Anthocyanin and Non-Anthocyanin Phenolic Levels in Pomegranate Juices by High Resolution Mass Spectrometry Approach, *Food*s, 2020, **9**, 1161.



82 R. S. Bindra, N. K. Satti and O. P. Suri, Isolation and structures of ellagic acid derivatives from *Euphorbia acaulis*, *Phytochemistry*, 1988, **27**, 2313–2315.

83 M. Maldini, P. Montoro and C. Pizza, Phenolic compounds from *Byrsonima crassifolia* L. bark: Phytochemical investigation and quantitative analysis by LC-ESI MS/MS, *J. Pharm. Biomed. Anal.*, 2011, **56**, 1–6.

84 T. Bu, M. Liu, L. Zheng, Y. Guo and X. Lin, α -glucosidase inhibition and the in vivo hypoglycemic effect of butyl-isobutyl-phthalate derived from the *Laminaria japonica* rhizoid, *Phytother. Res.*, 2010, **24**, 1588–1591.

85 S.-M. Seo, *et al.*, Furofuran lignans from the bark of *Magnolia kobus*, *Chem. Nat. Compd.*, 2008, **44**, 419–423.

86 K. Z. W. Lae, S. S. Su, N. N. Win, N. N. Than and H. Ngwe, Isolation of Lasiodiplodin and Evaluation of some Biological Activities of the Stem Barks of *Phyllanthus Albizzoides* (Kurz) Hook.f., *SciMed. J.*, 2019, **1**, 199–216.

87 M. E. Hergueta-Castillo, R. López-Ruiz, A. Garrido Frenich and R. Romero-González, Characterization of the composition of plant protection products in different formulation types employing suspect screening and unknown approaches, *J. Sci. Food Agric.*, 2022, **102**, 5995–6004.

88 Y.-M. Chiang and Y.-H. Kuo, Two Novel α -Tocopheroids from the Aerial Roots of *Ficus microcarpa*, *J. Cheminf.*, 2003, **34**, 5125–5128.

89 R. Schmid, *et al.*, Ion identity molecular networking for mass spectrometry-based metabolomics in the GNPS environment, *Nat. Commun.*, 2021, **12**, 3832.