RSC Advances



View Article Online

View Journal | View Issue

PAPER

Check for updates

Cite this: RSC Adv., 2023, 13, 35569

Received 21st September 2023 Accepted 28th November 2023

DOI: 10.1039/d3ra06440a

rsc.li/rsc-advances

1. Introduction

Biomass is a valuable carbon-neutral resource with unique properties, such as tailored surface functional groups and a large surface area. It is readily available and can be used as an inexpensive starting material in various fields. One of the most important uses of biomass, is the preparation of biochar using the pyrolysis process. Biochar obtained from the pyrolysis of biomass has low porosity, high specific surface area, and limited surface functional groups.¹⁻³ To overcome these problems to maximize the use of biochar, there are several ways to activation and functionalization of biochar. In the chemical activation method, fresh biochar is mixed with activation agents such as ZnCl₂, KOH, H₂SO₄, K₂CO₃, H₃PO₄, etc., and heated at high temperatures in an inert gas flow.4-6 In a physical activation method, biochar is pyrolyzed and exposed to carbon dioxide or a controlled stream-flow, or a combination of both at above 700 °C.7 In this way, most of the reactive points of the carbon can be removed, and the pores can open and interconnect.8 Therefore, its surface area can significantly be modified.9,10 Several factors can be affected on the chemicalactivation of biochar such as the concentration and

Aluminium-based ionic liquid grafted on biochar as a heterogeneous catalyst for the selective synthesis of tetrazole and 2,3-dihydroquinazolin 4(1*H*)-one derivatives[†]

Masoomeh Norouzi, 🗅 * Parisa Moradi 🕩 and Maryam Khanmoradi

2,3-Dihydroquinazolin-4(1*H*)-one and tetrazole are a class of nitrogen-containing heterocyclic compounds that play an important role in drug design and are an important part of many biological and industrial compounds. In this study, aluminium-based ionic liquid grafted onto a biochar surface (BC/[TESPMI] AlCl₄) was synthesized and used as a catalyst for the synthesis of a series of biological derivatives including 2,3-dihydroquinazolin-4(1*H*)-ones in ethanol at reflux conditions and tetrazoles in PEG-400 at 110 °C. All products were obtained with good selectivity and yield of 90 to 97%. The catalyst was thoroughly characterized using various techniques such as thermogravimetric analysis (TGA), energy-dispersive X-ray spectroscopy (EDX), and powder X-ray diffraction (PXRD), scanning electron microscopy (SEM), inductively coupled plasm (ICP), and transmission electron microscopy (TEM), which confirmed the successful chemical grafting. This methodology has several advantages, including shorter reaction time, high yield of product, and recyclability of the catalyst. The catalyst remained active for five reaction cycles.

temperature of the activation agent, feedstock type, and other factors.¹¹ In addition, biochar is a low-cost, big surface area, high water-holding capacity, and pyrogenic carbon-rich material.¹²⁻¹⁹ Therefore, these carbon-based materials have a good potential to be used as heterogeneous catalysts even in many industrial applications, and they can be used as catalysts directly.^{20–23}

Besides, ionic liquids have emerged as more environmentally friendly alternative solvents than conventional solvents and have attracted major attention from researchers.^{24–26} Typically, ionic liquid formation occurred when an organic cation was combined with an inorganic or organic anion.²⁷ According to qualitative and semi-qualitative studies, ionic liquids can replace conventional catalysts with the advantage that they do not cause any toxicity in the reaction mixture.²⁸ Immobilization of ionic liquid on the surface of biochar has emerged as a superb platform for the synthesis of unique heterogeneous catalysts.²⁹

Tetrazole compounds have been investigated for over 100 years and applied in a variety field of science such as gasgenerating compositions, organic synthesis, biochemical and pharmaceutical applications, and some others.^{30–32} The low toxicity and high physiological activity of tetrazoles led to the fact that these materials continue to be highly regarded as widely used materials.^{33–35}

2,3-Dihydroquinazolin-4(1H)-ones are a significant class of heterocyclic compounds that contain nitrogen. They have

Department of Chemistry, Faculty of Science, Ilam University, PO Box 69315516, Ilam, Iran. E-mail: m.norozi@ilam.ac.ir

[†] Electronic supplementary information (ESI) available. See DOI: https://doi.org/10.1039/d3ra06440a

gained considerable interest due to their varied therapeutic and pharmacological properties, which include hepatoprotective, vasodilator, antidebrillatory, antipyretic antiatherosclerotic, and analgesic effects.³⁶

Several reports have been published on the use of nano catalysts in the synthesis of tetrazole derivatives. For example, a review summarized the most important nano-based catalyst approaches used in the synthesis of 5-substituted 1H-tetrazole described by Mittal and Awasthi.37 Mohammad Hosein Afsarian and co-workers reported copper bis(diacetylcurcumin) 1,2-diaminobenzene Schiff base complex, SiO₂-[Cu-BDACDABSBC] as a heterogeneous catalyst in the presence of ascorbic acid and a solution of water/i-PrOH (50:50, V/V) media at reflux condition is described. The yields of product have been achieved in the range of 75-95% for 3-8 hours.³⁸ Dehghani and co-workers reported salen complex of Cu(II) supported on superparamagnetic Fe₃O₄(a)SiO₂ nanoparticles for the synthesis of tetrazole in DMF at 120 °C in 83-92% of product.³⁹ Sharghi and co-workers reported 4'-phenyl-2,2':6',2' '-terpyridine-copper(11) complex immobilized onto activated multi-walled carbon nanotubes [AMWCNTs-O-Cu(II)-PhTPY] (in DMF at 70 °C, 75-95% yield of products).⁴⁰ And the other work has been published using ZnO, Zn hydroxyapatite, Zn/Al HT (in DMF, at 120-130 °C, yields: 62-91%), FeCl₃-SiO₂, Sb₂O₃, γ-Fe₂O₃, BaWO₄ Cu₂O, CdCl₂ and ZnS (solvent: DMF/ MeOH, 2.5 mol% catalyst, 12 h).41-45 Many of these works have disadvantages such as low product efficiency, high reaction temperature, use of toxic solvents to promote the reaction, use of high amounts of catalyst, use of catalysts that cannot be recycled, etc. Also, there are several reports on the use of nano catalysts for the synthesis of 2,3-dihydroquinazolin-4(1H)-one derivatives such as solvent-free, mechanochemically scalable synthesis of 2,3-dihydroquinazolin-4(1H)-one using Brønsted acid catalyst reported by Gauravi Yashwantrao and co-workers (homogenous catalyst).46 The other strategies for the preparation of these compounds have been reported by Sun et al. in 2018 (110 °C, DMSO, 12 h)⁴⁷ and Pathare et al. in 2019 (toluene, R.T, 2 h).⁴⁸ Aqueous Facile Synthesis of 2,3-Dihydroquinazolin-4(1H)-One Derivatives by Reverse Zinc Oxide was reported by Jie Mou and the other work in 2020 (homogenous catalyst, 90 °C, 5 h).49 However, most of these protocols have limitations, such as excess oxidant, complicated reactions, harsh reaction conditions (up to 100 °C), tedious work-ups, non-renewable and toxic solvents, non-reusable catalyst, low yields and long reaction times. All of these approaches reveal that the catalytic formation of these compounds is still challenging and that the area demands to be developed further. Consequently, based on our research on heterogeneous supported catalysts⁵⁰⁻⁵³ to introduce a catalyst that does not have any of the drawbacks stated for the mentioned catalytic systems, herein, we reported aluminium-based 1-(triethoxysilyl) propyl-3-methylimidazolium chloride ionic liquid grafted on ([TESPMI]Cl)-biochar (BC/[TESPMI]AlCl₄) as a high active, low-cost, reusable, environmentally friendly, and a heterogeneous nature catalyst. Furthermore, the catalytic performance of BC/[TESPMI]AlCl₄ was investigated in the synthesis of 5substituted 1*H*-tetrazole and 2,3-dihydroquinazolin-4(1H)-one derivatives.

2. Experimental

2.1. Materials

1-methylimidazole, (3-chloropropyl) triethoxysilane, AlCl₃, and all other reagents, chemicals, and solvents were purchased from Merck, Aldrich, Fluka and were used without additional purification.

2.2. Instrumental measurements

Powder X-ray diffraction (PXRD) of catalyst were recorded on a Holland Philips diffractometer (Cu Ka, radiation at 40 kV and 30 mA). SEM, EDS and EDS-mapping analysis were performed by using a FESEM-TESCAN MIRA3 scanning electron microscope. The thermogravimetric analysis (TGA) curves are recorded between 30 and 800 °C in air with heating rate of 10 °C min⁻¹. FT-IR spectra were recorded with KBr pellets using a Bruker FT-IR spectrometer (model VRTEX 70). And the transmission electron microscope with operating voltage at 100 kV. The elemental Al content catalyst was determined by PerkinElmer Optima 7300D inductively coupled plasma (ICP). The BET test was performed by using a BELSORP MINI II instrument.

2.3. Synthesis of catalyst

Biochar was obtained by placing dried chicken manure (500 g) in a porcelain crucible and pyrolyzed at the common temperature range of pyrolysis means 400 °C to 800 °C. We reached the required temperature after approximately 30 minutes of heating with the carrier N_2 gas that scanned at 0.3 L min⁻¹. In continuous, it introduced the porcelain crucibles into the heating zone with a 0.03 L min⁻¹ rate of nitrogen flow. This process continues for 1 and then 2 hours. After that, the heating zone of porcelain crucibles was cut down, and cooled with N_2 sweep with 0.3 L min⁻¹ for 30 minutes. Resulting, a black solid is obtained which is biochar.⁵⁴

In a round-bottomed flask equipped with a condenser, a mixture of 1-methylimidazole (1 mmol) and (3-chloropropyl) triethoxysilane (1 mmol) was reflexed at 100 °C for 12 h, according to recent research.²⁶ After that, the reaction mixture was washed with *n*-hexane several times to get a pure yellowish viscous ionic liquid ([TESPMI]Cl).

BC/[TESPMI]Cl was prepared by dissolving 1.5 mmol of [TESPMI]Cl and 1 g of biochar in toluene and stirred at 90 °C for 24 h. The resulting product was purified by washing with EtOH for several times and drying at 60 °C and labeled as BC/ [TESPMI]Cl.

The ionic liquid content (weight stability) was determined by gravimetric method after complete drying. The amount of [TESPMI] Cl per gram of support was measured to be 1.13 mmol g^{-1} .

Finally, the nanocatalyst was prepared by dissolving of 1 mmol of $AlCl_3$ and 0.5 g of BC/[TESPMI]Cl in EtOH (20 mL), and the reaction mixture was stirred at 60 °C for 20 h. The desirable product was obtained after filtering, washing with the



 $\label{eq:scheme-for-preparation-of-BC/[TESPMI]AlCl_4} catalyst.$

mixture of ethanol/water, and dried at room temperature. This product labeled as BC/[TESPMI]AlCl₄.

2.4. General procedures for preparation of 5-substituted-1*H*-tetrazoles

5-Substituted-1*H*-tetrazoles were synthesized from a mixture of 1 mmol of benzonitrile and 1.2 mmol of sodium azide in PEG-400 solvent in the presence of 50 mg BC/[TESPMI]AlCl₄ for enough time at 110 °C. The end of reactions has been determined by TLC method and the obtained data for several derivatives were summarized in Table 2. In the end of each reaction, hot ethyl acetate has been added to the reaction mixture, and BC/[TESPMI]AlCl₄ catalyst has been separated from the reaction mixture by a simple filtration. After that, the reaction mixture was transferred to a decanter funnel containing water, HCl and ethyl acetate. The combined organic layers were dried over anhydrous Na₂SO₄, then filtered, and the solvent was removed



Fig. 1 The XRD pattern of $BC/[TESPMI]AlCl_4$ catalyst.



View Article Online

RSC Advances



under reduced pressure and finally the crude product was purified using a silica gel plate.

2.5. General procedures for preparation of 2,3dihydroquinazolin-4(1*H*)-ones

To a mixture of 2-aminobenzamide (1 mmol) and aldehyde (1 mmol) in ethanol (5 mL), 30 mg of BC/[TESPMI]AlCl₄ was added and stirred under reflux condition for appropriate reaction time (Table 4). The end of the reaction was determined by TLC. After this, the reaction mixture was cooled down and dissolved in hot ethanol. Then it filtered off and dried at 50 °C. The product was further purified by re-crystallization form water/ethanol mixture (1 : 1).



Fig. 3 The SEM images of BC/[TESPMI]AlCl₄ catalyst.





Fig. 5 The WDX analysis for the BC/[TESPMI]AlCl₄ catalyst.

3. Results and discussion

The BC/[TESPMI]AlCl₄ nanocatalyst was afforded according to a three-step operationally simple process, as shown in Scheme 1. It involves the discrete synthesis of a biochar and ionic liquid ([TESPMI]Cl) followed by its immobilization on the prepared support (BC/[TESPMI]Cl) and further complexation with the metal salt to obtain the final immobilized aluminum (BC/ [TESPMI]AlCl₄).^{72,73} The synthesized BC/[TESPMI]AlCl₄ catalyst has been characterized by various techniques such as BET, SEM, FT-IR, EDAX, TGA, WDX, XRD, ICP and TEM.

3.1. Catalyst characterizations

The internal structures of BC/[TESPMI]AlCl₄ was examined by XRD analyses, as shown in Fig. 1. The XRD pattern of biochar



Fig. 6 Representative TEM micrographs of BC/[TESPMI]AlCl₄ catalyst.



Fig. 7 $\,N_2$ adsorption–desorption isotherms and BJH-Plot of BC/ [TESPMI]AlCl_4 catalyst.



Fig. 8 The FT-IR spectra for (a) biochar (BC), (b) [TESPMI]Cl, (c) BC/ [TESPMI]Cl, and (d) BC/[TESPMI]AlCl₄ catalyst.

shows broad diffraction peaks around $2\theta = 17-30^{\circ}$, which reflects amorphous carbon.^{54,55} Furthermore, peaks observed at $2\theta = 19.87^{\circ}$, 21.90°, 26.66° and 31.74° in the XRD pattern of BC/ [TESPMI]AlCl₄ confirm the synthesis of the catalyst.

The thermal stability of inorganic and organic components can be obtained by thermogravimetric analysis (TGA) at different temperatures. Therefore, this technique was used to identify the thermal stability of the BC/[TESPMI]AlCl₄ catalyst, which results are shown in Fig. 2. As shown in Fig. 2, three stages of weight loss can be seen in the TGA diagram of catalyst, which correspond to 18%, 20%, and 27% of the weight, respectively. The first mass loss of catalyst is corresponding to the release of water and other solvents from the nanocatalyst network.⁵⁴ The second weight loss is related to the elimination of organic groups that are immobilized on the surface of biochar.⁵⁴ The third mass loss are corresponding to the continued biochar pyrolysis.

The morphology of $BC/[TESPMI]AlCl_4$ catalyst was explored using scanning electron microscopy (SEM) technique and depicted in Fig. 3. As shown in Fig. 3, the dense and interconnected nanoparticles with a nanometer size have a quasispherical morphology uniformly. These results confirm that biochar has not undergone any changes during the modification.

Energy-dispersive X-ray spectroscopy (EDS) is widely used in the characterizations of the content of different elements in the nanostructure's frameworks. Based on this analysis, C, Cl, Si, O, N and Al species are present on the structure of the catalyst (Fig. 4).

To further characterize the catalyst, wavelength dispersive Xray spectroscopy (WDX) was employed to determine the elemental distributed of BC/[TESPMI]AlCl₄ catalyst, as illustrated in Fig. 5. This analysis showed that all elements were homogeneously distributed in the structure of BC/[TESPMI] AlCl₄.

The exact amount of aluminium in BC/[TESPMI]AlCl₄ was measured by inductively coupled plasma optical emission spectroscopy (ICP-OES) analysis. Based on the ICP results, 0.73 mmol of Al was loaded onto 1.0 g of the catalyst which shows a high loading capacity for this nanocatalyst.

Also, the microscopic analysis of $BC/[TESPMI]AlCl_4$ was investigated by TEM (Fig. 6). The TEM technique showed that most biochar particles were roughly spherical, and its size distribution diagram shows a uniform size distribution with an average size less than 100 nm.

 Table 1
 Definition of conditions for the cycloaddition reaction of sodium azide (1.2 mmol) and benzonitrile (1 mmol) in the presence of BC/

 [TESPMI]AlCl4 nanocatalyst

| $\begin{array}{c} & \text{CN} \\ + & \text{NaN}_3 \end{array} \xrightarrow{\text{Amount of nano cat (mg)}} \\ & \text{Solvent, Temperature °C} \end{array} \xrightarrow{\text{N-N}} \\ & \text{NaN}_3 \end{array}$ | | | | | |
|--|-----------------------------|------------------------------------|------------------|------------|------------------------|
| Entry | Amount of the catalyst (mg) | Solvent | Temperature (°C) | Time (min) | Yield ^a (%) |
| 1 | 20 | PEG-400 | 110 | 230 | 67 |
| 2 | 30 | PEG-400 | 110 | 230 | 80 |
| 3 | 40 | PEG-400 | 110 | 150 | 89 |
| 4 | 50 | PEG-400 | 110 | 100 | 96 |
| 5 | 50 | EtOH | Reflux | 100 | Trace |
| 6 | 50 | H ₂ O | Reflux | 100 | 40 |
| 7 | 50 | Glysrole : coline chloride (1 : 1) | 110 | 100 | 45 |
| 8 | 50 | PEG-400 | 80 | 100 | 52 |

Table 2 Preparation of 1H-tetrazole derivatives using BC/[TESPMI]AlCl₄

| | R + NaN ₃ | BC/[TESPMI]AlCl ₄ PEG-400, 110°C | | |
|-------------------------------------|-----------------------|--|------------|------------------------|
| Entry | Nitrile | Product | Time (min) | Yield ^a (%) |
| 1 | NO ₂ CN | | 480 | 90 |
| 2 | CN OH | N N N N N N N N N N N N N N N N N N N | 20 | 97 |
| 3 | CN CN | | 60 | 94 |
| 4 | CN CN | | 100 | 96 |
| 5 | CN | | 60 | 95 |
| 6 | | | 360 | 92 |
| 7 | CN CF ₃ | CF ₃ | 100 | 95 |
| ^{<i>a</i>} Isolated yield. | | | | |

The porosity properties of BC/[TESPMI]AlCl₄ catalyst were explored by the nitrogen ads/des isotherm analysis (Fig. 7). According to previous reports on these analyses of biochar materials, the surface area and pore volume of BC/[TESPMI] AlCl₄ catalyst are lower than surface area and pore volume of biochar. The values of these parameters for this catalyst are 6 m² g⁻¹ (surface area) and 0.01 cm³ g⁻¹ (total pore volume).

Meanwhile, the pore diameter value of BC/[TESPMI]AlCl₄ catalyst is 8.0 nm, which is higher than of the BC substrate.⁵⁴ Considering that the pore diameter of BC/[TESPMI]AlCl₄ catalyst is between 2 and 50 nm.

To identify the functional groups in biochar (BC), [TESPMI]Cl, ionic liquid functionalized biochar (BC/[TESPMI]Cl) and BC/ [TESPMI]AlCl₄ catalyst, the FTIR spectra were taken, as shown in

Table 3 Evaluation of the reaction parameter on the synthesis of 2,3-dihydroquinazolin-4(1H)-one derivatives over the catalysis of (BC/[TESPMI] AlCl₄)



| Entry | Catalyst | Catalyst (mg) | Solvent | Temperature (°C) | Time (min) | Yield ^{a,b} (%) |
|------------|---------------------------------------|---------------|--------------|-------------------------------|------------|---------------------------------------|
| 1 | _ | _ | EtOH | Reflux | 5 h | N.R |
| 2 | Biochar | 45 | EtOH | Reflux | 120 | Trace |
| 3 | BC/[TESPMI]AlCl ₄ | 45 | EtOH | Reflux | 25 | 93 |
| 4 | BC/[TESPMI]AlCl ₄ | 30 | EtOH | Reflux | 25 | 92 |
| 5 | BC/TESPMIAlCl4 | 15 | EtOH | Reflux | 60 | 68 |
| 6 | BC/[TESPMI]AlCl ₄ | 5 | EtOH | Reflux | 90 | 20 |
| 7 | BC/TESPMIAlCl4 | 30 | $CHCl_3$ | Reflux | 85 | 0 |
| 8 | BC/TESPMIAlCl4 | 30 | DMSO | Reflux | 60 | 30 |
| 9 | BC/[TESPMI]AlCl ₄ | 30 | MeOH | Reflux | 60 | 16 |
| 10 | BC/TESPMIAlCl4 | 30 | DMF | 120 | 60 | 40 |
| 11 | BC/[TESPMI]AlCl ₄ | 30 | Acetonitrile | Reflux | 60 | 35 |
| 12 | BC/[TESPMI]AlCl ₄ | 30 | DI water | Reflux | 60 | 50 |
| 13 | BC/[TESPMI]AlCl ₄ | 30 | EtOH | 50 | 180 | 15 |
| 14 | BC/[TESPMI]AlCl ₄ | 30 | EtOH | r.t | 180 | Trace |
| a Taslatad | wield ^b Conditions 4 shler | | -1) | and de (1 mensel) estatent (2 | | -+ (5T) |

Isolated yield. ^b Conditions: 4-chlorobenzaldehyde (1 mmol) and 2-aminobenzamide (1 mmol), catalyst (30 mg) in EtOH solvent (5 mL).

Table 4 Synthesis of 2,3-dihydroquinazolin-4(1H)-one derivatives^a



ž

(00)



| Entry | Aldehyde | Product | Time (min) | Yield ^a (%) |
|-------|------------------------------------|--|------------|------------------------|
| 4 | Br H OH | NH OH NH OH H Br | 55 | 92 |
| 5 | H ₃ CO OCH ₃ | NH NH OCH ₃ OCH ₃ | 65 | 93 |
| 6 | H NO ₂ | NH NH NO2 | 120 | 90 |
| 7 | ОН | NH NH | 35 | 98 |
| 8 | H ₃ CO H | NH NH NH OCH3 | 45 | 94 |
| 9 | H ₃ C | NH NH H CH ₃ | 35 | 96 |
| 10 | O ₂ N H | NNH NNH NO2 | 55 | 91 |



Fig. 8. In Fig. 8a, which corresponds to BC, several characteristic peaks can be observed at the 3430 cm^{-1} wave number, reveling oxygen-containing functional groups stretching vibrations such as phenol, alcohol, and carboxylic acid groups. The peaks around 3000 cm^{-1} are assigned to CH₂ stretching vibrations.⁵⁴ The IR spectrum of [TESPMI]Cl (Fig. 8b) shows absorption bands at 3431, 3083, 2974 and 2891 cm^{-1} , which are related to the asymmetrical stretching of amine salt, CH₂ units in the imidazolium ring and aliphatic groups, respectively. The peaks at $1631 \text{ and } 1570 \text{ cm}^{-1}$

are associated with the stretching vibrations of C=N and C=C groups in imidazolium rings, respectively. In addition, the bands at the interval 1085, 783 and 485 cm⁻¹ are attributed to the stretching vibrations of Si-O-Si. As can be observed in Fig. 8c, the peaks at 1602 cm⁻¹ and 3405 cm⁻¹ are related to the presence of – C=N- and O-H groups on the surface of BC/[TESPMI]Cl. In Fig. 8d, some of the peaks that existed in the previous stage (Fig. 8c) have shifted to other frequencies and appeared a new



Scheme 2 A plausible hypothetical mechanism for the formation of 5-aryl 1H-tetrazole catalyzed by BC/[TESPMI]AlCla.



Scheme 3 A plausible hypothetical mechanism for the formation of 2,3-dihydroquinazolin-4(1H)-one catalyzed by BC/[TESPMI]AlCl4.

peak at 609 cm⁻¹ and 904 cm⁻¹ which confirms the formation of BC/[TESPMI]AlCl₄ catalyst.

3.2. Catalytic study

The catalytic performance of $BC/[TESPMI]AlCl_4$ catalyst was investigated in the synthesis of 1*H*-tetrazole derivatives. First, optimization of the reaction condition is considered in the reaction between benzonitrile and sodium azide. In this stage, the effect of solvent, amount of catalyst, and temperature were examined to the synthesis of corresponding tetrazole.

First, the amount of the catalyst was optimized as an important factor in catalytic reactions. In the presence of 20 mg of the catalyst, the reaction afforded a 67% yield (Table 1, entry 1). Therefore, higher amounts of the catalyst were checked (Table 1, entries 2–4). Among different amounts of the catalyst (30, 40, and 50 mg), it was found that the best results were observed in the presence of 50 mg of BC/[TESPMI]AlCl₄ catalyst.

After optimizing the amount of catalyst, we attempted to optimize the best solvent to perform this reaction in the presence of 50 mg of $BC/[TESPMI]AlCl_4$ (Table 1, entries 4–7). Our findings revealed that the solubility of sodium azide plays a significant role in the yield of the desired product. In aprotic polar and nonpolar solvents, the reaction resulted in a very low yield, likely due to the poor solubility of sodium azide. However, in polar protic solvents, we observed a higher yield of the corresponding triazole (Table 1, entries 6 and 7). Among the solvents examined, PEG-400 was found to be the best reaction medium (Table 1, entry 4).

Further the reaction temperature was optimized. As shown in Table 1 (entries 4 and 8), at low temperatures, the reaction rate was found to be very slow. Finally, 110 °C provided the best results in the synthesis of tetrazole products. Therefore, the best conditions for this selected reaction model in the presence of Paper



Fig. 9 Recycling of nanocatalyst for the preparation of 2,3-dihy-droquinazolin-4(1*H*)-one derivatives under optimized conditions.



Fig. 10 XRD pattern of BC/[TESPMI]AlCl₄ after five times reuse.

 $BC/[TESPMI]AlCl_4$ catalyst were obtained using 50 mg of the catalyst at 110 °C in PEG-400 as solvent.

Further, several benzonitrile derivatives were reacted with sodium azide in the presence of BC/[TESPMI]AlCl₄ catalyst under optimized conditions. The benzonitrile derivatives with different electron-donating or electron-withdrawing functional groups were investigated and produced corresponding tetrazoles with excellent yields (Table 2).

Also, the catalytic efficiency of $BC/[TESPMI]AlCl_4$ catalyst has been studied for the synthesis of 2,3-dihydroquinazolin-4(1*H*)one derivatives. To get the optimized reaction conditions, several parameters have been explored in the reaction condensation of 4-chlorobenzaldehyde and 2-aminobenzamide.

To investigate the role of the solvent in promoting the reaction and find the most suitable amount of the catalyst, the reaction was carried out in the absent of $BC/[TESPMI]AlCl_4$ catalyst and in the presence of biochar as catalyst. The results show that there was no reaction in either of these two reactions even after long reaction time (Table 3, entries 1 and 2). Next, we investigated the role of $BC/[TESPMI]AlCl_4$ as catalyst by performing different amounts of this catalyst including 45, 30, 15, and 5 mg (Table 3, entries 3–6). The results show that 30 mg of $BC/[TESPMI]AlCl_4$ revealed 92% yield of desired product. Significant efficiency was not obtained in the presence of lower amounts of the catalyst.

In next attempt to find the best solvent, the same reaction in two polar protic solvents (water, ethanol, and methanol) and polar aprotic solvents (DMF, $CHCl_3$, DMSO and acetonitrile) were explored (Table 3, entries 6–12), which can effectively dissolve both polar and non-polar compounds and provide a suitable environment for the reaction to occur.

As shown in Table 3, when the reaction is carried out in $CHCl_3$, as a solvent, the desired product was not observed after 85 min under reflux conditions. However, when the solvents were changed to DMF, DMSO, MeOH and H_2O , were shown moderate to good yield.

Preliminary results illustrated the maximum yields in EtOH (Table 3, entry 4). We chose EtOH as solvent for the synthesis of 2,3-dihydroquinazolin-4(1*H*)-one because it is green, safe, environmentally friendly, and easier work-up.

The optimized reaction conditions were extended to a series of aldehydes with electron donor and electron acceptor functional groups. As shown in Table 4, the corresponding 2,3-dihydroquinazolin-4(1H)-one derivatives were synthesized in 90–97% yields under optimized conditions.

A purposed mechanism for the tetrazole formation catalyzed by BC/[TESPMI]AlCl₄ is illustrated in Scheme 2.⁵⁶ At first, the catalyst is coordinated with C \equiv N functional groups, intermediate (1) formed. N₃ ion attacked on intermediate (1) by [3 + 2] cycloaddition reaction and formed intermediate (2), which finally formed the corresponding tetrazole (3).

Also, Scheme 3 presented a proposed mechanism pathway for the formation of 2,3-dihydroquinazolin-4(1*H*)-ones in the presence of BC/[TESPMI]AlCl₄ catalyst. Based on this mechanism, initially, the imidazolium cations and Al-based nanocatalyst interacted with carbonyl group in aldehyde. Next, the activated carbonyl is attacked by 2-aminobenzamide's NH₂, which imine intermediate (2) is formed. Afterwards, imine intermediate (2) and catalyst produces the intermediate (3). Finally, the target 2,3-dihydroquinazolin-4(1*H*)-ones (4) was formed by the nucleophilic cycloaddition of amide nitrogen to the imine active group.



Fig. 11 FE-SEM image of BC/[TESPMI]AICl₄ after five times reuse.

Table 5 Comparison results for BC/[TESPMI]AlCl₄ in the synthesis of 5-phenyl-1H-tetrazole with previously reported procedures

| Entry | Catalyst | Time (min) | Yield (%) | Ref. |
|-------|---|------------|-----------|-----------|
| 1 | CoY zeolite | 14 h | 95 | 57 |
| 2 | $[Fe_3O_4(a)TAM$ -Schiff-base-Cu(II)] complex | 100 | 98 | 58 |
| 3 | ZrO-SB-APT@MCM-41 | 130 | 89 | 59 |
| 4 | PdArg@boehmite | 420 | 97 | 60 |
| 5 | Fe ₃ O ₄ /ZnS HNSs | 24 h | 86 | 61 |
| 6 | Co-(PYT) ₂ @BNPs | 120 | 98 | 62 |
| 7 | CuFe ₂ O ₄ | 720 | 90 | 63 |
| 8 | Cu-TBA@biochar | 420 | 98 | 54 |
| 9 | La-Schiff base@MCM-41 | 120 | 98 | 64 |
| 10 | BC/[TESPMI]AlCl4 | 100 | 96 | This work |

The recyclability test of BC/[TESPMI]AlCl₄ catalyst has been done to prove the heterogeneous behavior and no metal leaching of the BC/[TESPMI]AlCl₄ catalyst in the preparation of 2,3-dihydroquinazolin 4(1*H*)-one derivatives. In this stage, the reaction between 4-chlorobenzaldehyde with 2-aminobenzamide has been tested under the optimal conditions. At the end of the reaction, the catalyst was recovered from the reaction by filtration. The recovered catalyst washed with ethanol and drying it at 50 °C. The recovered catalyst was reused in the next similar reaction. We continued this experiment up to 5 times. Finally, it was observed that the catalytic activity of BC/[TESPMI]AlCl₄ was not reduced significantly (Fig. 9).

To assess the stability of BC/[TESPMI]AlCl₄, the leaching of active species in the reaction mixture was examined using ICP analysis after five cycles. The findings indicated that the leaching amounts of aluminum is 0.032 mmol g⁻¹, which indicates that a small accumulation may lead to a slight decline in recycled catalyst's catalytic performance. It is believed that strong interactions between the Al and the atoms of the BC/ [TESPMI] play a significant role in preventing metal leaching during the reaction. This illustrates that no substantial leaching occurred during the reaction.

SEM and XRD analyses for $BC/[TESPMI]AlCl_4$ were investigated after five runs. The result of XRD analysis of the reused catalyst showed that the structure of the catalyst was preserved during the reactions (Fig. 10). Fig. 11 illustrates SEM analysis of $BC/[TESPMI]AlCl_4$ after five runs. As can be seen, almost the BC/

Table 6 Comparison results for BC/[TESPMI]AlCl₄ in the synthesis of 2-phenyl-2,3-dihydroquinazolin-4(1H)-one with previously reported procedures

| Entry | Catalyst | Time (min) | Yield (%) | Ref. |
|-------|--|------------|-----------|------------|
| 4 | To Q Qabiff have of Qu(-) | 60 | 00 | 6 5 |
| 1 | Fe_3O_4 -Schiff base of $Cu(II)$ | 60 | 99 | 65 |
| 2 | CuCl/Fe ₃ O ₄ -TEDETA | 30 | 97 | 66 |
| 3 | SBA/AuNP | 90 | 91 | 67 |
| 4 | Amberlyst-15 | 60 | 85 | 68 |
| 5 | Cu(NO ₃) ₂ /F ₃ O ₄ -DETA | 50 | 97 | 69 |
| 6 | H ₃ BO ₃ -MCM-41 | 45 | 90 | 70 |
| 7 | Ni-CP | 100 | 96 | 71 |
| 8 | BC/[TESPMI] AlCl ₄ | 35 | 98 | This work |

[TESPMI]AlCl₄ particles with the same size and morphology as the fresh catalyst confirm the spherical shape.

To demonstrate the benefit of $BC/[TESPMI]AlCl_4$ catalyst compared to other catalysts that have been used in this field, the obtained results in this work was compared with previous catalysts. For this aim, the cycloaddition of benzonitrile and sodium azide (Table 5) and condensation of benzaldehyde with 2-aminobenzamide (Table 6) were considered. Based on the obtained data, $BC/[TESPMI]AlCl_4$ catalyst can be compared with some reported catalysts in terms of efficiency and reaction time. As shown, $BC/[TESPMI]AlCl_4$ catalyst provides better performance in reaction time than others. Also, $BC/[TESPMI]AlCl_4$ can easily be recycled and reused.

In some reagents reported, toxic or expensive solvents are used. But in this work, PEG-400 is used as a green and environmentally friendly solvent.

4. Conclusions

In summary, aluminium-based ionic liquid immobilized biochar surface was used for the sustainable and green synthesis of 5-substituted 1*H*-tetrazole and 2,3-dihydroquinazolin-4(1*H*)-one derivatives as a reusable, highly effective, and eco-friendly nanocatalyst. The nanocatalyst was prepared through a threestep procedure and characterized using various techniques including EDX, BET, FT-IR, SEM, PXRD, TEM, ICP and TGA. The nanocatalyst could be easily separated from the reaction mixtures and reused up to five times while maintaining its effectiveness. This approach has several benefits, including the use of a reusable and eco-friendly catalyst, a green solvent, mild reaction conditions, low catalyst loading, and the ability to achieve high yields without requiring a chromatographic column to isolate products.

Conflicts of interest

There is no conflict of interest for authors.

Acknowledgements

This work was supported by the research facilities of Ilam University, Ilam, Iran.

References

- 1 J. Huang, Y. Jian, H. Li and Z. Fang, *Catal. Today*, 2022, **404**, 35.
- 2 M. Zuo, Q. Niu, Y. Zhu, S. Li, W. Jia, Z. Zhou, X. Zeng and L. Lin, *Ind. Crops Prod.*, 2023, **192**, 115953.
- 3 S. Tsyganova, E. Mazurova, G. Bondarenko and N. Chesnokov, *Wood Sci. Technol.*, 2016, **50**, 963.
- 4 J. T. Yu, A. M. Dehkhoda and N. Ellis, *Energy Fuels*, 2011, 25, 337.
- 5 K. Qian, A. Kumar, H. Zhang, D. Bellmer and R. Huhnke, *Renewable Sustainable Energy Rev.*, 2015, **42**, 1055.
- 6 Y. S. Ok, S. X. Chang, B. Gao and H.-J. Chung, *Environ. Technol. Innovation*, 2015, 4, 206.
- 7 S. Das, S. Ghosh, T. Kuila, N. C. Murmu and A. Kundu, *Biomass*, 2022, 2, 155.
- 8 I. M. Atadashi, M. K. Aroua, A. R. Abdul Aziz and N. M. N. Sulaiman, *J. Ind. Eng. Chem.*, 2013, **19**, 14.
- 9 D. Hu, M. Zhang, H. Xu, Y. Wang and K. Yan, *Renewable Sustainable Energy Rev.*, 2021, 147, 111253.
- R. P. Premalatha1, J. Poorna Bindu, E. Nivetha,
 P. Malarvizhi, K. Manorama, E. Parameswari and
 V. Davamani, *Front. Energy Res.*, 2023, 11, 1092637.
- 11 X. Cao, S. Sun and R. Sun, RSC Adv., 2017, 7, 48793.
- 12 T. M. E. Shareef and B. Zhao, *J. Agric. Chem. Environ.*, 2017, 6, 38–61.
- 13 G. S. Rani, A. Jyotsna and B. L. A. Prabhavathi Devi, *Asian J. Green Chem.*, 2019, **3**, 125.
- 14 J. Lee, K. H. Kim and E. E. Kwon, *Renewable Sustainable Energy Rev.*, 2017, 77, 70.
- 15 S. Abrishamkesh, M. Gorji, H. Asadi, G. H. Bagheri-Marandi and A. A. Pourbabaee, *Plant, Soil Environ.*, 2015, **61**, 475.
- 16 K. Angalaeeswari and S. P. B. Kamaludeen, *Int. J. Chem. Stud.*, 2017, 5, 442.
- 17 A. E. A. Z. Amin, Arab. J. Geosci., 2018, 11, 379.
- 18 T. T. S. Matos, J. Schultz, M. Y. Khan, E. F. Zanoelo, A. S. Mangrich, B. R. Araújo, S. Navickiene and L. P. C. Romão, *J. Braz. Chem. Soc.*, 2017, 28, 1975.
- 19 J. P. Bindu, D. D. Reddy, C. C. S. Rao, K. Manorama, L. Prasad, R. Mathur and D. V. L. Satyavathi, *Tob. Res.*, 2019, 45, 70.
- 20 H. Veisi, A. Nikseresht, N. Ahmadi, K. Khosravi and F. Saeidifar, *Polyhedron*, 2019, **162**, 240.
- 21 E. Akhavan, S. Hemmati, M. Hekmati and H. Veisi, *New J. Chem.*, 2018, **42**, 2782.
- 22 J. P. Bindu, D. D. Reddy, C. C. S. Rao, K. Manorama, L. Prasad and R. Mathur, *Tob. Res.*, 2020, 46, 47.
- 23 J. Lee, K.-H. Kim and E. E. Kwon, *Renewable Sustainable Energy Rev.*, 2017, 77, 70.
- 24 S. K. Singh and A. W. Savoy, J. Mol. Liq., 2020, 297, 112038.
- 25 M. Saraiva, S. Costa, P. Pinto and A. Azevedo, *ChemSusChem*, 2017, **10**, 2321.
- 26 A. V. Nakhate and G. D. Yadav, *ChemistrySelect*, 2018, **16**, 4547.
- 27 V. P. Natalia and R. Seddon Kenneth, *Chem. Soc. Rev.*, 2008, 37, 123.

- 28 A. Ghorbani-Choghamarani and M. Norouzi, J. Magn. Magn. Mater., 2016, 401, 832.
- 29 A. M. Al-Sabagh, F. Z. Yehia, D. R. K. Harding, G. Eshaq and A. E. ElMetwally, *Green Chem.*, 2016, **18**, 3997.
- 30 E. Lukowska-Chojnacka, J. Mierzejewska, M. Milner-Krawczyk, M. Bondaryk and M. Staniszewska, *Bioorg. Med. Chem.*, 2016, 24, 6058.
- 31 A. Qian, Y. Zheng, R. Wang, J. Wei, Y. Cui, X. Cao and Y. Yang, *Bioorg. Med. Chem. Lett.*, 2018, **28**, 344.
- 32 C. G. Neochoritis, T. Zhao and A. Domling, *Chem. Rev.*, 2019, 119, 1970.
- 33 F. Gao, J. Xiao and G. Huang, Eur. J. Med. Chem., 2019, 184, 111744.
- 34 K. Kalantari, A. M. Afifi, H. Jahangirian and T. J. Webster, *Carbohydr. Polym.*, 2019, 207, 588.
- 35 H. N. Tran, S. J. You and H. P. Chao, *Waste Manage. Res.*, 2016, **34**, 129.
- 36 M. Norouzi and Sh. Beiranvand, J. Chem. Sci., 2023, 135, 86.
- 37 R. Mittal and S. K. Awasthi, Synthesis, 2019, 51, 3765.
- 38 M. H. Afsarian, M. Farjam, E. Zarenezhad, S. Behrouz and M. N. Soltani Rad, *Acta Chim. Slov.*, 2019, 66, 874.
- 39 F. Dehghani, A. R. Sardarian and M. Esmaeilpour, J. Organomet. Chem., 2013, 743, 87.
- 40 H. Sharghi, S. Ebrahimpourmoghaddam and M. M. Doroodmand, J. Organomet. Chem., 2013, 738, 41.
- 41 M. L. Kantam, K. B. Shiva Kumar and K. P. Raja, *J. Mol. Catal. A*, 2006, **247**, 186.
- 42 T. Jin, F. Kitahara, S. Kamijo and Y. Yamamoto, *Tetrahedron Lett.*, 2008, **49**, 2824.
- 43 L. Lang, B. Li, W. Liu, L. Jiang, Z. Xu and G. Yin, *Chem. Commun.*, 2010, **46**, 448.
- 44 M. Nasrollahzadeh, Y. Bayat, D. Habibi and S. Moshaee, *Tetrahedron Lett.*, 2009, **50**, 4435.
- 45 G. Venkateshwarlu, A. Premalatha, K. C. Rajanna and P. K. Saiprakash, *Synth. Commun.*, 2009, **39**, 4479.
- 46 G. Yashwantrao, V. P. Jejurkar, R. Kshatriya and S. Saha, *ACS Sustainable Chem. Eng.*, 2019, 7, 13551.
- 47 J. W. Sun, T. Tao, D. Xu, H. Cao, Q. G. Kong and X. Wang, *Tetrahedron Lett.*, 2018, **59**, 2099.
- 48 R. S. Pathare, A. K. Maurya, A. Kumari, V. K. Agnihotri, V. P. Verma and D. M. Sawant, *Org. Biomol. Chem.*, 2019, 17, 363.
- 49 J. Mou, N. Chen, Y. Zhao, H. Qi, S. Meng, R. Xiang and D. Pei, *Front. Chem.*, 2020, **8**, 2020.
- 50 M. Ghorbani-Choghamarani and M. Norouzi, *New J. Chem.*, 2016, **40**, 6299.
- 51 M. Ghorbani-Choghamarani, M. Nikoorazm and M. Norouzi, *Org. Chem. Res.*, 2018, 4, 11.
- 52 A. Ghorbani-Choghamarani and M. Norouzi, *Appl.* Organomet. Chem., 2016, **30**, 140.
- 53 M. Norouzi and A. Ghorbani-Choghamarani, *React. Kinet.*, *Mech. Catal.*, 2016, **119**, 537.
- 54 P. Moradi, M. Hajjami and B. Tahmasbi, *Polyhedron*, 2020, 175, 114169.
- 55 L. Ronghua, W. Liang, H. Huang, S. Jiang, D. Guo, M. Li, Z. Zhang, A. Ali and J. J. Wang, *J. Appl. Polym. Sci.*, 2018, 135, 46239.

Paper

- Open Access Article. Published on 06 December 2023. Downloaded on 7/29/2025 12:00:40 PM. This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.
- 56 M.-S. Mashhoori and R. Sandaroos, *Sci. Rep.*, 2022, **12**, 15364.
- 57 V. Rama, K. Kanagaraj and K. Pitchumani, *J. Org. Chem.*, 2011, **76**, 9090.
- 58 M. Norouzi, N. Noormoradi and M. Mohammadi, *Nanoscale Adv.*, 2023, **5**, 6594–6605.
- 59 M. Nikoorazm, Z. Rezaei and B. Tahmasbi, *J. Porous Mater.*, 2020, 27, 671.
- 60 B. Tahmasbi and A. Ghorbani-Choghamarani, *Appl. Organomet. Chem.*, 2017, **31**, e3644.
- 61 G. Qi, W. Liu and Z. Bei, Chin. J. Chem., 2011, 29, 131.
- 62 A. Jabbari, P. Moradi and B. Tahmasbi, *RSC Adv.*, 2023, **13**, 8890–8900.
- 63 B. Sreedhar, A. S. Kumar and D. Yada, *Tetrahedron Lett.*, 2011, 52, 3565.
- 64 B. Tahmasbi, M. Nikoorazm, P. Moradi and Y. Abbasi Tyula, *RSC Adv.*, 2022, **12**, 34303.
- 65 A. Ghorbani-Choghamarani, Z. Darvishnejad and M. Norouzi, *Appl. Organomet. Chem.*, 2015, **29**, 707.

- 66 A. Ghorbani-Choghamarani and M. Norouzi, *J. Mol. Catal. A: Chem.*, 2014, **395**, 172.
- 67 C. A. Didó, E. B. Mass, M. B. Pereira, R. Hinrichs, M. G. D'Oca, T. M. Costa and E. V. Benvenutti, *Colloids Surf.*, A, 2020, 589, 124455.
- 68 S. B. Bharate, N. Mupparapu, S. Manda, J. B. Bharate, R. Mudududdla, R. R. Yadav and R. A. Vishwakarma, *Arkivoc*, 2012, **8**, 308.
- 69 L. Shiri, A. Ghorbani-Choghamarani and M. Kazemi, *Appl.* Organomet. Chem., 2016, **31**, 9.
- 70 P. Sivaguru, K. Parameswaran, M. Kiruthiga, P. Vadivel and A. Lalitha, J. Iran. Chem. Soc., 2015, 12, 95.
- 71 M. Koolivand, M. Nikoorazm, A. Ghorbani-Choghamarani,
 R. Azadbakht and B. Tahmasbi, *Sci. Rep.*, 2021, 11, 24475.
- 72 V. V. Gaikwad, V. S. Saptal, K. Harada, T. Sasaki, D. Nishio-Hamane and M. B. M. Bhanage, *ChemNanoMat*, 2018, **6**, 575–582.
- 73 R. Soengas, Y. Navarro, M. J. Iglesias and F. López-Ortiz, Molecules, 2018, 23, 2975.