

**CORRECTION**
[View Article Online](#)  
[View Journal](#) | [View Issue](#)


Cite this: RSC Adv., 2023, 13, 23818

## Correction: Effects of Ni precursors on the formation of Mg–Fe–Ni intermetallic hydrides, kinetics, and reversibility

Palmarin Dansirima,<sup>a</sup> Sophida Thiangviriya,<sup>a</sup> Praphatsorn Plerdsranoy,<sup>a</sup> Narong Chanlek<sup>b</sup> and Rapee Utke<sup>\*a</sup>

DOI: 10.1039/d3ra90070c  
[rsc.li/rsc-advances](http://rsc.li/rsc-advances)

Correction for 'Effects of Ni precursors on the formation of Mg–Fe–Ni intermetallic hydrides, kinetics, and reversibility' by Palmarin Dansirima *et al.*, RSC Adv., 2023, 13, 16926–16934, <https://doi.org/10.1039/D3RA01914D>.

The authors regret that there was an error in Table 1. The correct version of Table 1 is presented below.

**Table 1** Reaction pathways and phase compositions of S1 and S2 during the 1st de/rehydrogenation

Samples	Possible reaction pathways and phase compositions
<b>S1</b>	
As-prepared	$\text{MgH}_2 + \text{Mg}_2\text{FeH}_6 + \text{Mg}_2\text{NiH}_4$
1st desorbed	$\text{Mg}_2\text{FeH}_6 \rightarrow 2\text{MgH}_2 + \text{Fe} + \text{H}_2$ $\text{MgH}_2 \rightarrow \text{Mg} + \text{H}_2$ $\text{Mg}_2\text{NiH}_4 \rightarrow \text{Mg}_2\text{Ni} + 2\text{H}_2$ $\text{Fe} + \text{Ni} \rightarrow \text{Fe–Ni}$ $\text{Mg} + \text{H}_2 \rightarrow \text{MgH}_2$ $2\text{MgH}_2 + \text{Fe} + \text{H}_2 \rightarrow \text{Mg}_2\text{FeH}_6$ $\text{Mg}_2\text{Ni} + 2\text{H}_2 \rightarrow \text{Mg}_2\text{NiH}_4$ $2\text{MgH}_2 + \text{Fe–Ni} \rightarrow \text{Mg}_2\text{NiH}_4 + \text{Fe}^{33}$
1st absorbed	
<b>S2</b>	
As-prepared	$\text{Mg}_2\text{FeH}_6 + \text{Mg}_2\text{NiH}_4 + \text{Fe–Ni}$
1st desorbed	$\text{Mg}_2\text{FeH}_6 \rightarrow 2\text{MgH}_2 + \text{Fe} + \text{H}_2$ $\text{MgH}_2 \rightarrow \text{Mg} + \text{H}_2$ $\text{Mg}_2\text{NiH}_4 \rightarrow \text{Mg}_2\text{Ni} + 2\text{H}_2$ $\text{Fe–Ni}$ (comparable to as-prepared state) $\text{Mg} + \text{H}_2 \rightarrow \text{MgH}_2$ $2\text{MgH}_2 + \text{Fe} + \text{H}_2 \rightarrow \text{Mg}_2\text{FeH}_6$ $x\text{Mg}_2\text{Ni} + (1 - x)\text{Mg}_2\text{FeH}_6 + 3x\text{H}_2 \rightarrow \text{Mg}_2\text{Fe}_{(1-x)}\text{Ni}_x\text{H}_6$ $\text{Fe} + \text{Ni} \rightarrow \text{Fe–Ni}$
1st absorbed	

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

<sup>a</sup>School of Chemistry, Institute of Science, Suranaree University of Technology, Nakhon Ratchasima 30000, Thailand. E-mail: rapee.g@sut.ac.th

<sup>b</sup>Synchrotron Light Research Institute (Public Organization), Nakhon Ratchasima 30000, Thailand

