

Cite this: *Chem. Sci.*, 2023, 14, 4564

All publication charges for this article have been paid for by the Royal Society of Chemistry

## Water molecule switching heterogeneous proton-coupled electron transfer pathway†

Zhonghuan Liu,<sup>b</sup> Wei Peng,<sup>ae</sup> Yuhan Lin,<sup>ae</sup> Xinyu Lin,<sup>b</sup> Shikang Yin,<sup>b</sup> Shuhan Jia,<sup>b</sup> Dongge Ma,<sup>id</sup> Yan Yan,<sup>id</sup>\*<sup>b</sup> Peng Zhou,<sup>id</sup>\*<sup>c</sup> Wanhong Ma\*<sup>ae</sup> and Jincai Zhao<sup>id</sup><sup>ae</sup>

Figuring out the specific pathway of semiconductor-mediated proton-coupled electron transfer (PCET) driven by light is essential to solar energy conversion systems. In this work, we reveal that the amount of adsorbed water molecules determines the photo-induced PCET pathway on the TiO<sub>2</sub> surface through systematic kinetic solvent isotope effect (KSIE) experiments. At low water content (<1.7 wt%), the photo-induced single-proton/single-electron transfer on TiO<sub>2</sub> nanoparticles follows a stepwise PT/ET pathway with the formation of high-energy H<sup>+</sup>/D<sup>+</sup>-O=C or H<sup>+</sup>/D<sup>+</sup>-O-C intermediates, resulting in an inverse KSIE (H/D) ~0.5 with <sup>t</sup>Bu<sub>3</sub>ArO· and KSIE (H/D) ~1 with TEMPO in methanol-*d*<sub>0</sub>/*d*<sub>4</sub> systems. However, at high water content (>2 wt%), the PCET reaction follows a concerted pathway with a lower energy barrier, leading to normal KSIEs (H/D) ≥ 2 with both reagents. *In situ* ATR-FTIR observation and DFT calculations suggest that water molecules' existence significantly lowers the proton/electron transfer energy barrier, which coincides with our experimental observations.

Received 23rd December 2022

Accepted 30th March 2023

DOI: 10.1039/d2sc07038c

rsc.li/chemical-science

## Introduction

Proton-coupled electron transfer (PCET) reactions are pervasive in natural energy transfer processes, including photosynthesis, nitrogen fixation, and respiration.<sup>1–4</sup> Metal oxide semiconductor (*e.g.*, ZnO, TiO<sub>2</sub>, and SnO<sub>2</sub>) nanoparticle-mediated photo-induced electron transfer has been directly proven to be a proton-coupled process.<sup>5</sup> Figuring out the internal relation between specific intermediate reaction pathways and the interfacial chemical environment is critical to the profound understanding and future design of targeted semiconductor nanoparticle-based photocatalytic systems with high efficiency.

Typically, under continuous light irradiation, metastable photo-induced electrons can be generated and constantly trapped on oxide semiconductor (*e.g.*, TiO<sub>2</sub>) nanoparticles in small-molecule hole-scavengers (*e.g.*, H<sub>2</sub>O, amines, or alcohols).<sup>6,7</sup> Stabilized by protons (H<sup>+</sup>), photo-induced electrons (e<sub>cb</sub><sup>-</sup>) on TiO<sub>2</sub> nanoparticles were proved to be trapped at the outer surface as

Ti<sub>4c</sub><sup>3+</sup> species,<sup>8</sup> which can be consumed *in situ* or *ex situ* as reactive reducing reagents to accomplish many valuable and challenging chemical reactions without extra high-temperature and high-pressure conditions.<sup>9–11</sup> The corresponding intermediate reaction pathway directly determines such a liquid/solid heterogeneous PCET process driven by photo-excitation. Compared to the stepwise electron-transfer followed by the proton-transfer (ET/PT) pathway and *vice versa* (PT/ET), concerted gain or loss of coupled e<sub>cb</sub><sup>-</sup> and H<sup>+</sup> (CPET) typically favors redox reactions. The advantage of such a reaction pathway is that it avoids high-energy intermediates caused by localized charge accumulation,<sup>12–14</sup> which makes it consistently exhibit a lower reaction energy barrier.<sup>13</sup> Generally, the PCET pathway selectivity was long considered only determined by the proton concentration and specific activation path built by the reactive sites in reaction systems.<sup>15</sup> Previous observations have revealed that the reduction potential of e<sub>cb</sub><sup>-</sup> varies sharply with the proton concentration near TiO<sub>2</sub> electrodes, which exhibit different reactivities.<sup>16</sup> Our previous work further distinguished the binding energy difference that reflects energy levels of electron trapping states on TiO<sub>2</sub> nanoparticles between proton (1.3 eV) and proton-free (1.8 eV) systems by synchrotron radiation UV photoelectron spectroscopy.<sup>17</sup> Furthermore, hybrid functional periodic density functional theory (DFT) calculations elucidate the impact of proton-coupled defects on the bond dissociation-free energies (BDFEs) of O–H bonds on anatase TiO<sub>2</sub> surfaces, which are directly related to interfacial PCET thermochemistry.<sup>18</sup> In dye-sensitized solar cells, the PCET model has been applied in monitoring geometric parameters, excitation, and electronic structures of free and Ti<sup>4+</sup>-bound squaraine dye solar cells.<sup>19</sup>

<sup>a</sup>Key Laboratory of Photochemistry, Beijing National Laboratory for Molecular Sciences, Institute of Chemistry, Chinese Academy of Sciences, Beijing, 100190, PR China. E-mail: whma@iccas.ac.cn

<sup>b</sup>Institute of Green Chemistry and Chemical Technology, School of Chemistry and Chemical Engineering, Jiangsu University, Zhenjiang 212013, PR China. E-mail: dgy5212004@163.com

<sup>c</sup>Department of Electrical Engineering and Computer Science, University of Michigan, Ann Arbor, MI 48109, USA. E-mail: dpzhou@umich.edu

<sup>d</sup>Department of Chemistry, College of Chemistry and Materials Engineering, Beijing Technology and Business University, Beijing 100048, China

<sup>e</sup>University of Chinese Academy of Sciences, Beijing, 100049, China

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d2sc07038c>



However, in a proton-excessive PCET system, *e.g.*, the most common solar energy conversion path of  $\text{H}_2\text{O} \rightarrow \text{H}_2$  or  $\text{CO}_2 + \text{H}_2\text{O} \rightarrow \text{C}_x\text{H}_y\text{O}_z$  involving water participation, the impact of the aqueous interfacial chemical environment on the photo-induced PCET reaction pathway on semiconductor nanoparticles remains unclear. For instance, in some photo-reduction scenarios (*e.g.*, dehalogenation of aromatic halides,  $\text{H}_2$  production, and  $\text{CO}_2$  reduction), reaction efficiencies on semiconductor nanoparticle catalysts in  $\text{H}_2\text{O}$ /organic mixtures were reported to be much higher than those in the pure organic phase.<sup>20–22</sup> Although the effect of water as a direct reactant on photocatalytic performances has been reported in many terms, we still lack a consensual in-depth understanding of the cause of such phenomena. Therefore, determining whether these water-enhanced performances were directly achieved by regulating the PCET pathway is of utmost importance.


## Results and discussion

Herein, by using a 2,4,6-tri-*tert*-butylphenoxy radical ( ${}^t\text{Bu}_3\text{ArO}\cdot$ ) and 2,2,6,6-tetramethyl-piperidin-1-yl-oxyl (TEMPO) as single-electron/single-proton reagents and isotopically labeled methanol- $d_0/d_4$  as both the electron/proton donor and solvent, a series of delicate kinetic solvent isotope effect (KSIE (H/D)) experiments were carried out on  $\text{TiO}_2$  nanoparticles as catalysts (Scheme 1). We found an unexpected switch of the photo-induced PCET pathway from the stepwise PT/ET to CPET manipulated by a trace amount of adsorbed water molecules on  $\text{TiO}_2$  nanoparticles. *In situ* attenuated total reflection Fourier transform infrared spectroscopy (ATR-FTIR) observations confirmed a strong interaction between water molecules and photo-induced surface protons (mainly from the oxidation of methanol solvent) in the presence of  $e_{\text{cb}}^-$ . Such an interreaction directly affects the bond-dissociation of these protons from the  $\text{TiO}_2$  surface and favors proton-coupled reduction *via* a CPET pathway. Density functional theory (DFT) calculations were further employed to investigate the impact of water molecules on the proton bond-dissociation and electron transfer on the  $\text{TiO}_2$  surface.

## Determining the PCET pathway by KSIE (H/D) measurements

We first performed the KSIE (H/D) experiments on  $\text{TiO}_2$  nanoparticles under UV with the well-reported single-electron/single-proton radical reduction of  ${}^t\text{Bu}_3\text{ArO}\cdot$  ( ${}^t\text{Bu}_3\text{ArO}\cdot + \text{H}^+ + e_{\text{cb}}^- \rightarrow {}^t\text{Bu}_3\text{ArOH}$ ) and TEMPO ( $\text{TEMPO} + \text{H}^+ + e_{\text{cb}}^- \rightarrow \text{TEMPOH}$ )<sup>15,23</sup> by using isotopically labeled methanol- $d_0/d_4$  as the H/D solvent and electron/proton donor ( $\text{CH}_3\text{OH} + h\nu_{\text{cb}}^+ \rightarrow \text{CH}_2\text{O} + \text{H}^+$ ).  ${}^1\text{H}$  NMR spectra (Figure S1†) and mass spectra (Figure S2†) confirm no other major products except for the reduction product. All measurements were carried out under anaerobic conditions.

The KSIE (H/D) was determined using concentration-time profiles of  ${}^t\text{Bu}_3\text{ArO}\cdot/\text{TEMPO}$  that were *in situ* monitored in methanol- $d_0/d_4$  (H/D systems) by electron spin resonance (ESR) spectroscopy (for details of the KSIE experiments, see the ESI†). Control experiments were performed without  $\text{TiO}_2$  nanoparticles, confirming no reaction without a catalyst (Figure S3†). The commonly used commercial  $\text{TiO}_2$  nanoparticles (anatase-phase, without further treatment) were employed as a typical semiconductor nanoparticle catalyst. A direct normal KSIE (H/D) of 2.18 was observed with  ${}^t\text{Bu}_3\text{ArO}\cdot$  between methanol- $d_0/d_4$  systems (Figure S4†), suggesting a one-step CPET reaction pathway on pristine  $\text{TiO}_2$  nanoparticles.<sup>24</sup> Moreover, with TEMPO as the reagent, a normal KSIE (H/D) of 3.73 was observed (Figure S5†). Our observation is consistent with the previously reported result by the Mayer group<sup>5</sup> that the single-electron/single-proton transfer induced by photo-generated  $\text{H}^+/e_{\text{cb}}^-$  on  $\text{TiO}_2$  and  $\text{ZnO}$  nanoparticles follows a concerted pathway. However, when freshly high-temperature (773 K) calcined  $\text{TiO}_2$  nanoparticles were used as catalysts under otherwise identical conditions, the slower reaction kinetics of  ${}^t\text{Bu}_3\text{ArO}\cdot$  in methanol- $d_0$  compared to methanol- $d_4$  were observed with an unexpected inverse KSIE (H/D) of 0.34 (Figure S6†), indicating that the PCET reaction pathway was significantly changed. Noticeably, only when atomic hybrid modes adjacent to -H/D coordination were changed from  $\text{sp}^2$  to  $\text{sp}^3$ , the subsequent bond-dissociation kinetic isotope effect (KIE) would appear with an inverse value, indicating the



**Scheme 1** Mechanisms of  $\text{TiO}_2$  photocatalysis: a well-established CPET pathway induced free-radical reaction for (a)  ${}^t\text{Bu}_3\text{ArO}\cdot$  and (b) TEMPO substrates (left) and PTET pathway induced free-radical reaction (right).



formation of a certain protonation intermediate configuration. Moreover, with TEMPO as the reagent, the KSIE (H/D) on calcined TiO<sub>2</sub> nanoparticles was approaching  $\sim 1$  (Figure S7†) because the transformation of TEMPO into TEMPOH does not involve the hybridization change from sp<sup>2</sup> to sp<sup>3</sup>. Control experiments have been performed with TiO<sub>2</sub> nanoparticles calcined in an air, oxygen, or argon atmosphere. An inverse KSIE (H/D)  $\leq 0.5$  with <sup>t</sup>Bu<sub>3</sub>ArO· and KSIE (H/D)  $\sim 1$  with TEMPO were always observed on calcined TiO<sub>2</sub> nanoparticles regardless of atmosphere (Figures S8 and S9†). These results indicate that other effects from calcination rather than defect vacancies determine the switch of the KSIE (H/D) and corresponding PCET pathways.

The XRD result shows that the crystalline structure of anatase TiO<sub>2</sub> nanoparticles was not changed after calcination and the crystallite diameter did not severely change (Figure S10 and Table S1†), which further excludes the impact of rutile formation as the major effect. We further compared the sedimentation behavior of a pristine/calcined TiO<sub>2</sub> (2.5 g L<sup>-1</sup>) suspension by *in situ* UV-vis optical-fiber spectroscopy. As shown in Figure S11,† the light transmittance hardly changed within 90 s in both systems, suggesting that sedimentation is not the major effect for the KIE in our system.

The decreased KSIE value caused by calcination would recover to the normal KSIE value after the calcined TiO<sub>2</sub> catalyst is stored in the air for one week (Figures S12 and S13†). We argue that the loss of trace water on TiO<sub>2</sub> nanoparticles after calcination is the determining factor for the KSIE <1. Through the *in situ* diffuse reflection infrared Fourier transform spectroscopy (DRIFTS) characterization of TiO<sub>2</sub> nanoparticles during calcination (298–773 K), we found that the featured signals of adsorbed water molecules (O–H stretching vibration band at 3200 cm<sup>-1</sup> and H–O–H bending vibration at 1642 cm<sup>-1</sup>) severely decreased until they wholly disappeared with the increasing temperature. When the temperature exceeded 773 K, the loss of dissociated water (O–H stretching vibration band at 3694/3459 cm<sup>-1</sup> without 1642 cm<sup>-1</sup> bending vibration) was observed (Figure S14a†). However, with the temperature further returned to room temperature, although the ambient humidity can recover a certain amount of water adsorption on TiO<sub>2</sub>

nanoparticles, it would never be able to reach the pre-calcination level (Figure S14b†), demonstrating the inevitable loss of water content in TiO<sub>2</sub> nanoparticles after high-temperature calcination. Moreover, by using thermogravimetric analysis (TGA) measurements, we found that the water content in pristine TiO<sub>2</sub> nanoparticles long stored in the air was  $\sim 3.76$  wt% before calcination. In contrast, after 773 K calcination and returning to room temperature, the water content of the freshly calcined TiO<sub>2</sub> nanoparticles was significantly reduced to  $\sim 1.33$  wt% (Figure S15†). To directly demonstrate the relationship between the water content of the catalyst and the KSIE (H/D) of photo-induced PCET reactions, we prepared a series of TiO<sub>2</sub> nanoparticle catalysts with well-controlled water contents (1.33–3.80 wt%) by treating dried TiO<sub>2</sub> nanoparticles in airflows with different humidities (Figure S16†) (for preparation methods see the ESI†). The coverage of water layers was calculated according to the surface Ti atoms on TiO<sub>2</sub> nanoparticles (in terms of a perfect anatase {101} surface) and is summarized in Table S2,† corresponding to an increasing water coverage from  $\sim$  one layer to  $\sim$  three layers. With the increasing water content of TiO<sub>2</sub> nanoparticles, the KSIE (H/D) with <sup>t</sup>Bu<sub>3</sub>ArO· significantly increased from an inverse value of 0.34 to a normal value of 3.33 (Fig. 1a). In contrast with the KSIE(H/D) with <sup>t</sup>Bu<sub>3</sub>ArO·, the KSIE (H/D) with TEMPO increased from a recessive value of  $\sim 1$  to a normal value of 3 (Fig. 1b), which provides solid evidence that the KSIE (H/D) value was directly determined using the water content in the TiO<sub>2</sub> nanoparticle catalyst. Reaction rate parameters in methanol ( $k(H)$ ) with both radical reagents show a similar trend that increases with the increasing water content (Fig. 1a and b), which is consistent with the water-determined KSIE (H/D). We further employed a well-reported Fe(III)-1,10-phenanthroline spectrometric titration method (for details of the titration method see the ESI†) to quantify the oxidation of CH<sub>3</sub>OH/CD<sub>3</sub>OD on different TiO<sub>2</sub> samples.<sup>26</sup> As shown in Figure S17,† KIEs for methanol-d<sub>0</sub>/d<sub>4</sub> oxidation on both dry (water content  $\sim 1.3$  wt%) and wet (water content  $\sim 3.8$  wt%) TiO<sub>2</sub> are identical at  $\sim 1.3$ , indicating that the changed KIE value with increasing water content in the TiO<sub>2</sub> catalyst is not from the methanol oxidation.




Fig. 1 Profiles of the KSIE (H/D) and reaction rate parameter  $k(H)$  as functions of the water content in TiO<sub>2</sub> nanoparticles with (a) <sup>t</sup>Bu<sub>3</sub>ArO· and (b) TEMPO as the single-proton/single-electron acceptors under UV irradiation. Error bars were obtained by repeating the experiment three times under identical conditions.




The water-determined KSIE (H/D) reversal directly reflects the change of the photo-induced PCET reaction pathway on TiO<sub>2</sub> nanoparticles. Typically, the <sup>t</sup>Bu<sub>3</sub>ArO· radical has three possible configurations, as shown in Scheme 2a. Type I is the O–C single bond radical structure, while type II and type III are O=C double bond structures. Based on the reported single-crystalline structure analysis, type III is the major contributor to the stable <sup>t</sup>Bu<sub>3</sub>ArO· radical, *i.e.*, in the O=C double bond configuration.<sup>27</sup> Therefore, the single-proton/single-electron transfer of <sup>t</sup>Bu<sub>3</sub>ArO· → <sup>t</sup>Bu<sub>3</sub>ArOH is accompanied by the dissociation of the C=O double bond and the hybrid configuration transition of O=C sp<sup>2</sup> → O–C sp<sup>3</sup>. The inverse KIE from sp<sup>2</sup>–sp<sup>3</sup> hybridization is a very classic phenomenon in reaction kinetics.<sup>25,28</sup> The secondary KIE is mainly due to the difference in bending vibrations. Due to symmetry, the in-plane and out-of-plane bending of sp<sup>3</sup> hybridized carbon atoms are equivalent. The in-plane bending vibrations of sp<sup>2</sup> hybridized carbon or oxygen atoms are stiffer than out-of-plane bending vibrations (there is a minor site resistance for out-of-plane bending), leading to a more significant difference in zero-point energy in the O–H and O–D bonding in the reaction of the hybrid state change. In the sp<sup>3</sup> state as the substrate, the O–D bond has lower zero-point energy and needs to cross a larger energy barrier to reach the intermediate state, while in the sp<sup>2</sup> state it is the opposite (Fig. S18†). In this process, an inverse KIE (H/D) can only occur when the protonated intermediate of H<sup>+</sup>/D<sup>+</sup>–O=C is formed and subsequently converted to the H/D–O–C product by the further e<sub>cb</sub><sup>–</sup> transfer (Scheme 2b), reflecting a stepwise PT/ET pathway at low water content (Scheme 2b). Conversely, at high water content, <sup>t</sup>Bu<sub>3</sub>ArO· was directly reduced to <sup>t</sup>Bu<sub>3</sub>ArOH through one-step CPET without forming such H<sup>+</sup>/D<sup>+</sup>–O sp<sup>2</sup> protonated intermediates, resulting in a normal KIE ≥ 2 from the direct bond-dissociation of protons

from bridging-O sites on the TiO<sub>2</sub> surface (Scheme 2b).<sup>29–31</sup> However, due to the high electronegativity and coordination limits of the adjacent N atoms that hinder the transfer of the unpaired electron from the O atom to the carbon ring, TEMPO as another single-proton/single-electron acceptor always retains the N–O single bond free radical structure without the O=C → O–C transition during the reaction. Thus, even with the PT intermediate formed at low water content, a KIE ~1 rather than an inverse KIE should be observed (Scheme S2†), entirely consistent with our experimental results (Fig. 1b). Our KSIE experiments show that the participation of water can directly switch the photo-induced PCET pathway on the TiO<sub>2</sub> nanoparticle catalyst from water-free stepwise PT/ET to water-participated CPET.

### In situ ATR-FTIR experiments

Isotopic-labeled ATR-FTIR spectroscopy was further used to *in situ* observe water molecules on the TiO<sub>2</sub> surface and their interaction with photo-induced protons/electrons under continuous UV irradiation. Setting the equilibrium conditions as the blank background, a positive or negative IR response reflects the intermediate species' gradual gain or loss at the methanol-*d*<sub>0</sub>/*d*<sub>4</sub> (CH<sub>3</sub>OH/CD<sub>3</sub>OD)/H<sub>2</sub>O/TiO<sub>2</sub> interface during the reaction. Before measurements, TiO<sub>2</sub> nanoparticles, as the typical catalyst sample, were first calcined at 773 K for 30 minutes and then cooled to room temperature under a constant dried nitrogen flow. In the water-participated system, the calcined TiO<sub>2</sub> sample was further treated in water vapor under a continuous nitrogen flow for 15 min, resulting in a water content of 3.80 wt%. The experiments were carried out in a nitrogen atmosphere to prevent trapped e<sub>cb</sub><sup>–</sup> from being quenched by oxygen.

As shown in Fig. 2a, when TiO<sub>2</sub>/CH<sub>3</sub>OH was used as the ATR-FTIR sample and irradiated *in situ* with a 365 nm LED lamp, a broad negative peak at 3272 cm<sup>–1</sup> and signal bands at 2941 cm<sup>–1</sup> and 2830 cm<sup>–1</sup> emerged from the background and increased in intensity with increasing irradiation time, corresponding to the stretching vibration signals of O–H and C–H in methanol and representing the oxidation of methanol. This is further verified by the ground-state signal on the TiO<sub>2</sub>/CH<sub>3</sub>OH system. (Figure S20†). Moreover, a strong baseline upshift absorption band from 2500 cm<sup>–1</sup> to 1100 cm<sup>–1</sup> was observed, which is consistent with our previous observations and can be ascribed to the absorption of trapped e<sub>cb</sub><sup>–</sup> on the TiO<sub>2</sub> nanoparticles.<sup>17,26,32,33</sup> These trapped e<sub>cb</sub><sup>–</sup> were localized in the surface defect layer in the form of four-coordinated Ti<sub>4c</sub><sup>3+</sup> under the stabilization of protons provided by the methanol oxidation.<sup>8,17,31</sup> When the water vapor was first introduced to TiO<sub>2</sub> nanoparticles, a broad negative absorption of O–H at 3220 cm<sup>–1</sup> and a mild negative peak at 1677 cm<sup>–1</sup> corresponding to the water molecule H–O–H bending vibration emerged at the TiO<sub>2</sub>/H<sub>2</sub>O/CH<sub>3</sub>OH interface under continuous UV irradiation. These signals are accompanied by the methanol oxidation and the baseline upshift from the absorption of trapped e<sub>cb</sub><sup>–</sup> (Fig. 2b), further verified by the ground-state water and methanol signal on TiO<sub>2</sub> (Figure S20†). More importantly,



Scheme 2 (a) Configurations of the <sup>t</sup>Bu<sub>3</sub>ArO· resonance structure with type III as the major contributor;<sup>27</sup> (b) schematic diagram of the CPET and PT/ET reaction pathways of the single-proton/single-electron transfer on TiO<sub>2</sub> with <sup>t</sup>Bu<sub>3</sub>ArO· as the acceptor.






Fig. 2 ATR-FTIR spectra *in situ* monitored at (a)  $\text{TiO}_2/\text{CH}_3\text{OH}$ ; (b)  $\text{TiO}_2/\text{H}_2\text{O}/\text{CH}_3\text{OH}$ ; (c)  $\text{TiO}_2/\text{D}_2\text{O}/\text{CD}_3\text{OD}$ ; and (d)  $\text{TiO}_2/\text{CD}_3\text{OD}$  interface under constant 365 nm (3 W, LED) irradiation.

a positive peak at  $1633\text{ cm}^{-1}$  corresponding to the H–O–H bending vibration with a lower frequency (from  $1677\text{ cm}^{-1}$ ) and a positive absorption band of O–H stretching vibration at  $3594\text{ cm}^{-1}$  with a higher frequency (from  $3220\text{ cm}^{-1}$ ) emerged. These vibrations reflect a strong interaction between water molecules and protons/ $e_{\text{cb}}^-$  from methanol oxidation on  $\text{TiO}_2$  nanoparticles. Such a strong interaction causes the stretching vibration of the water to be faster while the bending vibration slows down due to the steric effect. All these effects manifested as a change in infrared absorption wavenumbers. The negative peak at  $3707\text{ cm}^{-1}$  corresponds to the terminal hydrogen on  $\text{TiO}_2$ .<sup>32–34</sup> Moreover, when the deuterium-labeled  $\text{TiO}_2/\text{D}_2\text{O}/\text{CD}_3\text{OD}$  sample is used, the strong interaction between water and protons remains. The O–H stretching vibration peak shifts towards higher frequencies ( $2394\text{ cm}^{-1} \rightarrow 2713\text{ cm}^{-1}$ ) and H–O–H bending vibration towards lower frequencies ( $1232\text{ cm}^{-1} \rightarrow 1130\text{ cm}^{-1}$ ) in terms of H/D replacement effects (Fig. 2c), which is consistent with the ground-state methanol- $\text{d}_4$  and  $\text{D}_2\text{O}$  signals on  $\text{TiO}_2$  (Figure S20†). The displacement of the deuterium-labeled proton peak indicates that the proton peak is indeed produced by the interaction of water and protons. In water-free systems, similar peak shift phenomena were no longer observed on  $\text{TiO}_2/\text{CH}_3\text{OH}$  or  $\text{TiO}_2/\text{CD}_3\text{OD}$  samples (Fig. 2a and d).

To further confirm that the observed peak shift was due to the interaction between water molecules and proton/ $e_{\text{cb}}^-$  on the  $\text{TiO}_2$  surface, hydrochloric acid was used as an external proton donor to replace methanol, forming an  $\text{H}_2\text{O}/\text{HCl}/\text{TiO}_2$  interface (pH = 6). When the light was off, the addition of HCl only induced mild water desorption from the  $\text{TiO}_2$  surface with negative peaks at  $3377\text{ cm}^{-1}$  and  $1644\text{ cm}^{-1}$ , and no positive peaks were observed (Figure S19b†), indicating that water

molecules were hardly affected by only the presence of additional protons. However, when the light was on, the negative peak at  $1704\text{ cm}^{-1}$  and positive peak at  $1663\text{ cm}^{-1}$  that correspond to the shift of H–O–H bending vibration towards lower frequencies were immediately observed, as well as the negative band at  $3333\text{ cm}^{-1}$  and the emerging positive band at  $3555\text{ cm}^{-1}$  that correspond to the shift of O–H stretching vibration towards higher frequencies (Figure S19d†), which is identical to the observed phenomenon at the  $\text{TiO}_2/\text{H}_2\text{O}/\text{CH}_3\text{OH}$  interface. Moreover, a broad negative band at around  $2964\text{ cm}^{-1}$  emerged, which can be ascribed to the destruction of hydrogen-bonds between water molecules, further demonstrating the changed water molecules in the presence of protons and photo-induced  $e_{\text{cb}}^-$ . Without the addition of HCl, no other peaks appeared except for the molecular water H–O–H bending vibration at  $1640\text{ cm}^{-1}$  and the molecular water O–H stretching vibration at  $3210\text{ cm}^{-1}$  under constant irradiation (Figure S19a†). We further reduced pH to 5 and repeated the ATR-FTIR experiment. As shown in Figure S19e,† the observed phenomenon is consistent with pH = 6. Furthermore, it offers a more significant blueshift signal of –OH stretching vibration and a redshift signal of H–O–H bending vibration. Such a result indicates that high proton concentrations accelerate the formation of hydrated protons, which is consistent with our conclusion. Our ATR-FTIR results solidly confirm a strong interaction between water and protons/ $e_{\text{cb}}^-$  intermediates on the  $\text{TiO}_2$  surface, possibly achieved by destroying hydrogen bonds between water or forming hydrated protons<sup>35</sup> which would undoubtedly affect the subsequent PCET dynamics.

Typically, protons tend to be localized at bridging –O atoms on  $\text{TiO}_2$  nanoparticles. At the same time, the trapping of  $e_{\text{cb}}^-$  at neighboring Ti atoms on  $\text{TiO}_2$  nanoparticles under UV in the presence of methanol to quench photo-induced holes occurs.<sup>31</sup> In that case, a stepwise PT/ET route with the formation of high-energy protonation intermediates was inevitable. However, according to our observations in this work, surface protons/ $e_{\text{cb}}^-$  can form a strong interaction with water (in the form of hydrogen bonds or hydrated protons), serving as a reaction relay to favor the concerted transfer of proton/ $e_{\text{cb}}^-$  pairs by avoiding the formation of high-energy intermediates. In other words, the presence of water lowered the reaction energy barrier of photo-induced PCET on  $\text{TiO}_2$  nanoparticles by switching the reaction pathway from stepwise PT/ET to CPET.

### DFT calculations

DFT calculations were further employed to investigate the impact of water molecules on the single-proton/single-electron transfer on the  $\text{TiO}_2$  surface (for details of the computational methods, see the ESI†). As shown in Fig. 3a, the calculated free energy profiles of electron and proton transfer suggest that the existence of water molecules on the  $\text{TiO}_2$  surface can significantly promote the transfer of electrons and protons from the surface –OH group due to more negative reaction energy ( $-1.36\text{ eV}$ ) in comparison with the water-free system ( $-0.63\text{ eV}$ ). The projected density of states (PDOS) profiles of the surface –OH group further indicate that the edge line of the main peak of






Fig. 3 Redox reactions were simulated on (a) water-free system and (b) water-participated system. (c) Free energy profiles of hydrogen in both water-participated/water-free systems at pH = 7 and  $U = 0$  V vs. SHE; and (d) PDOS of the surface-OH group in both water-participated/water-free systems. The dashed line stands for the Fermi level.

O 2p states in the -OH group with a water layer on the  $\text{TiO}_2$  surface is closer to the Fermi level than that without a water layer, which is often active in the catalysis reaction. This also suggests the higher reaction activity of the -OH group in the presence of a water layer on the  $\text{TiO}_2$  surface (Fig. 3b). As a result, the surface -OH group with neighboring water is more active than that without water, which coincides with our experimental observations.

## Conclusion

An unexpected water-controlled switch of the photo-induced single-proton/single-electron transfer pathway on  $\text{TiO}_2$  nanoparticles was discovered. On dried  $\text{TiO}_2$  nanoparticles (water content <1.7 wt%,  $\sim$ monolayer coverage), the photo-induced PCET follows a stepwise PT/ET pathway with the formation of a high-energy protonation intermediate, resulting in an inverse KSIE (H/D)  $\sim$ 0.5 with  ${}^t\text{Bu}_3\text{ArO}\cdot$  and KSIE (H/D)  $\sim$ 1 with TEMPO in methanol- $d_0/d_4$  systems. However, in the presence of a trace amount of water (>2 wt% in  $\text{TiO}_2$ ,  $\sim$ two layer coverage), the single-proton/single-electron transfer reaction on  $\text{TiO}_2$  nanoparticles follows a concerted pathway with a lower energy barrier, leading to a normal KSIE (H/D)  $\geq$ 2 with both reagents. Our work is the first to clarify the genuine impact of water on the PCET pathway responsible for the performance enhancement in water-participated systems in terms of PCET, which dispels previous confusion on this issue and can be used for the design of efficient semiconductor-based photocatalytic systems.

## Data availability

The data supporting the findings of this study are available within the article and its ESI† files. All other relevant source

data are available from the corresponding author upon request following the data management specifications of Jiangsu University and University of Michigan. Source data are provided with this paper.

## Author contributions

Z. Liu: investigation, data curation, and writing of the original draft. W. Peng, Y. Lin, X. Lin, S. Yin, S. Jia, and D. Ma: resources. Y. Yan and P. Zhou: writing-review. Y. Yan, W. Ma, and J. Zhao: conceptualization, supervision, writing-review and editing.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

We gratefully acknowledge the financial support of the National Natural Science Foundation of China (Grant No. 21776117, 22176192, and 21806060).

## References

- 1 E. Tsui, A. J. Metrano, Y. Tsuchiya and R. R. Knowles, *Angew. Chem., Int. Ed.*, 2020, **59**, 11845–11849.
- 2 G. J. Choi, Q. Zhu, D. C. Miller, C. J. Gu and R. R. Knowles, *Nature*, 2016, **539**, 268–271.
- 3 K. T. Tarantino, P. Liu and R. R. Knowles, *J. Am. Chem. Soc.*, 2013, **135**, 10022–10025.
- 4 J. Shangguan, A. J. R. Hensley, M. V. Gradiski, N. Pfriend, J. McEwen, R. H. Morris and Y. H. C. Chin, *ACS Catal.*, 2019, **9**, 1763–1778.



- 5 J. N. Schrauben, R. Hayoun, C. N. Valdez, M. Braten, L. Fridley and J. M. Mayer, *Science*, 2012, **336**, 1298–1301.
- 6 C. C. Mercado, F. J. Knorr and J. L. McHale, *J. Phys. Chem. C*, 2008, **112**, 12786–12794.
- 7 C. C. D. Zhao, C. Yu, W. Ma and J. Zhao, *J. Phys. Chem. C*, 2009, **113**, 13160–13165.
- 8 F. G. Santomauro, A. Lubcke, J. Rittmann, E. Baldini, A. Ferrer, M. Silatani, P. Zimmermann, S. Grubel, J. A. Johnson, S. O. Mariager, P. Beaud, D. Grolimund, C. Borca, G. Ingold, S. L. Johnson and M. Chergui, *Sci. Rep.*, 2015, **5**, 14834.
- 9 C. N. Valdez, M. Braten, A. Soria, D. R. Gamelin and J. M. Mayer, *J. Am. Chem. Soc.*, 2013, **135**, 8492–8495.
- 10 S. Qiao, Y. Zhou, H. Hao, X. Liu, L. Zhang and W. Wang, *Green Chem.*, 2019, **21**, 6585–6589.
- 11 S. Zeng, L. Zhang, W. Wang, D. Shao and H. Hao, *Phys. Chem. Chem. Phys.*, 2017, **19**, 29053–29056.
- 12 D. R. Weinberg, C. J. Gagliardi, J. F. Hull, C. F. Murphy, C. A. Kent, B. C. Westlake, A. Paul, D. H. Ess, D. G. McCafferty and T. J. Meyer, *Chem. Rev.*, 2012, **112**, 4016–4093.
- 13 R. Tyburski, T. Liu, S. D. Glover and L. Hammarstrom, *J. Am. Chem. Soc.*, 2021, **143**, 560–576.
- 14 A. A. Stuchebrukhov and S. H. Schiffer, *Chem. Rev.*, 2010, **110**, 6939–6960.
- 15 C. Chen, T. Shi, W. Chang and J. Zhao, *ChemCatChem*, 2015, **7**, 724–731.
- 16 G. J. Meyer and P. Qu, *Langmuir*, 2001, **17**, 6720–6728.
- 17 Y. Yan, W. Shi, W. Peng, Y. Lin, C. Zhang, L. Li, Y. Sun, H. Ju, J. Zhu, W. Ma and J. Zhao, *Chem. Commun.*, 2019, **2**, 88.
- 18 R. E. Warburton, J. M. Mayer and S. H. Schiffer, *J. Phys. Chem. Lett.*, 2021, **12**, 9761–9767.
- 19 S. G. Aziz, O. I. Osman, S. A. Elroby, W. M. I. Hassan, A. Jedidi and R. H. Hilal, *Struct. Chem.*, 2018, **29**, 983–997.
- 20 M. Lei, Z. Wang, L. Zhu, W. Nie and H. Tang, *Appl. Catal., B*, 2020, **261**, 118236.
- 21 Y. Kuramochi, M. Kamiya and H. Ishida, *Inorg. Chem.*, 2014, **53**, 3326–3332.
- 22 Y. Li, P. Ren, D. Zhang, W. Qiao, D. Wang, X. Yang, X. Wen, M. H. Rummeli, H. Niemantsverdriet, J. P. Lewis, F. Besenbacher, H. Xiang, Y. Li and R. Su, *ACS Catal.*, 2021, **11**, 4338–4348.
- 23 C. Wang, H. Groenzin and M. J. Shultz, *J. Am. Chem. Soc.*, 2005, **127**, 9736–9744.
- 24 Y. Lin, Y. Yan, W. Peng, X. Qiao, D. Huang, H. Ji, C. Chen, W. Ma and J. Zhao, *J. Phys. Chem. Lett.*, 2020, **11**, 3941–3946.
- 25 W. Chang, C. Sun, X. Pang, H. Sheng, Y. Li, H. Ji, W. Song, C. Chen, W. Ma and J. Zhao, *Angew. Chem., Int. Ed.*, 2015, **54**, 2052–2056.
- 26 Y. Yan, W. Shi, Z. Yuan, S. He, D. Li, Q. Meng, H. Ji, C. Chen, W. Ma and J. Zhao, *J. Am. Chem. Soc.*, 2017, **139**, 2083–2089.
- 27 V. W. Manner, T. F. Markle, J. H. Freudenthal, J. P. Roth and J. M. Mayer, *Chem. Commun.*, 2008, 256–258.
- 28 D. Dougherty and E. Anslyn, *Modern Physical Organic Chemistry*, 2005.
- 29 Q. Guo, C. Zhou, Z. Ma and X. Yang, *Adv. Mater.*, 2019, **31**, e1901997.
- 30 Q. Guo, Z. Ma, C. Zhou, Z. Ren and X. Yang, *Chem. Rev.*, 2019, **119**, 11020–11041.
- 31 Q. Guo, C. Zhou, Z. Ma, Z. Ren, H. Fan and X. Yang, *Chem. Soc. Rev.*, 2016, **45**, 3701–3730.
- 32 J. A. Moss, S. H. Szczepankiewicz and M. R. Hoffmann, *J. Phys. Chem. B*, 2002, **106**, 2922–2927.
- 33 D. S. Warren and A. J. McQuillan, *J. Phys. Chem. B*, 2004, **108**, 19373–19379.
- 34 H. Sheng, H. Zhang, W. Song, H. Ji, W. Ma, C. Chen and J. Zhao, *Angew. Chem., Int. Ed.*, 2015, **54**, 5905–5909.
- 35 Q. Yu, W. B. Carpenter, N. H. C. Lewis, A. Tokmakoff and J. M. Bowman, *J. Phys. Chem. B*, 2019, **123**, 7214–7224.

