



Cite this: *Chem. Sci.*, 2023, 14, 214

## Correction: An exchange coupled *meso*–*meso* linked vanadyl porphyrin dimer for quantum information processing

Davide Ranieri,<sup>a</sup> Fabio Santanni,<sup>a</sup> Alberto Privitera,<sup>a</sup> Andrea Albino,<sup>a</sup> Enrico Salvadori,<sup>b</sup> Mario Chiesa,<sup>b</sup> Federico Totti,<sup>a</sup> Lorenzo Sorace\*<sup>a</sup> and Roberta Sessoli<sup>a</sup>

DOI: 10.1039/d2sc90246j

Correction for 'An exchange coupled *meso*–*meso* linked vanadyl porphyrin dimer for quantum information processing' by Davide Ranieri *et al.*, *Chem. Sci.*, 2022, <https://doi.org/10.1039/d2sc04969d>.

rsc.li/chemical-science

During the editorial production of the finished article, the graphics associated with Fig. 3(a) and (b) were inadvertently switched. The intended version of Fig. 3 is shown below, and replaces that of the original article:




Fig. 3 Room temperature angular-dependent CW EPR X-band spectra of (a) *m*-[VO(TrPP)]<sub>2</sub> and (b) *o*-[VO(TrPP)]<sub>2</sub> for crystal rotations around three orthogonal axes. For both panels, the upper row shows the 2D experimental EPR contour plots for the three rotations, acquired with a 3° step; the lower row shows representative EPR spectra (black lines) for the three rotations – from 0° to 180° every 15° – together with the best spectral simulations (red lines) obtained by using  $|J| = 0.01$  (0.005) cm<sup>-1</sup> and  $|J| = 0.05$  (0.01) cm<sup>-1</sup> for (a) and (b), respectively. Experimental frequency: 9.40 GHz for (a), 9.87 GHz for (b).

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

<sup>a</sup>Department of Chemistry "Ugo Schiff" & INSTM RU, University of Florence, Via della Lastruccia 3, 50019 Sesto Fiorentino, Italy. E-mail: lorenzo.sorace@unifi.it

<sup>b</sup>Department of Chemistry, NIS, University of Turin, Via P. Giuria 7, I10125 Torino, Italy

