



Correction: Tyrosine bioconjugation with hypervalent iodine

Cite this: *Chem. Sci.*, 2023, 14, 393

Nina Declas,^a John R. J. Maynard,^b Laure Menin,^c Natalia Gasilova,^c Sebastian Götze,^d Jakob L. Sprague,^e Pierre Stallforth,^d Stefan Matile^b and Jerome Waser^{*a}

DOI: 10.1039/d2sc90252d

Correction for 'Tyrosine bioconjugation with hypervalent iodine' by Nina Declas *et al.*, *Chem. Sci.*, 2022, 13, 12808–12817, <https://doi.org/10.1039/D2SC04558C>.

rsc.li/chemical-science

The authors regret that due to a technical issue with the ChemDraw file, part of the reaction conditions in Scheme 2 on page 12811 of the original article were not visible. A revised Scheme 2 with full reaction conditions is shown below:




^aLaboratory of Catalysis and Organic Synthesis, Institut des Sciences et Ingénierie Chimique, Ecole Polytechnique Fédérale de Lausanne, CH-1015, Lausanne, Switzerland. E-mail: jerome.waser@epfl.ch

^bDepartment of Organic Chemistry, University of Geneva, 1211 Geneva, Switzerland. E-mail: stefan.matile@unige.ch

^cInstitut des Sciences et Ingénierie Chimique, Ecole Polytechnique Fédérale de Lausanne, EPFL, 1015 Lausanne, Switzerland

^dDepartment of Paleobiotechnology, Leibniz Institute for Natural Product Research and Infection Biology, Hans Knöll Institute (HKI), 07745 Jena, Germany

^eDepartment of Microbial Pathogenicity Mechanisms, Leibniz Institute for Natural Product Research and Infection Biology, Hans Knöll Institute (HKI), 07745 Jena, Germany



Scheme 2 (a) Scope of tetramers on 20.0 μmol . (b) Scope of bioactive peptides on 1.0 μmol or 1.0 mg. Tyr* = Tyr modified with EBX. HPLC-MS yields are given, determined as indicated in Scheme 1. ^aIsolated yield. ^bObtained as a 1 : 2 mixture with 4.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

