

Cite this: *Chem. Sci.*, 2023, 14, 4351

All publication charges for this article have been paid for by the Royal Society of Chemistry

# Copper-catalyzed asymmetric allylic alkylation of racemic inert cyclic allylic ethers under batch and flow conditions†

Jun Li,<sup>a</sup> Xiao Song,<sup>a</sup> Yan Wang,<sup>a</sup> Junrong Huang,<sup>ab</sup> Hengzhi You<sup>ab</sup> and Fen-Er Chen<sup>abc</sup>

Highly enantioselective Cu-catalyzed asymmetric allylic alkylation of racemic inert cyclic allylic ethers is accomplished in this work. The use of Grignard reagents in combination with  $\text{BF}_3 \cdot \text{OEt}_2$  and  $\text{CuBr} \cdot \text{SMe}_2 / \text{L}_2$  is the key to enable employment of long-challenging low reactive allylic substrates in this AAA reaction. In addition, the approach exhibited a remarkable superiority in the construction of a stereogenic center involving challenging sterically hindered nucleophiles for both primary and secondary alkyl groups. Extension of this method under continuous flow conditions was also performed with excellent enantioselectivity in a short residence time. Mechanistic experiments indicated that the reaction proceeded through an *in situ* generated allylic bromide intermediate. All these advantages demonstrated that this chiral catalytic system is a valuable complement for existing asymmetric catalytic methods.

Received 9th January 2023  
Accepted 22nd March 2023

DOI: 10.1039/d3sc00127j

rsc.li/chemical-science

## Introduction

Asymmetric allylic alkylation (AAA) reactions are powerful synthetic tools to strategically install C–C bonds.<sup>1–4</sup> Generally, there are two main strategies for generating asymmetric C–C bonds in AAA reactions: (1) introduction of a chiral center from a prochiral substrate through the  $\text{S}_{\text{N}}2'$  pathway<sup>5–15</sup> (Scheme 1a); (2) resolution,<sup>16,17</sup> dynamic kinetic processes<sup>18–22</sup> (DyKAT), or direct enantioconvergent transformation<sup>23–27</sup> (DECT) from racemic chiral starting materials. Compared to prochiral substrates, AAA reactions from racemic substrates were far less studied. There are two equal amounts of enantiomers in the starting materials which complicated the catalytic processes and made the reaction more challenging.<sup>21</sup> Even though the use of stabilized nucleophiles ( $\text{p}K_{\text{a}} < 25$ ) in Pd-catalyzed AAA of racemic starting materials has been particularly successful since 1994,<sup>22,28</sup> the use of non-stabilized nucleophiles ( $\text{p}K_{\text{a}} > 50$ ) in Cu-catalyzed AAA of racemic starting materials with high yields and enantioselectivities was not reported until 2009.<sup>24</sup> Many pioneers have made significant contributions to the development of this chemistry since then. For example,

different copper–ligand complexes were synthesized,<sup>20,23</sup> and various non-stabilized nucleophiles such as Grignard reagents,<sup>24,29</sup> organolithium reagents,<sup>26</sup> and organozirconium reagents<sup>17,30</sup> were developed; versatile allylic substrates, such as bromides,<sup>20,23,24</sup> chlorides,<sup>17,18,29</sup> acetates<sup>25</sup> and phosphates<sup>19</sup> were employed (Scheme 1b). However, there still remain long-challenging issues in this field. First, Cu-catalyzed AAA reactions normally use allylic substrates with good leaving groups (such as Br, Cl, OCOR *etc.*) to enhance the reactivity and enantioselectivity.<sup>24</sup> However, these substrates are mostly sensitive, prone to deterioration, and hard to purify and store, which limits their applications in multistep reactions and synthesis.<sup>8,31</sup> Meanwhile, racemic inert substrates such as allylic ethers<sup>31–34</sup> have shown unsatisfactory performance in AAA reactions. The pioneering research reported by Consiglio and co-workers revealed moderate enantioselectivities in Ni-catalyzed AAA using phenyl-ethers.<sup>35</sup> Later on, Okamoto and co-workers achieved moderate results by employing pyridyl-ethers in Cu-catalyzed AAA with Grignard reagents.<sup>36</sup> In 2012, Feringa and co-workers developed a highly enantioselective Cu-catalyzed AAA reaction with prochiral linear allylic ethers using organolithium reagents.<sup>8</sup> However, the nucleophile scope was limited to  $n\text{BuLi}$  and  $n\text{HexLi}$  in this beautiful chemistry. To the best of our knowledge, despite extensive endeavours, highly efficient Cu-catalyzed AAA reactions with racemic inert substrates have not yet emerged.

Another key challenge in Cu-catalyzed AAA reactions is the generality of the non-stabilized nucleophile scope used in the catalytic systems. Employment of functionalized and sterically hindered nucleophiles was found to be extremely difficult


<sup>a</sup>School of Science, Harbin Institute of Technology (Shenzhen), Taoyuan Street, Nanshan District, Shenzhen, 518055, China. E-mail: [hjunrong@hit.edu.cn](mailto:hjunrong@hit.edu.cn); [youthengzhi@hit.edu.cn](mailto:youthengzhi@hit.edu.cn); [rfchen@fudan.edu.cn](mailto:rfchen@fudan.edu.cn)

<sup>b</sup>Green Pharmaceutical Engineering Research Center, Harbin Institute of Technology (Shenzhen), Taoyuan Street, Nanshan District, Shenzhen, 518055, China

<sup>c</sup>Engineering Center of Catalysis and Synthesis for Chiral Molecules, Department of Chemistry, Fudan University, Shanghai, 200433, China

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d3sc00127j>





Scheme 1 Asymmetric allylic alkylation (AAA) procedures.

during the process.<sup>24</sup> Crucial pioneering work was first conducted by Alexakis and co-workers where sterically hindered secondary alkyl groups were accomplished in moderate enantioselectivities with racemic allylic bromides.<sup>23</sup> Later on, the improvement of the performance of secondary alkyl metal reagents was achieved with the same group by a well-designed catalytic system using racemic allylic trifluoroacetates, though it was not compatible with low reactive simple primary alkyl groups (such as MeMgBr).<sup>25</sup> At the same time, Fletcher and co-workers described the breakthrough of a Cu-catalyzed AAA of racemic allylic chlorides with organozirconium reagents. The reaction was able to tolerate a variety of functionalized primary alkyl groups under convenient mild conditions.<sup>18,19</sup> To the best of our knowledge, despite all these excellent advances, a general universal catalytic system which could enable efficient Cu-catalyzed AAA with challenging sterically hindered secondary and simple primary alkyl nucleophiles has not emerged. Herein, we reported a highly efficient Cu-catalyzed AAA reaction using racemic inert cyclic allylic ethers in combination with BF<sub>3</sub>·OEt<sub>2</sub> and CuBr·SMe<sub>2</sub>/L2. Noteworthy, this catalytic system delivered general efficiency for challenging sterically hindered Grignard reagents with both primary and secondary alkyl groups.

## Results and discussion

We started our investigation by examining the AAA reactions of 3-benzoyloxycyclohexene (**1a**) with 1.5 equivalents of MeLi as a nucleophile in the presence of BF<sub>3</sub>·OEt<sub>2</sub> (2 eq.), catalyzed by CuBr·SMe<sub>2</sub> (5 mol%) and phosphoramidites (*S,S,S*)-L2 (11 mol%) in DCM. The reaction gave 3-methylcyclohexene (**3a**) in 70% yield with 48% ee (Table 1, entry 1). To enhance the enantioselectivity of **3a**, we screened different copper sources,


ligands and Lewis acids. However, employment of copper sources (such as CuCl, CuTc and Cu(OTf)<sub>2</sub>), ligands (such as (*S,S,S*)-L1 and (*S,S,R*)-L3), and Lewis acids (BF<sub>2</sub>OTf generated *in situ* from BF<sub>3</sub>·OEt<sub>2</sub> and TMSOTf) gave poor results (Table 1, entries 1–7). Encouragingly, the enantioselectivity of **3a** improved to 87% ee by using less reactive MeMgBr (Table 1, entry 8). Meanwhile, the copper/ligand ratio was found to have little effect on this catalytic system (Table 1, entries 8 vs. 9). When diluted MeMgBr (0.5 M in Et<sub>2</sub>O) was used, the enantioselectivity increased further to 96% ee in 46% yield (Table 1, entry 10). Delightfully, the yield of **3a** was also optimized to 85% maintaining the high enantioselectivity by adjusting –OBn to the less hindered –OMe group (Table 1, entry 11). Benefiting from this low activity of the ether substrate, MeMgBr could be added dropwise for only 1 min instead of the 0.5 h or 2 h reported in previous work<sup>8,25</sup> (Table 1, entries 11 vs. 12). Finally, single isomer **3a** was produced in excellent yield and enantioselectivity with 10 mol% catalyst loading (99% yield, 98% ee, Table 1, entry 13). It was worth noting that the enantioselectivity of the product (Table S2,† entry 3, 94% ee) remained excellent even at a low catalyst loading (1 mol%), but the conversion rate of 3-methoxycyclohex-1-ene (**1b**) showed a significant decline at 1 mol% catalyst loading (Table S2,† entry 3, 78% conversion rate), despite extending the reaction time to 4 hours.

Table 1 Selected optimization experiments<sup>a</sup>

Entry	Nu	R	CuX	L	Yield <sup>d</sup>	ee <sup>f</sup>
1	MeLi	Bn	CuBr·SMe <sub>2</sub>	L2	70%	48%
2	MeLi	Bn	CuCl	L2	57%	24%
3	MeLi	Bn	CuTc	L2	61%	37%
4	MeLi	Bn	CuBr·SMe <sub>2</sub>	L1	74%	<10%
5	MeLi	Bn	CuBr·SMe <sub>2</sub>	L3	54%	15%
6	MeLi	Bn	Cu(OTf) <sub>2</sub>	L3	N.D.	—
7 <sup>c</sup>	MeLi	Bn	CuBr·SMe <sub>2</sub>	L2	N.D.	—
8 <sup>b,f</sup>	MeMgBr	Bn	CuBr·SMe <sub>2</sub>	L2	48%	87%
9 <sup>b</sup>	MeMgBr	Bn	CuBr·SMe <sub>2</sub>	L2	45%	87%
10 <sup>b</sup>	MeMgBr <sup>e,h</sup>	Bn	CuBr·SMe <sub>2</sub>	L2	46%	96%
11 <sup>b</sup>	MeMgBr <sup>e,h</sup>	Me	CuBr·SMe <sub>2</sub>	L2	85%	96%
12 <sup>b</sup>	MeMgBr <sup>h</sup>	Me	CuBr·SMe <sub>2</sub>	L2	88%	96%
13 <sup>b,g</sup>	MeMgBr <sup>h</sup>	Me	CuBr·SMe <sub>2</sub>	L2	99%	98%

<sup>a</sup> Conditions: racemic **1** (0.4 mmol), CuX (5% mol), L (11% mol), BF<sub>3</sub>·OEt<sub>2</sub> (2 eq.), CH<sub>2</sub>Cl<sub>2</sub> (4 mL), –78 °C, MeLi (1.5 eq.), 2 h addition time. <sup>b</sup> Conditions: racemic **1** (0.4 mmol), CuX (5% mol), L (6% mol), BF<sub>3</sub>·OEt<sub>2</sub> (1.5 eq.), CH<sub>2</sub>Cl<sub>2</sub> (4 mL), –78 °C, MeMgBr (3 M in Et<sub>2</sub>O, 1.5 eq.), 1 min addition time. <sup>c</sup> 2 eq. BF<sub>3</sub>·OEt<sub>2</sub> and 6 eq. TMSOTf were used. <sup>d</sup> Yield was determined by GC. <sup>e</sup> 2 h addition time. <sup>f</sup> CuX (5% mol) and L (11% mol) were used. <sup>g</sup> CuX (10% mol) and L (11% mol) were used. <sup>h</sup> MeMgBr (0.5 M in Et<sub>2</sub>O) was used. <sup>i</sup> ee was determined by chiral GC.






Scheme 2 Substrate scope. <sup>a</sup>Conditions: racemic **1** (0.4 mmol),  $\text{CuBr}\cdot\text{SMe}_2$  (10% mol), **L2** (11% mol),  $\text{BF}_3\cdot\text{OEt}_2$  (1.5 eq.),  $\text{CH}_2\text{Cl}_2$  (4 mL),  $-78\text{ }^\circ\text{C}$ ,  $\text{MeMgBr}$  (1.5 eq.). ee was determined by chiral GC. <sup>b</sup>Yield was determined by GC or NMR. <sup>c</sup>The highest ee reported for the alkylation of analogous allylic bromides is given in parenthesis. <sup>d</sup>**L4** was used. <sup>e</sup>Isolated yield of the corresponding diastereoisomeric epoxides.

With the optimal conditions in hand, the nucleophile scope of this novel protocol was then evaluated. Several enantiomerically enriched cycloalkenes were obtained with excellent enantioselectivities by using simple unfunctionalized primary alkyl (**3a–3f**, **4a**), alkenyl (**3g**) and aromatic Grignard reagents (**3i–3k**, **5a**). Surprisingly, bulky alkyl groups such as secondary alkyl Grignard reagents also reacted smoothly to give the corresponding products **3l–3p** in 83–93% yields with excellent enantioselectivities. In particular, decent improvements were achieved for previously challenging sterically hindered nucleophiles in AAA of the reported allylic bromide approaches,<sup>23</sup> such as **3h** (65% ee vs. 28% ee), **3l** (82% ee vs. 40% ee), **3m** (90% ee vs. 74% ee), and **3n** (88% ee vs. 70% ee) and for different ring sizes **4a** (91% ee vs. 60% ee) and **5a** (83% ee vs. 44% ee). To further extend the generality of this catalytic system, the reaction conditions were applied to challenging acyclic ether. However, only 15% ee was obtained in the reaction (Scheme S1<sup>†</sup>), which was similar to the previous result with acyclic bromide (8% ee).<sup>23</sup> Preliminary experiments of AAA reactions using challenging 3-methoxy-1-methylcyclohex-1-ene to form a more complex quaternary cycloalkene were also performed. A mixture of 1,1- and 1,3-disubstituted alkylation products was obtained with moderate enantioselectivities (Scheme S2, <sup>†</sup> 35% yield, 83% ee for


1,1-disubstituted quaternary cycloalkene and 12% yield, 53% ee for 1,3-disubstituted cycloalkene). Further research focused on the construction of a stereogenic quaternary center is currently under study and will be reported in due course.

Although satisfactory results were achieved under batch conditions, an intractable question worth thinking about is whether Grignard reagents are too reactive to scale up.<sup>37–39</sup> The recent emerging continuous flow technology demonstrated many advantages, including higher safety and efficiency, accurate control, better heat and mass transfer, easier amplification and better sustainability.<sup>40–43</sup> Thus, we also explored the application of this protocol under continuous flow conditions. However, the enantioselectivities of the products dropped significantly under continuous flow conditions when using allylic bromide as the substrate in previous reports (up to 33% ee).<sup>26</sup> Generally, the organometallic reagents were added dropwise over 2 h (ref. 26) or 0.5 h (ref. 25) to a solution of substrates to avoid side-reactions under batch conditions. In contrast, substrates and organometallic reagents are mixed simultaneously under flow conditions; the highly active allylic bromides may undergo uncatalyzed reactions<sup>6</sup> or react with “ate” and “aggregated” copper species<sup>25</sup> generated by a chiral catalyst exposed to an excess of organometallic reagents during the process. We suspected that the difference in contact mode between batch and flow mainly resulted in a decrease in enantioselectivities. Thus, we started our investigation by employing inert allylic ethers instead of high reactive allylic substrates here. The flow system was set up as shown in Scheme 3. Solutions of  $\text{CuBr}\cdot\text{SMe}_2$ , ligand **L2** and cyclic racemic substrate **1b** were first mixed with a  $\text{BF}_3\cdot\text{OEt}_2$  stream, and then, another stream of phenylethyl Grignard reagent was pumped into a microchip reactor. A



Scheme 3 Cu-catalyzed AAA reaction of **1b** under continuous flow conditions.





Scheme 4 Mechanistic analysis and a plausible reaction mechanism.

sonication device was employed to increase reaction mixing and prevent solid aggregation. Encouragingly, the desired products were obtained in high yield with excellent enantioselectivity within 3.6 s under optimized flow conditions (Table S1 and Scheme S3<sup>†</sup>). And the system could continue operating for over 700 min without any loss of activity or enantioselectivity to provide 1.09 g product (84% yield, 91% ee).

We then paid our attention to the mechanism of this new Cu-catalyzed AAA protocol. A Lewis acid was confirmed to be the key for the formation of desired products from racemic inert cyclic allylic ethers here.<sup>44</sup> We carried out the AAA reaction of **1b** without  $\text{BF}_3 \cdot \text{OEt}_2$ , and the starting material **1b** could be almost completely recovered (Scheme 4 – (1)). It suggested that ether groups could act as a robust protecting group for former stages of synthetic schemes in the absence of a Lewis acid. Conversely, the ether groups could become leaving groups to perform enantioselective C–C bond formation in the presence of a Lewis acid and copper catalyst. This provided new opportunities for the application of the Cu-catalyzed AAA protocol using racemic inert cyclic allylic ethers in multistep reactions and synthesis.<sup>31</sup> In addition, we examined the AAA reaction of **1b** using cyclohexylmagnesium chloride as a nucleophile instead of cyclohexylmagnesium bromide (88% ee, Scheme 2, **3n**). Surprisingly, the enantioselectivity of product **3n** using cyclohexylmagnesium chloride was very low (Scheme 4 – (2), 33% ee). This phenomenon indicated that halogen anions from Grignard reagents might be involved in the reaction and influence the enantioselectivity of the product. A similar phenomenon was also observed in Fletcher and co-workers's work.<sup>18,19</sup> They considered that allyl halide formed *in situ* from a halogen anion and phosphate in the reaction and reacted with nucleophiles to prove enantio-enriched alkylation products. We suspected that the allylic ethers might undergo analogical transformation here. To gain more insight into the reaction mechanism,  $\text{MgBr}_2$ , instead of Grignard reagents was added to the mixture of **1b** and  $\text{BF}_3 \cdot \text{OEt}_2$ . As we expected, 45% cyclic allylic bromide (**1c**) was observed in 1 h (Fig. S3<sup>†</sup>), indicating that **1c** could be generated *in situ* from **1b** in this catalytic system (Scheme 4 – (2) and (3)). Furthermore, since we were not able to observe the intermediate **1c** in the reaction of racemic allylic ethers, an even lower reactive **1a** substrate was employed in our

experiments. Inspiringly, trace **1c** was observed during our standard AAA reaction (Scheme 4 – (4), Fig. S1 and S2<sup>†</sup>). In combination with the inspiration from Alexakis's and co-workers pioneering work for DECT progress,<sup>25</sup> we proposed a plausible mechanistic route for this new Cu-catalyzed AAA protocol using racemic inert cyclic allylic ethers. First, racemic **1c** is generated from **1b** in the presence  $\text{BF}_3 \cdot \text{OEt}_2$  and  $\text{Br}^-$ . Then, (*R*)-**1c** and (*S*)-**1c** underwent a DECT process to form the same intermediate **Int-1**. Finally, **Int-1c** underwent reductive elimination to provide the desired single product **3** in high yield and enantioselectivity (Scheme 4 – (5)).

## Conclusions

In summary, a highly efficient Cu-catalyzed AAA reaction of racemic inert cyclic allylic ethers was developed here. The use of  $\text{BF}_3 \cdot \text{OEt}_2$  with  $\text{CuBr} \cdot \text{SMe}_2/\text{L2}$  was confirmed to be crucial in this catalytic system through the proposed allylic bromide intermediate. Meanwhile, the new method showed significant advantages in nucleophile scope over existing protocols with racemic sensitive allylic substrates. In addition, application of the method under continuous flow conditions was also conducted successfully with excellent enantioselectivity and yield in a short residence time. We anticipate that this research will aid the development of a new AAA reaction with inert substrates and provide a valuable tool for existing asymmetric catalytic methods. Intensive efforts are now directed toward the extension of this process to other inert allylic substrates in our lab.

## Data availability

The ESI<sup>†</sup> is available and contains experimental and analytical data.

## Author contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.



## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

This work was supported by the Shenzhen Science and Technology Research Fund No. JCYJ20190806142203709; JSGG20191129114029286; JSGG20201103153807021; GXWD20220811173736002. We are also grateful to the Guangdong Basic and Applied Basic Research Foundation (No. 2021A1515110366) and Shenzhen Key Laboratory of Advanced Functional Carbon Materials Research and Comprehensive Application. We appreciate the guidance and opinions of Jingyuan Liao and Junwen Wang.

## Notes and references

- C. A. Falciola and A. Alexakis, Copper-Catalyzed Asymmetric Allylic Alkylation, *Eur. J. Org. Chem.*, 2008, **2008**, 3765–3780.
- L. Susse and B. M. Stoltz, Enantioselective Formation of Quaternary Centers by Allylic Alkylation with First-Row Transition-Metal Catalysts, *Chem. Rev.*, 2021, **121**, 4084–4099.
- T. Fan, J. Song and L. Z. Gong, Asymmetric Redox Allylic Alkylation to Access 3,3'-Disubstituted Oxindoles Enabled by Ni/NHC Cooperative Catalysis, *Angew. Chem., Int. Ed.*, 2022, **61**, e202201678.
- X. J. Liu, W. Y. Zhang, C. Zheng and S. L. You, Iridium-Catalyzed Asymmetric Allylic Substitution of Methyl Azaarenes, *Angew. Chem., Int. Ed.*, 2022, **61**, e202200164.
- K. Tissot-Croset, D. Polet and A. Alexakis, A highly effective phosphoramidite ligand for asymmetric allylic substitution, *Angew. Chem., Int. Ed.*, 2004, **43**, 2426–2428.
- M. Perez, M. Fananas-Mastral, P. H. Bos, A. Rudolph, S. R. Harutyunyan and B. L. Feringa, Catalytic asymmetric carbon-carbon bond formation via allylic alkylations with organolithium compounds, *Nat. Chem.*, 2011, **3**, 377–381.
- M. Fananas-Mastral, M. Perez, P. H. Bos, A. Rudolph, S. R. Harutyunyan and B. L. Feringa, Enantioselective synthesis of tertiary and quaternary stereogenic centers: copper/phosphoramidite-catalyzed allylic alkylation with organolithium reagents, *Angew. Chem., Int. Ed.*, 2012, **51**, 1922–1925.
- M. Perez, M. Fananas-Mastral, V. Hornillos, A. Rudolph, P. H. Bos, S. R. Harutyunyan and B. L. Feringa, Asymmetric allylic alkylation of acyclic allylic ethers with organolithium reagents, *Chem.–Eur. J.*, 2012, **18**, 11880–11883.
- V. Hornillos, M. Perez, M. Fananas-Mastral and B. L. Feringa, Copper-catalyzed enantioselective allyl-allyl cross-coupling, *J. Am. Chem. Soc.*, 2013, **135**, 2140–2143.
- W.-H. Deng, F. Ye, X.-F. Bai, L. Li, T. Song, Y.-L. Wei and L.-W. Xu, Chlorotrimethylsilane (TMSCl): an efficient silicon-based Lewis acid mediator in allylic alkylation using a diethylzinc reagent, *RSC Adv.*, 2014, **4**, 479–483.
- M. Fananas-Mastral, R. Vitale, M. Perez and B. L. Feringa, Enantioselective synthesis of all-carbon quaternary stereogenic centers via copper-catalyzed asymmetric allylic alkylation of (*Z*)-allyl bromides with organolithium reagents, *Chem.–Eur. J.*, 2015, **21**, 4209–4212.
- S. Guduguntla, V. Hornillos, R. Tessier, M. Fananas-Mastral and B. L. Feringa, Chiral Diarylmethanes via Copper-Catalyzed Asymmetric Allylic Arylation with Organolithium Compounds, *Org. Lett.*, 2016, **18**, 252–255.
- W. Xiong, G. Xu, X. Yu and W. Tang, P-Chiral Monophosphorus Ligands for Asymmetric Copper-Catalyzed Allylic Alkylation, *Organometallics*, 2019, **38**, 4003–4013.
- R. Tarrieu, A. Dumas, J. Thongpaen, T. Vives, T. Roisnel, V. Dorcet, C. Crévisy, O. Baslé and M. Mauduit, Readily Accessible Unsymmetrical Unsaturated 2,6-Diisopropylphenyl N-Heterocyclic Carbene Ligands. Applications in Enantioselective Catalysis, *J. Org. Chem.*, 2017, **82**, 1880–1887.
- S. Guduguntla, J.-B. Gualtierotti, S. S. Goh and B. L. Feringa, Enantioselective Synthesis of Di- and Tri-Arylated All-Carbon Quaternary Stereocenters via Copper-Catalyzed Allylic Arylations with Organolithium Compounds, *ACS Catal.*, 2016, **6**, 6591–6595.
- A. Alexakis and R. Millet, SimplePhos as Efficient Ligand for the Copper-Catalyzed Kinetic Resolution of Cyclic Vinyloxiranes with Grignard Reagents, *Synlett*, 2008, **2008**, 1797–1800.
- S. Karabiyikoglu, A. V. Brethomé, T. Palacin, R. S. Paton and S. P. Fletcher, Enantiomerically enriched tetrahydropyridine allyl chlorides, *Chem. Sci.*, 2020, **11**, 4125–4130.
- H. You, E. Rideau, M. Sidera and S. P. Fletcher, Non-stabilized nucleophiles in Cu-catalysed dynamic kinetic asymmetric allylic alkylation, *Nature*, 2015, **517**, 351–355.
- E. Rideau, H. You, M. Sidera, T. D. W. Claridge and S. P. Fletcher, Mechanistic Studies on a Cu-Catalyzed Asymmetric Allylic Alkylation with Cyclic Racemic Starting Materials, *J. Am. Chem. Soc.*, 2017, **139**, 5614–5624.
- X. Y. Cui, Y. Ge, S. M. Tan, H. Jiang, D. Tan, Y. Lu, R. Lee and C. H. Tan, (Guanidine)copper Complex-Catalyzed Enantioselective Dynamic Kinetic Allylic Alkynylation under Biphasic Condition, *J. Am. Chem. Soc.*, 2018, **140**, 8448–8455.
- R. He, X. Huo, L. Zhao, F. Wang, L. Jiang, J. Liao and W. Zhang, Stereodivergent Pd/Cu Catalysis for the Dynamic Kinetic Asymmetric Transformation of Racemic Unsymmetrical 1,3-Disubstituted Allyl Acetates, *J. Am. Chem. Soc.*, 2020, **142**, 8097–8103.
- V. Bhat, E. R. Welin, X. Guo and B. M. Stoltz, Advances in Stereoconvergent Catalysis from 2005 to 2015: Transition-Metal-Mediated Stereoablative Reactions, Dynamic Kinetic Resolutions, and Dynamic Kinetic Asymmetric Transformations, *Chem. Rev.*, 2017, **117**, 4528–4561.
- J.-B. Langlois and A. Alexakis, Copper-Catalyzed Asymmetric Allylic Alkylation of Racemic Cyclic Substrates: Application of Dynamic Kinetic Asymmetric Transformation (DYKAT), *Adv. Synth. Catal.*, 2010, **352**, 447–457.



- 24 J. B. Langlois and A. Alexakis, Dynamic kinetic asymmetric transformation in copper catalyzed allylic alkylation, *Chem. Commun.*, 2009, 3868–3870, DOI: [10.1039/b907722g](https://doi.org/10.1039/b907722g).
- 25 J.-B. Langlois, D. Emery, J. Mareda and A. Alexakis, Mechanistic identification and improvement of a direct enantioconvergent transformation in copper-catalyzed asymmetric allylic alkylation, *Chem. Sci.*, 2012, **3**, 1062–1069.
- 26 J. Li, X. Song, F. Wu, H. You and F.-E. Chen, Cu-Catalyzed Asymmetric Allylic Alkylation of Racemic Cyclic Allyl Bromides with Organolithium Compounds, *Eur. J. Org. Chem.*, 2022, **2022**, e202200860.
- 27 Y. Ge, X. Y. Cui, S. M. Tan, H. Jiang, J. Ren, N. Lee, R. Lee and C. H. Tan, Guanidine-Copper Complex Catalyzed Allylic Borylation for the Enantioconvergent Synthesis of Tertiary Cyclic Allylboronates, *Angew. Chem., Int. Ed.*, 2019, **58**, 2382–2386.
- 28 B. M. Trost and R. C. Bunt, Asymmetric induction in allylic alkylations of 3-(acyloxy)cycloalkenes, *J. Am. Chem. Soc.*, 1994, **116**, 4089–4090.
- 29 J. B. Langlois and A. Alexakis, Identification of a valuable kinetic process in copper-catalyzed asymmetric allylic alkylation, *Angew. Chem., Int. Ed.*, 2011, **50**, 1877–1881.
- 30 M. Sidera and S. P. Fletcher, Cu-catalyzed asymmetric addition of sp<sup>2</sup>-hybridized zirconium nucleophiles to racemic allyl bromides, *Chem. Commun.*, 2015, **51**, 5044–5047.
- 31 N. A. Butt and W. Zhang, Transition metal-catalyzed allylic substitution reactions with unactivated allylic substrates, *Chem. Soc. Rev.*, 2015, **44**, 7929–7967.
- 32 N. Suzuki, D. Y. Kondakov and T. Takahashi, Zirconium-catalyzed highly regioselective carbon-carbon bond formation reactions, *J. Am. Chem. Soc.*, 1993, **115**, 8485–8486.
- 33 H. Yasui, K. Mizutani, H. Yorimitsu and K. Oshima, Cobalt- and rhodium-catalyzed cross-coupling reaction of allylic ethers and halides with organometallic reagents, *Tetrahedron*, 2006, **62**, 1410–1415.
- 34 H. F. Tu, C. Zheng, R. Q. Xu, X. J. Liu and S. L. You, Iridium-Catalyzed Intermolecular Asymmetric Dearomatization of beta-Naphthols with Allyl Alcohols or Allyl Ethers, *Angew. Chem., Int. Ed.*, 2017, **56**, 3237–3241.
- 35 G. Consiglio, O. Piccolo, L. Roncetti and F. Morandini, Asymmetric nickel-catalyzed cross-coupling reaction of allylic substrates with grignard reagents, *Tetrahedron*, 1986, **42**, 2043–2053.
- 36 S. Tominaga, Y. Oi, T. Kato, D. K. An and S. Okamoto,  $\gamma$ -Selective allylic substitution reaction with Grignard reagents catalyzed by copper N-heterocyclic carbene complexes and its application to enantioselective synthesis, *Tetrahedron Lett.*, 2004, **45**, 5585–5588.
- 37 G. Laudadio, Y. Deng, K. van der Wal, D. Ravelli, M. Nuño, M. Fagnoni, D. Guthrie, Y. Sun and T. Noël, C(sp<sup>3</sup>)-H functionalizations of light hydrocarbons using decatungstate photocatalysis in flow, *Science*, 2020, **369**, 92–96.
- 38 Y. Saito and S. Kobayashi, Chiral Heterogeneous Scandium Lewis Acid Catalysts for Continuous-Flow Enantioselective Friedel-Crafts Carbon-Carbon Bond-Forming Reactions, *Angew. Chem., Int. Ed.*, 2021, **60**, 26566–26570.
- 39 J. Wang, J. Li, Y. Wang, S. He, H. You and F.-E. Chen, Polymer-Supported Chiral Heterogeneous Copper Catalyst for Asymmetric Conjugate Addition of Ketones and Imines under Batch and Flow, *ACS Catal.*, 2022, **12**, 9629–9637.
- 40 J. Liao, S. Zhang, Z. Wang, X. Song, D. Zhang, R. Kumar, J. Jin, P. Ren, H. You and F.-E. Chen, Transition-metal catalyzed asymmetric reactions under continuous flow from 2015 to early 2020, *Green Synth. Catal.*, 2020, **1**, 121–133.
- 41 Y. Wu, W. Q. Chen, Y. Q. Zhao and H. R. Piao, Efficient synthesis of panaxadiol derivatives using continuous-flow microreactor and evaluation of anti-tumor activity, *Chin. Chem. Lett.*, 2015, **26**, 334–338.
- 42 Y. Xin, S. Peng, J. X. Chen, Z. J. Yang and J. Y. Zhang, Continuous flow synthesis of porous materials, *Chin. Chem. Lett.*, 2020, **31**, 1448–1461.
- 43 L. Wan, G. Kong, M. Liu, M. Jiang, D. Cheng and F. Chen, Flow chemistry in the multi-step synthesis of natural products, *Green Synth. Catal.*, 2022, **3**, 243–258.
- 44 Y. Yamamoto, S. Yamamoto, H. Yatagai and K. Maruyama, Lewis acid mediated reactions of organocopper reagents. A remarkably enhanced regioselective gamma-attack of allylic halides and direct alkylation of allylic alcohols via RCu·BF<sub>3</sub> system, *J. Am. Chem. Soc.*, 1980, **102**, 2318–2325.

