# Chemical Science



# **EDGE ARTICLE**

View Article Online
View Journal | View Issue



Cite this: Chem. Sci., 2023, 14, 6970

dll publication charges for this article have been paid for by the Royal Society of Chemistry

Received 13th March 2023 Accepted 5th June 2023

DOI: 10.1039/d3sc01353g

rsc li/chemical-science

# Sequencing palladium-catalyzed cycloisomerization cascades in a synthesis of the gelsemine core†

Guoduan Liang and Edward A. Anderson 🕩 \*

Transition metal-catalyzed cycloisomerization is a powerful strategy for the construction of cyclic organic molecules, and the use of palladium catalysts can deliver a wide range of monocyclic and bicyclic products. However, applications of cycloisomerizations in complex target synthesis in which more than one cycloisomerization process is deployed in a cascade context are rare. Here we report investigations of the relative rates of two different types of ene-ynamide cycloisomerization that form fused and spirocyclic rings, and use of these results to design a sequence-controlled cascade cycloisomerization that prepares the tetracyclic core of gelsemine in a single step. Crucial to this work was an evaluation of the kinetics of each cycloisomerization in competition experiments, which revealed a key influence of the ynamide electron-withdrawing group on the cycloisomerization reaction.

#### Introduction

Transition metal-catalyzed cycloisomerizations have been widely explored as atom-economic processes to form cyclic organic molecules from acyclic polyunsaturated starting materials.1-4 Among many metals, palladium catalysts are wellestablished as versatile promoters of such transformations.5-12 When substrates feature multiple unsaturated bonds, elegant cascade processes have been developed whereby intermediate alkyl- or alkenylpalladium species undergo multiple C-C bond forming events in a single step;13 one pioneering example from the Trost group is shown in Scheme 1a  $(1 \rightarrow 2)$ .<sup>14</sup> Pd-catalyzed cycloisomerizations have also been widely used in total synthesis, 1,15-22 however applications of cascade cycloisomerizations are rare in this context. Sequential cycloisomerizations have been reported, such as an elegant approach to dendrobine described by the Chen group in which palladium- and rhodium-catalyzed cycloisomerizations were implemented in consecutive steps to construct the bicyclic pyrrolidine 3 from dienyne 4.23 A palladium-catalyzed cycloisomerization cascade was described by Trost et al. in which dienyne 5 underwent cyclization to triene 6 in the synthesis of various terpenoid natural products.24 However, to our knowledge, no examples of cascade cycloisomerizations have been reported in which a transition metal catalyst operates twice on

Scheme 1 (a) Cascade or sequenced cycloisomerizations. (b) Previous work from our group on palladium-catalyzed ynamide cycloisomerization. (c) This work: time-resolved palladium-catalyzed cascade cycloisomerization towards the synthesis of the gelsemine core.

a Trost et al. (Ref. 14) cat. Pd(dba)•CHCl<sub>3</sub> e / PPh<sub>3</sub>, AcOH PhO<sub>2</sub>S PhO<sub>2</sub>S PhO<sub>2</sub>S cascade Lee et al. (Ref. 23) cat. Pd(OAc)<sub>2</sub> / PPh<sub>3</sub> cat. RhCl(PPh<sub>3</sub>)<sub>3</sub> / AgSbF<sub>6</sub> sequential cvcloisomerizations Trost et al. (Ref. 24) cat. Pd(OAc)<sub>2</sub> single step 5 cascade cat. Pd(OAc)<sub>2</sub> / cat. Pd(OAc)<sub>2</sub> TsŅ bbeda bbeda OPG double-ΡĠ step cascade 7: (+)-Gelsemine

Chemistry Research Laboratory, Department of Chemistry, University of Oxford, 12 Mansfield Road, Oxford OX1 3TA, UK. E-mail: edward.anderson@chem.ox.ac.uk

 $<sup>\</sup>dagger$  Electronic supplementary information (ESI) available: Experimental procedures, details of reaction optimization, copies of  $^{\rm I}{\rm H}$  and  $^{\rm 13}{\rm C}$  NMR spectra (pdf). See DOI: https://doi.org/10.1039/d3sc01353g

**Edge Article** 

reaction success.


a reaction substrate, where the order of events is critical for

Our group has developed a variety of metal-catalyzed cyclo-isomerizations of alkenyl ynamides (*e.g.*, Scheme 1b).<sup>25–27</sup> These compounds are readily accessed by alkynylation of appropriate carbamate and sulfonamide precursors, <sup>28–36</sup> and undergo Pd-catalyzed cyclizations to azacycles under mild reaction conditions. We further studied the mechanism of these cyclo-isomerizations, where a series of deuterium-labelling studies demonstrated that the so-called 'ligand' bis(benzylidene)ethylenediamine (bbeda) in fact also serves a source of a palladium(II) hydride species that initiates the catalysis.<sup>37</sup> Building from this work, we questioned whether we could develop a cyclo-isomerization cascade in which two discrete cyclo-isomerizations would be sequenced in a time-resolved manner, thus generating products of greater complexity with higher efficiency.

We were particularly attracted to the challenge of differentiating between fused-ring and spiro-ring formation (as shown in Scheme 1b), as we recognized that a sequence of these reactions could generate the core scaffold of the indole alkaloid gelsemine (7, Scheme 1c). This natural product features a hexacyclic core with seven contiguous stereocenters, including two quaternary carbons, whose intriguing structure has attracted the attention of many chemists.<sup>38-41</sup> We questioned whether the challenging tetracyclic spirooxindole core of gelsemine could be obtained using a one-pot cascade polycyclization sequence. Specifically, tetracycle 8 might derive from cyclization of bisynamide 9 by initial fused-ring formation (Step 1), and then spiro-ring formation (Step 2), 'walking' the double bond of the linking cyclohexene around the six-membered ring. To achieve this time-resolved process, an understanding of the relative rates of each process would be critical, as if Step 2 occurs before Step 1 then a totally different skeleton would be formed. We were aware that the relative rates of these processes would likely depend not only on the conformational demands of the cyclizations, and the relative rates of the elementary steps of the cycloisomerization, but also on the nature of the ynamide electron-withdrawing groups, which had not previously been studied. Here we describe the exploration of these factors, and the successful execution of this cascade cycloisomerization, which to our knowledge represents the first example of the sequencing of two independent ring-forming events in palladium-catalyzed cycloisomerization chemistry.

## Results and discussion

As the relative rate of spiro ring formation (Step 2 in Scheme 1c) compared to fused ring formation was expected to be crucial for the success of the cascade, we first synthesized two model 'spiro' enynamide substrates  ${\bf 10}$  and  ${\bf 11}$  (Scheme 2a and b respectively), differing in the nature of the ynamide electron-withdrawing group, which as noted above could make a significant difference to the rate of cyclization. The synthesis of N-aryl ynamides can be challenging, and we selected an approach based on dichloroenamide precursors developed previously by our group.  $^{42,43}$  Towards sulfonamide ynamide  ${\bf 10}$ , lithiation of



Scheme 2 Model spiro-ring synthesis (a), model fused-ring synthesis (b), and reaction kinetics.

bromide **12** <sup>44</sup> followed by addition to cyclohexanone gave tertiary alcohol **13** which, without purification, underwent dehydration to afford alkene **14** (63% over two steps). Treatment of **14** with trichloroethene and Cs<sub>2</sub>CO<sub>3</sub> gave dichloroenamide **15** (97%), which was converted to *N*-tosyl ynamide **10** on treatment with PhLi, then iodomethane (90%). Subjection of **10** to Pd(OAc)<sub>2</sub>/bbeda (10 mol%) at room temperature for 12 h gave the spiroindoline **16** in 80% yield as a single alkene diastereomer. Other catalyst systems tested (*e.g.* Pd(OAc)<sub>2</sub>/2PPh<sub>3</sub>, or Pd(OAc)<sub>2</sub> alone) also proceeded efficiently. An NMR reaction profile obtained at 45 °C revealed that the reaction reached completion after around 2 hours at that temperature.

Synthesis of the equivalent *N*-carbamate ynamide **11** was attempted using a similar route, but proved unsuccessful due to

formation of a cyclic carbamate in the first step (by cyclization of the tertiary alcohol onto the *N*-carbamate in the equivalent of intermediate 13).<sup>45</sup> Instead, Suzuki coupling of boronic ester 17 <sup>46</sup> and enol triflate 18 afforded aniline 19 in 73% yield, which was converted to dichloroenamide 20 in two steps (62%). Treatment of 20 with LiHMDS generated an intermediate chloroynamide 21, which underwent *in situ* copper-catalyzed cross-coupling with Me<sub>2</sub>Zn to give ynamide 11 in 48% yield. Pd(OAc)<sub>2</sub>/bbeda-catalyzed cyclization of 11 at room temperature afforded the spirocyclic product 22 in 70% yield, as a 2.5:1 (Z:E) mixture of diastereomers. Interestingly, the two cycloisomerizations (of 10 and 11) appeared to proceed at quite similar reaction rates; an NMR reaction profile obtained at 45 °C revealed the reaction of 11 reached completion after around 1 hour

We next targeted a model fused-ring system (23, Scheme 3). An N-tosyl electron-withdrawing group was selected due to the efficiency of ynamide formation for aliphatic amides with this group, compared to carbamates. This synthesis commenced with an Ac<sub>2</sub>O-promoted Diels-Alder reaction<sup>47</sup> between commercially-available aldehyde 24 and acrylonitrile (60%). Reduction of the cyclohexene (H<sub>2</sub>, Pd/C) and the nitrile (LiAlH<sub>4</sub>) followed by tosylation of the resulting amine afforded 26 (53% over three steps). Elimination of the alcohol using the Burgess reagent, and base-promoted isomerization of the alkene48 gave 27 exclusively, with the alkene in conjugation with the aromatic ring. Sulfonamide 27 was then coupled33 with bromoalkyne 28 to afford ynamide 29 (86%). Cycloisomerization of 29 using Pd(OAc)2/bbeda did not proceed at room temperature, but on heating to 45 °C for 3 h gave the desired fused-ring bicyclic product 23 in 50% yield. Surprisingly, use of Pd(OAc)<sub>2</sub> alone<sup>6</sup> proved similarly effective, generating 23 in 54% yield. An NMR reaction profile obtained at 45 °C revealed the reaction of 29 reached completion at around 3 hours.

Pd/C. THF. 98% 3. TsCl, Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub> tol. 110 °C 24 1. Burgess reagent, tol, 100 °C, 90% 2. KO*t*-Bu, DMF, 60 °C, Pd(OAc)<sub>2</sub> (10 mol%) bbeda (10 mol%) toluene, 45 °C, 3 h OTBDPS **TBDPSO** OTBDPS 50% 28 CuSO<sub>4</sub>•5H<sub>2</sub>O, Pd(OAc)<sub>2</sub> (10 mol%) ,10-phenanthroline, K<sub>3</sub>PO, toluene, 80 °C, 86% 23 100 NMR reaction profile for formation of 23 from 29 80 (0.167 M) 60 OTBDPS 40 20 29 2000 4000 6000 8000 10000 12000

Scheme 3 Pd-catalyzed cycloisomerization to fused-ring model 23.

The individual NMR timecourse experiments (at 45 °C) suggested that spiro-ring synthesis should be favoured over fused-ring synthesis, which is contrary to our synthetic design. To further explore this, competition experiments were carried out between ynamides 10 and 29, and ynamides 11 and 29, each in a 1:1 ratio (Fig. 1). We first compared the reactivity of the two sulfonamide-substituted ynamides 10 and 29 (Fig. 1a). This revealed somewhat similar rates of ynamide consumption and product formation for the two substrates. However, to our surprise, the equivalent competition between N-carbamate ynamide 11 and N-tosyl ynamide 29 (Fig. 1b) resulted in quite distinct reaction profiles, in which sulfonamide-substituted fused-ring formation outcompeted carbamate-substituted spiro-ring formation, in spite of the significantly higher rate of reaction of the latter when conducted in isolation (compare Schemes 2b and 3). In a competition setting, this appears to suggest preferential complexation of the sulfonamidesubstituted ynamide to the palladium(II) catalyst over the carbamate-substituted ynamide. Cycloisomerization of the former may proceed via a low-energy intermediate (i.e. a catalyst resting state) that retards the overall observed rate of reaction for both compounds.

This fortunate finding set the stage for execution of the double cycloisomerization cascade, in which the spiro-ring ynamide would feature a carbamate, and the fused-ring ynamide a sulfonamide (*i.e.*, bis-ynamide **30**, Scheme **4**). The




Fig. 1 Competition experiments for Pd-catalyzed cycloisomerization of 10 and 11 vs. 29.

**Edge Article Chemical Science** 

Scheme 4 Pd-catalyzed cascade cycloisomerization to the tetracyclic core of gelsemine, 36

synthesis of 30 began with a Heck reaction<sup>49</sup> of 2-iodoaniline with cyclohexenone, which after carbamoylation of the aniline and Wittig olefination of the ketone gave diene 31 (19% yield over three steps). Regioselective hydroboration/oxidation of 31 and subsequent Mitsunobu reaction gave 32 (57% yield over two steps). After selective carbamate deprotection of the sulfonamide-bearing nitrogen, the resulting sulfonamide was coupled<sup>33</sup> with bromoalkyne 28 to obtain 33 (78% yield over two steps). In preparation for installation of the second ynamide, dichloroenamide 34 was first formed; 42,43 however, treatment of 34 under ynamide-forming conditions (using LiHMDS) led only to unexpected cleavage of the sulfonamide ynamide, 45 with no formation of the desired bis-ynamide 30 being observed.

Pleasingly, this obstacle could be overcome by switching the order of ynamide formation. Thus, dichloroenamide synthesis from 32 (95%) was followed by conversion to the methyl ynamide 35 via elimination (with LiHMDS), copper-catalyzed crosscoupling of the intermediate chloroalkyne with dimethylzine, and cleavage of the 'sulfonamide' carbamate. Coupling of 35 with bromoalkyne 28 successfully afforded bis-ynamide 30. To our delight, subjection of 30 to Pd(OAc)<sub>2</sub>/bbeda-catalyzed cyclization conditions gave tetracyclic compound 36 - the gelsemine core - in 30% yield. Surprisingly, this cascade was accompanied by migration of the double bond of the six membered ring to the (more-substituted) ring junction, presumably by chain-walking of the palladium(II) hydride species. 50-52 This yield is comparable with that of the two individual cyclizations (50% and 70% for 29 and 11 respectively); based on observations with the model system 29 we believe the low yield is mainly impacted by substrate degradation during the first fused-ring cycloisomerization. Finally, we demonstrated that differentiation of the two enamides in this product

could be achieved on reaction with m-CPBA,53 which led to selective oxidative cleavage of the indoline enamide double bond, furnishing the spirooxindole 37 in 68% yield, as required in the gelsemine framework.

### Conclusions

In summary, we have demonstrated a time-resolved palladiumcatalyzed cycloisomerization cascade, in which sequential fused- and then spiro-ring formation accessed the core of the natural product gelsemine. Critical to this chemistry was an understanding of the relative reactivity of the different ynamides involved, where kinetics studies revealed a preferential (but slower) reaction of a sulfonamide-substituted ynamide over a carbamate-substituted vnamide, perhaps suggesting this reaction may proceed via an intermediate resting state that acts as a catalyst reservoir. Application of these findings enabled a 10 step synthesis of the tetracyclic gelsemine core, but perhaps more significantly opens up opportunities for further invention and exploitation of cycloisomerization cascades in synthesis.

# Data availability

All data for experimental procedures and compound characterization are available in the paper and its ESI files.†

#### Author contributions

G. L. and E. A. A. conceived the work. G. L. performed the experiments and analysed the results. E. A. A. directed the project. The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

#### Conflicts of interest

There are no conflicts to declare.

# Acknowledgements

EAA thanks the EPSRC for support (EP/S013172/1).

#### Notes and references

- 1 Y. Hu, M. Bai, Y. Yang and Q. Zhou, Org. Chem. Front., 2017, 4, 2256-2275.
- 2 Y. Yamamoto, Chem. Rev., 2012, 112, 4736-4769.
- 3 V. Michelet, P. Y. Toullec and J. P. Genet, Angew. Chem., Int. Ed., 2008, 47, 4268-4315.
- 4 C. Aubert, O. Buisine and M. Malacria, Chem. Rev., 2002, 102, 813-834.
- 5 B. M. Trost, G. J. Tanoury, M. Lautens, C. Chan and D. T. Macpherson, J. Am. Chem. Soc., 1994, 116, 4255-4267.
- 6 B. M. Trost, D. L. Romero and F. Rise, J. Am. Chem. Soc., 1994, 116, 4268-4278.
- 7 J. Nugent, E. Matoušová, M. G. Banwell and A. C. Willis, J. Org. Chem., 2017, 82, 12569-12589.

- 8 B. M. Trost, Acc. Chem. Res., 1990, 23, 34-42.
- 9 S. Yadav and S. S. V. Ramasastry, *Chem.-Asian J.*, 2020, **15**, 2764–2774.
- 10 S. Mondal, T. Ballav, K. Biswas, S. Ghosh and V. Ganesh, *Eur. J. Org. Chem.*, 2021, 4566–4602.
- 11 M. Lanzi, T. Cañeque, L. Marchiò, R. Maggi, F. Bigi, M. Malacria and G. Maestri, ACS Catal., 2018, 8, 144–147.
- 12 J. Biemolt and E. Ruijter, *Adv. Synth. Catal.*, 2018, **360**, 3821–3871.
- 13 B. Gabriele, R. Mancuso, L. Veltri, I. Ziccarelli and N. Della Ca, *Eur. J. Org. Chem.*, 2019, 5073–5092.
- 14 B. M. Trost and Y. Shi, *J. Am. Chem. Soc.*, 1993, **115**, 9421–9438
- 15 N. Gao, M. G. Banwell and A. C. Willis, *Org. Lett.*, 2017, **19**, 162–165
- 16 X. Ma, N. Gao, M. G. Banwell, P. D. Carr and A. C. Willis, J. Org. Chem., 2017, 82, 4336–4341.
- 17 J. Nugent, E. Matoušová and M. G. Banwell, *Eur. J. Org. Chem.*, 2015, 3771–3778.
- 18 L. Zilke and D. G. Hall, Eur. J. Org. Chem., 2012, 4153-4163.
- 19 P. A. Peixoto, R. Severin, C.-C. Tseng and D. Y. K. Chen, *Angew. Chem., Int. Ed.*, 2011, **50**, 3013–3016.
- 20 B. M. Trost, L. Dong and G. M. Schroeder, J. Am. Chem. Soc., 2005, 127, 10259–10268.
- 21 B. M. Trost and Y. Li, *J. Am. Chem. Soc.*, 1996, **118**, 6625–6633.
- 22 B. M. Trost, P. A. Hipskind, J. Y. L. Chung and C. Chan, *Angew. Chem., Int. Ed.*, 1989, **28**, 1502–1504.
- 23 Y. Lee, E. M. Rochette, J. Kim and D. Y. K. Chen, *Angew. Chem.*, *Int. Ed.*, 2017, **56**, 12250–12254.
- 24 B. M. Trost and C. Min, Nat. Chem., 2020, 12, 568-573.
- 25 P. R. Walker, C. D. Campbell, A. Suleman, G. Carr and E. A. Anderson, *Angew. Chem., Int. Ed.*, 2013, **52**, 9139–9143.
- 26 R. N. Straker, Q. Peng, A. Mekareeya, R. S. Paton and E. A. Anderson, *Nat. Commun.*, 2016, 7, 10109.
- 27 S. J. Mansfield, K. E. Christensen, A. L. Thompson, K. Ma, M. W. Jones, A. Mekareeya and E. A. Anderson, *Angew. Chem., Int. Ed.*, 2017, 56, 14428–14432.
- 28 L. Zhao, H. Yang, R. Li, Y. Tao, X.-F. Guo, E. A. Anderson, A. Whiting and N. Wu, *J. Org. Chem.*, 2021, **86**, 1938–1947.
- 29 L. Andna and L. Miesch, *Org. Biomol. Chem.*, 2019, **17**, 5688–5692.
- 30 X. Zeng, Y. Tu, Z. Zhang, C. You, J. Wu, Z. Ye and J. Zhao, *J. Org. Chem.*, 2019, **84**, 4458–4466.
- 31 Y. Yang, X. Zhang and Y. Liang, *Tetrahedron Lett.*, 2012, 53, 6557–6560.

- 32 A. Coste, G. Karthikeyan, F. Couty and G. Evano, *Angew. Chem., Int. Ed.*, 2009, **48**, 4381–4385.
- 33 X. Zhang, Y. Zhang, J. Huang, R. P. Hsung, K. C. M. Kurtz, J. Oppenheimer, M. E. Petersen, I. K. Sagamanova, L. Shen and M. R. Tracey, J. Org. Chem., 2006, 71, 4170–4177.
- 34 S. Hirano, R. Tanaka, H. Urabe and F. Sato, *Org. Lett.*, 2004, 6, 727–729.
- 35 J. R. Dunetz and R. L. Danheiser, *Org. Lett.*, 2003, 5, 4011–4014.
- 36 B. Witulski and T. Stengel, Angew. Chem., Int. Ed., 1998, 37, 489–492.
- 37 A. Mekareeya, P. R. Walker, A. Couce-Rios, C. D. Campbell, A. Steven, R. S. Paton and E. A. Anderson, *J. Am. Chem. Soc.*, 2017, 139, 10104–10114.
- 38 A. Ghosh and R. G. Carter, *Angew. Chem., Int. Ed.*, 2019, 58, 681–694.
- 39 X. Chen, S. Duan, C. Tao, H. Zhai and F. G. Qiu, *Nat. Commun.*, 2015, **6**, 7204.
- 40 X. Zhou, T. Xiao, Y. Iwama and Y. Qin, *Angew. Chem., Int. Ed.*, 2012, 51, 4909–4912.
- 41 H. Lin and S. J. Danishefsky, *Angew. Chem., Int. Ed.*, 2003, **42**, 36–51.
- 42 S. J. Mansfield, R. C. Smith, J. R. J. Yong, O. L. Garry and E. A. Anderson, *Org. Lett.*, 2019, **21**, 2918–2922.
- 43 S. J. Mansfield, C. D. Campbell, M. W. Jones and E. A. Anderson, *Chem. Commun.*, 2015, 51, 3316–3319.
- 44 M. R. Paleo, L. Castedo and D. Dominguez, *J. Org. Chem.*, 1993, **58**, 2763–2767.
- 45 See the ESI† for details.
- 46 S. Ortgies and A. Breder, Org. Lett., 2015, 17, 2748-2751.
- 47 D. Strübing, A. J. von Wangelin, H. Neumann, D. Gördes, S. Hübner, S. Klaus, A. Spannenberg and M. Beller, *Eur. J. Org. Chem.*, 2005, 107–113.
- 48 M. Hassam, A. Taher, G. E. Arnott, I. R. Green and W. A. L. van Otterlo, *Chem. Rev.*, 2015, **115**, 5462–5569.
- 49 S. E. Reisman, J. M. Ready, M. M. Weiss, A. Hasuoka, M. Hirata, K. Tamaki, T. V. Ovaska, C. J. Smith and J. L. Wood, J. Am. Chem. Soc., 2008, 130, 2087–2100.
- 50 H. Sommer, F. Juliá-Hernández, R. Martin and I. Marek, *ACS Cent. Sci.*, 2018, 4, 153–165.
- 51 T. Kochi, T. Hamasaki, Y. Aoyama, J. Kawasaki and F. Kakiuchi, *J. Am. Chem. Soc.*, 2012, **134**, 16544–16547.
- 52 See also: L. A. Goj and R. A. Widenhoefer, *J. Am. Chem. Soc.*, 2001, **123**, 11133–11147.
- 53 Q. Wang, L. Zhang, J. Yao, G. Qiu, X. Li and H. Zhou, *J. Org. Chem.*, 2018, **83**, 4092–4098.