

Cite this: *Chem. Sci.*, 2023, 14, 7355

All publication charges for this article have been paid for by the Royal Society of Chemistry

Facile preparation of organosilanes from benzylboronates and *gem*-diborylalkanes mediated by $\text{KO}^t\text{Bu}^\dagger$

Man Tang,^a Wenyan Zhu,^a Huaxing Sun,^a Jing Wang,^a Su Jing,^{ib} Minyan Wang,^{*b} Zhuangzhi Shi^{ib} and Jiefeng Hu^{ib} ^{*a}

Methods to efficiently synthesize organosilanes are valuable in the fields of synthetic chemistry and materials science. During the past decades, boron conversion has become a generic and powerful approach for constructing carbon–carbon and other carbon–heteroatom bonds, but its potential application in forming carbon–silicon remains unexplored. Herein, we describe an alkoxide base-promoted deborylative silylation of benzylic organoboronates, geminal bis(boronates) or alkyltriboronates, allowing for straightforward access to synthetically valuable organosilanes. This selective deborylative methodology exhibits operational simplicity, broad substrate scope, excellent functional group compatibility and convenient scalability, providing an effective and complementary platform for the generation of diversified benzyl silanes and silylboronates. Detailed experimental results and calculated studies revealed an unusual mechanistic feature of this C–Si bond formation.

Received 15th May 2023
Accepted 10th June 2023
DOI: 10.1039/d3sc02461j
rsc.li/chemical-science


Introduction

Organosilanes not only play pivotal roles in medicinal chemistry¹ and material science,² but they are also highly functionalized and versatile building blocks for further transformation in synthetic chemistry.³ In the past few decades, considerable attention has been placed on developing synthetic strategies to construct $\text{C}(\text{sp}^3)\text{--Si}$ bonds by cross-coupling reactions in the presence of transition metals, such as palladium, nickel, and copper salts, as catalysts (Fig. 1A).⁴ Highly reactive organometallic species, including Grignard reagents, organolithiums, organozincs or organoaluminums, are emerging as the most attractive substrates for the construction of organosilanes.^{5,6} Although these transformations are synthetically useful, these organometallic reagents exhibit a high basicity and strong nucleophilicity, making them incompatible with many functional groups. Organohalides are often used as precursors to organometallic reagents; therefore, substantial progress has been recently achieved to access silicon-containing molecules by cross-coupling from these abundant and structurally diverse feedstocks to replace preformed organometallic reagents.^{7,8} Despite these advances, a method to easily prepare


organosilanes without reactive coupling partners and metal catalysts is more attractive.

Interconversion of the main group elements has been a long-standing goal in organic chemistry.⁹ In this area, organoboron compounds have become valuable building blocks in organic synthesis because of their flexibility, as they can be synthesized from multiple different functional groups.¹⁰ Significant developments have been achieved for the interconversion of carbon–boron bonds to carbon–carbon and carbon–heteroatom (F, Cl, Br, I, O, N, S ...) bonds (Fig. 1B).^{11,12} In this context, the conversion of C–Si bonds into C–B bonds has been developed through the reaction with boron trihalides (BCl_3 , BBr_3).¹³ At the

(A) Reported routes to organosilanes



(B) Interconversion of the main group elements



(C) Opening the route from organoboronates to silanes (this work)




Fig. 1 Background and discovery. (A) Classic synthesis methods of organosilanes; (B) boron conversion; (C) deborylative silylation.

^aSchool of Chemistry and Molecular Engineering, Nanjing Tech University, Nanjing, 211816, China. E-mail: iasjfh@njtech.edu.cn

^bState Key Laboratory of Coordination Chemistry, School of Chemistry and Chemical Engineering, Nanjing University, Nanjing, 210023, China

[†] Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d3sc02461j>

Table 1 Effect of reaction parameters^a

Entry	Variation from standard conditions	Yield ^b (%)
1	None	92 (84 ^c)
2	Without KO ^t Bu	0
3	KOMe instead of KO ^t Bu	21
4	NaO ^t Bu instead of KO ^t Bu	Trace
5	LiO ^t Bu instead of KO ^t Bu	0
6	KOH instead of KO ^t Bu	0
7	K ₃ PO ₄ instead of KO ^t Bu	0
8	Toluene as the solvent	69
9	ⁿ Octane as the solvent	58
10	MeCN as the solvent	0
11	DMF as the solvent	Trace
12	At 80 °C	64

^a Reaction conditions: **1a** (0.4 mmol), **2a** (0.8 mmol), 2 equiv. of KO^tBu in THF (2 mL), 5 h, at 100 °C under Ar. ^b The yields were determined by GC-MS analysis using an internal standard. ^c Isolated yield.

Table 2 The scope of alkylborons^{abc}

Benzyl substrates^a 3b 87% 3c 68% 3d 62% 3e 49% 3f 57% 3g 72% 3h 60% 3i 61% 3j 67% 3k 79% 3l 63% 3m 65%	Multiborylated substrates^a 3n 74% 3o 77% 3p 79% 3q 73% 3r 70% 3s 73% 3t 78% (82% ^c) 3u 77% 3v 42% 3aa 60% 3ab 64% 3ac 61% 3ad 61% 3ae 67% 3af 49% 3ag 62% 3ah 72% 3ai 69% 3aj 63% 3ak 67% 3al 67% 3am 72% 3an 69% 3ao 63% 3ap 67% 3aq 67% 3ar 67% 3as 67% 3at 67% 3au 67% 3av 67% 3aw 67% 3ax 67% 3ay 67% 3az 67% 3ba 67% 3bb 67% 3bc 67% 3bd 67% 3be 67% 3bf 67% 3bg 67% 3bh 67% 3bi 67% 3bj 67% 3bk 67% 3bl 67% 3bm 67% 3bn 67% 3bo 67% 3bp 67% 3bq 67% 3br 67% 3bs 67% 3bt 67% 3bu 67% 3bv 67% 3bw 67% 3bx 67% 3by 67% 3bz 67% 3ca 67% 3cb 67% 3cc 67% 3cd 67% 3ce 67% 3cf 67% 3cg 67% 3ch 67% 3ci 67% 3cj 67% 3ck 67% 3cl 67% 3cm 67% 3cn 67% 3co 67% 3cp 67% 3cq 67% 3cr 67% 3cs 67% 3ct 67% 3cu 67% 3cv 67% 3cw 67% 3cx 67% 3cy 67% 3cz 67% 3da 67% 3db 67% 3dc 67% 3dd 67% 3de 67% 3df 67% 3dg 67% 3dh 67% 3di 67% 3dj 67% 3dk 67% 3dl 67% 3dm 67% 3dn 67% 3do 67% 3dp 67% 3dq 67% 3dr 67% 3ds 67% 3dt 67% 3du 67% 3dv 67% 3dw 67% 3dx 67% 3dy 67% 3dz 67% 3ea 67% 3eb 67% 3ec 67% 3ed 67% 3ee 67% 3ef 67% 3eg 67% 3eh 67% 3ei 67% 3ej 67% 3ek 67% 3el 67% 3em 67% 3en 67% 3eo 67% 3ep 67% 3eq 67% 3er 67% 3es 67% 3et 67% 3eu 67% 3ev 67% 3ew 67% 3ex 67% 3ey 67% 3ez 67% 3fa 67% 3fb 67% 3fc 67% 3fd 67% 3fe 67% 3ff 67% 3fg 67% 3fh 67% 3fi 67% 3fj 67% 3fk 67% 3fl 67% 3fm 67% 3fn 67% 3fo 67% 3fp 67% 3fq 67% 3fr 67% 3fs 67% 3ft 67% 3fu 67% 3fv 67% 3fw 67% 3fx 67% 3fy 67% 3fz 67% 3ga 67% 3gb 67% 3gc 67% 3gd 67% 3ge 67% 3gf 67% 3gg 67% 3gh 67% 3gi 67% 3gj 67% 3gk 67% 3gl 67% 3gm 67% 3gn 67% 3go 67% 3gp 67% 3gq 67% 3gr 67% 3gs 67% 3gt 67% 3gu 67% 3gv 67% 3gw 67% 3gx 67% 3gy 67% 3gz 67% 3ha 67% 3hb 67% 3hc 67% 3hd 67% 3he 67% 3hf 67% 3hg 67% 3hi 67% 3hj 67% 3hk 67% 3hl 67% 3hm 67% 3hn 67% 3ho 67% 3hp 67% 3hq 67% 3hr 67% 3hs 67% 3ht 67% 3hu 67% 3hv 67% 3hw 67% 3hx 67% 3hy 67% 3hz 67% 3ia 67% 3ib 67% 3ic 67% 3id 67% 3ie 67% 3if 67% 3ig 67% 3ih 67% 3ii 67% 3ij 67% 3ik 67% 3il 67% 3im 67% 3in 67% 3io 67% 3ip 67% 3iq 67% 3ir 67% 3is 67% 3it 67% 3iu 67% 3iv 67% 3iw 67% 3ix 67% 3iy 67% 3iz 67% 3ja 67% 3jb 67% 3jc 67% 3jd 67% 3je 67% 3jf 67% 3jg 67% 3jh 67% 3ji 67% 3jj 67% 3jk 67% 3jl 67% 3jm 67% 3jn 67% 3jo 67% 3jp 67% 3jq 67% 3jr 67% 3js 67% 3jt 67% 3ju 67% 3jv 67% 3jw 67% 3jx 67% 3jy 67% 3jz 67% 3ka 67% 3kb 67% 3kc 67% 3kd 67% 3ke 67% 3kf 67% 3kg 67% 3kh 67% 3ki 67% 3kj 67% 3kl 67% 3km 67% 3kn 67% 3ko 67% 3kp 67% 3kq 67% 3kr 67% 3ks 67% 3kt 67% 3ku 67% 3kv 67% 3kw 67% 3kx 67% 3ky 67% 3kz 67% 3la 67% 3lb 67% 3lc 67% 3ld 67% 3le 67% 3lf 67% 3lg 67% 3lh 67% 3li 67% 3lj 67% 3lk 67% 3ll 67% 3lm 67% 3ln 67% 3lo 67% 3lp 67% 3lq 67% 3lr 67% 3ls 67% 3lt 67% 3lu 67% 3lv 67% 3lw 67% 3lx 67% 3ly 67% 3lz 67% 3ma 67% 3mb 67% 3mc 67% 3md 67% 3me 67% 3mf 67% 3mg 67% 3mh 67% 3mi 67% 3mj 67% 3mk 67% 3ml 67% 3mn 67% 3mo 67% 3mp 67% 3mq 67% 3mr 67% 3ms 67% 3mt 67% 3mu 67% 3mv 67% 3mw 67% 3mx 67% 3my 67% 3mz 67% 3na 67% 3nb 67% 3nc 67% 3nd 67% 3ne 67% 3nf 67% 3ng 67% 3nh 67% 3ni 67% 3nj 67% 3nk 67% 3nl 67% 3nm 67% 3no 67% 3np 67% 3nq 67% 3nr 67% 3ns 67% 3nt 67% 3nu 67% 3nv 67% 3nw 67% 3nx 67% 3ny 67% 3nz 67% 3oa 67% 3ob 67% 3oc 67% 3od 67% 3oe 67% 3of 67% 3og 67% 3oh 67% 3oi 67% 3oj 67% 3ok 67% 3ol 67% 3om 67% 3on 67% 3oo 67% 3op 67% 3oq 67% 3or 67% 3os 67% 3ot 67% 3ou 67% 3ov 67% 3ow 67% 3ox 67% 3oy 67% 3oz 67% 3pa 67% 3pb 67% 3pc 67% 3pd 67% 3pe 67% 3pf 67% 3pg 67% 3ph 67% 3pi 67% 3pj 67% 3pk 67% 3pl 67% 3pm 67% 3pn 67% 3po 67% 3pp 67% 3pq 67% 3pr 67% 3ps 67% 3pt 67% 3pu 67% 3pv 67% 3pw 67% 3px 67% 3py 67% 3pz 67% 3qa 67% 3qb 67% 3qc 67% 3qd 67% 3qe 67% 3qf 67% 3qg 67% 3qh 67% 3qi 67% 3qj 67% 3qk 67% 3ql 67% 3qm 67% 3qn 67% 3qo 67% 3qp 67% 3qq 67% 3qr 67% 3qs 67% 3qt 67% 3qu 67% 3qv 67% 3qw 67% 3qx 67% 3qy 67% 3qz 67% 3ra 67% 3rb 67% 3rc 67% 3rd 67% 3re 67% 3rf 67% 3rg 67% 3rh 67% 3ri 67% 3rj 67% 3rk 67% 3rl 67% 3rm 67% 3rn 67% 3ro 67% 3rp 67% 3rq 67% 3rr 67% 3rs 67% 3rt 67% 3ru 67% 3rv 67% 3rw 67% 3rx 67% 3ry 67% 3rz 67% 3sa 67% 3sb 67% 3sc 67% 3sd 67% 3se 67% 3sf 67% 3sg 67% 3sh 67% 3si 67% 3sj 67% 3sk 67% 3sl 67% 3sm 67% 3sn 67% 3so 67% 3sp 67% 3sq 67% 3sr 67% 3ss 67% 3st 67% 3su 67% 3sv 67% 3sw 67% 3sx 67% 3sy 67% 3sz 67% 3ta 67% 3tb 67% 3tc 67% 3td 67% 3te 67% 3tf 67% 3tg 67% 3th 67% 3ti 67% 3tj 67% 3tk 67% 3tl 67% 3tm 67% 3tn 67% 3to 67% 3tp 67% 3tq 67% 3tr 67% 3ts 67% 3tt 67% 3tu 67% 3tv 67% 3tw 67% 3tx 67% 3ty 67% 3tz 67% 3ua 67% 3ub 67% 3uc 67% 3ud 67% 3ue 67% 3uf 67% 3ug 67% 3uh 67% 3ui 67% 3uj 67% 3uk 67% 3ul 67% 3um 67% 3un 67% 3uo 67% 3up 67% 3uq 67% 3ur 67% 3us 67% 3ut 67% 3uu 67% 3uv 67% 3uw 67% 3ux 67% 3uy 67% 3uz 67% 3va 67% 3vb 67% 3vc 67% 3vd 67% 3ve 67% 3vf 67% 3vg 67% 3vh 67% 3vi 67% 3vj 67% 3vk 67% 3vl 67% 3vm 67% 3vn 67% 3vo 67% 3vp 67% 3vq 67% 3vr 67% 3vs 67% 3vt 67% 3vu 67% 3vv 67% 3vw 67% 3vx 67% 3vy 67% 3vz 67% 3wa 67% 3wb 67% 3wc 67% 3wd 67% 3we 67% 3wf 67% 3wg 67% 3wh 67% 3wi 67% 3wj 67% 3wk 67% 3wl 67% 3wm 67% 3wn 67% 3wo 67% 3wp 67% 3wq 67% 3wr 67% 3ws 67% 3wt 67% 3wu 67% 3wv 67% 3ww 67% 3wx 67% 3wy 67% 3wz 67% 3xa 67% 3xb 67% 3xc 67% 3xd 67% 3xe 67% 3xf 67% 3xg 67% 3xh 67% 3xi 67% 3xj 67% 3xk 67% 3xl 67% 3xm 67% 3xn 67% 3xo 67% 3xp 67% 3xq 67% 3xrr 67% 3xrs 67% 3xrt 67% 3xru 67% 3xrv 67% 3xrw 67% 3xrx 67% 3xry 67% 3xrz 67% 3ya 67% 3yb 67% 3yc 67% 3yd 67% 3ye 67% 3yf 67% 3yg 67% 3yh 67% 3yi 67% 3yj 67% 3yk 67% 3yl 67% 3ym 67% 3yn 67% 3yo 67% 3yp 67% 3yq 67% 3yr 67% 3ys 67% 3yt 67% 3yu 67% 3yv 67% 3yw 67% 3yx 67% 3yy 67% 3yz 67% 3za 67% 3zb 67% 3zc 67% 3zd 67% 3ze 67% 3zf 67% 3zg 67% 3zh 67% 3zi 67% 3zj 67% 3zk 67% 3zl 67% 3zm 67% 3zn 67% 3zo 67% 3zp 67% 3zq 67% 3zr 67% 3zs 67% 3zt 67% 3zu 67% 3zv 67% 3zw 67% 3zx 67% 3zy 67% 3zz 67%

^a Reaction conditions: **1** (0.4 mmol), **2a** (0.8 mmol), 2 equiv. of KO^tBu in THF (2 mL), 5 h, at 100 °C under Ar. ^b Reaction conditions: **1** (0.4 mmol), **2b** (0.8 mmol), 2 equiv. of KO^tBu in ⁿoctane (2 mL), 5 h, at 100 °C under Ar; yields given refer to isolated yields of product. ^c The reaction was carried out in THF; yields given refer to isolated yields of product.


outset of this project, no inverse approach from C–B bonds to C–Si bonds had been developed.

Because of the diagonal relationship between boron and silicon in the periodic table, the boryl and silyl groups exhibit related properties and reactivity for further functionalization, but the latter groups are more stable during storage and


handling.¹⁴ Furthermore, silicon reagents exhibit significant differences in functionality in materials, which are important precursors to commercial polymers and copolymers.¹⁵ Thus, developing a method to prepare organosilanes from related boronates is an attractive goal because it fills the gap in borane–silane interconversion. Here, we develop the first deborylative silylation of benzylic boronate esters or multiboronates to organosilanes mediated by an alkoxide base under transition metal-free conditions (Fig. 1C). In this surprisingly simple approach, the *in situ* formation of boron–ate complexes can react with chlorosilanes through a concerted process to access valuable benzyl silanes and silylboronates.

Results and discussion

We commenced our studies using commercially available 2-benzyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane **1a** and chlorotriisopropylsilane **2a** as model substrates. Based on a systematic examination of reaction conditions, we determined that the following settings were optimal: KO^tBu (2 equiv.) as a base in tetrahydrofuran (THF) at 100 °C. As a result, benzyl-triisopropylsilane **3a** was obtained in 84% isolated yield (Table 1, entry 1). KO^tBu was removed from standard conditions, and the target product could not be detected by GC-MS (Table 1, entry 2). While **3a** was formed in 21% yield with KOM



Scheme 1 Synthetic applications of the deboronative silylation.



Scheme 2 Mechanistic experiments.

synthesis of geminal silylboronates,¹⁶ which exhibit unique reactivity characteristics that allow them to participate in a variety of complexity-generating procedures. A range of carbon chain 1,1-bis(pinacolboronate) esters were tolerated under modified conditions, affording valuable silicated boronates in

70–79% yields (**3n–3t**). In addition, boronates **1u** underwent cross-coupling with chlorotriethylsilane to form highly congested α -quaternary carbon centres **3u** in 77% yield. Although oxygen-containing substrates participated in this reaction, their yields were low (42–46%; **3v–3x**). Substrates possessing fluorine, chlorine, bromine, alkenyl and naphthyl groups were well tolerated in this system (**3y–3ad**). Delightedly, the common heterocyclic cores could favourably carry out these deboronative cross-coupling transformations (**3ae–3ah**). Selective functionalization of 1,2-bis(boronate esters) is a challenging topic in organic synthesis.¹⁷ Therefore, we tried the selective cross-coupling reaction with these bis(boronate esters) and chlorosilanes under standard conditions. We found that this cross-coupling exclusively occurs at the benzyl boron, providing 1,2-silylboronates in moderate to good yields (**3ai–3ak**). Subsequently, 1,1,1-alkyltriboronates and 1,1,2-alkyltriboronates were selectively coupled with chlorosilanes to afford the corresponding products **3al–3ap** in 67–89% yield. Most of these silylboronate compounds are difficult to synthesize by known methods. Primary alkylboronates were also examined, but only trace amounts of silanes **3aq** were detected by GC-MS.

Next, we investigated the scope of chlorosilanes for cross-coupling with geminal boronates **1aa**. Substrates with different carbon chains were amenable to the cross-coupling reaction as well (**3ar–3au**). Notably, chlorosilane with a bulky substituent, such as a *tert*-butyl, showed lower yields (**3av**). The reaction of aryl chlorosilanes could afford target products **3aw**,






Fig. 2 The calculated energy profiles for the alkoxide-promoted cross-coupling of **1a** and **2a**.




Fig. 3 The additional stabilization for substrate geminal boronates.

3ax and **3ay** in satisfactory yields as well. Chlorodisilanes were also feasible substrates with good yields (**3az**). Additionally, the use of a chloro-containing example afforded product **3aaa** in 53% yield. Finally, the presence of alkenyl substituents was tolerated (**3aab** and **3aac**).

To showcase the synthetic applications of this robust deboronative platform, we performed late-stage modification of complex molecules and a series of sequential transformations (Scheme 1). Lithocholic acid derivative **1ar** reacted with Et_3SiCl (**2b**) under optimized conditions, affording product **3aad** in 62% yield. Oleic acid derivative **1as** bearing a *trans*-alkene group

provided silylboronate **3aae** in 71% yield. Interestingly, by continuously treating **1at** with chlorosilanes and KO^tBu , two deborylative silylations occurred, affording geminal disilanes **4** in 67% yield. When using 1,1,2-alkyltriboronate **1au** with two reactive sites as a substrate, the selective process of silylation was observed to give triethyl(1-phenyl-2,2-bis(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)ethyl) silane **3aag**, which can further undergo cross-coupling with alkyl bromide **5a** under the condition of sodium alkoxide (NaO^tBu),¹⁸ providing 1,2-silylboronate **6** in 84% yield. Moreover, successive silylation and allylation led to the desired (1,2-bis(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pent-4-en-1-yl)triethylsilane **7** in 87% yield. These results demonstrate the excellent chemoselectivity of this method. A gram-scale reaction proceeded with **1y**, affording a 76% isolated yield of silylated product **3y**, which can then undergo oxidation or amination processes to afford hydroxyl- or amino-products (**8**, **9**) in good yields, respectively.

To determine the reaction mechanism, we performed some experiments (Scheme 2). First, when **R-1b** (97% ee) was reacted with $\text{Me}_2\text{PhSi-Cl}$ **2h** under our conditions, racemic compound **10** was observed in 42% yield after oxidation treatment. GC-MS was used to detect the course of the reaction, and we found that byproduct **11** formed. Subsequently, *in situ* ^{11}B NMR of the reaction process was performed using benzyl boron ester **1a** and chlorosilane **2a** as substrates, and three new resonances (**I**, **II**, and **III**) appeared at 45 minutes. The resonance at 5.1 ppm (**II**) is

assigned to an “ate” complex.¹⁹ Over time, this characteristic peak gradually disappeared. Based on these observations, we speculate that this transformation involves two processes. The first step was the generation of a carbanion through a deborylative pathway of an “ate” complex **II**. The second step was the nucleophilic reaction of the formed alkyl anion with chlorosilane to generate the final product.

The proposed mechanism was further supported by a detailed density functional theory (DFT) study, as shown in Fig. 2. Initially, the boron atom of **1a** is coordinated with the oxygen of KO^tBu solvated by five tetrahydrofuran molecules, and then the oxygen of **1a** replaces one molecule of THF at potassium to generate **INT1A** with an exothermic energy of 18.3 kcal mol^{−1}. In **INT1A**, the boron centre is more electron-rich than that in **1a**, and the C–B distance is elongated to 1.66 Å from 1.57 Å, decreasing the bond dissociation energy of the C–B bond. Subsequent C–B bond cleavage proceeds smoothly through transition state **TS2A** with an energy barrier of 6.7 kcal mol^{−1}, in which the potassium cation binds to the aromatic ring of **1a** to stabilize the developing negative charge centre and simultaneously release one molecule of THF. The direct cleavage of the C–B bond *via* transition state **TS2B** without the assistance of potassium requires a higher activation free energy (25.3 kcal mol^{−1}). The negative charge centre in intermediate **INT2A** undergoes nucleophilic attack to chlorotriisopropylsilane **2a** *via* transition state **TS3A** with an activation free energy of 28.9 kcal mol^{−1}, producing the desired alkyl silane **3a** and byproduct **INT3A**. The calculational results indicate that the nucleophilic attack step is the rate-determining step, and the overall activation free energy for this transformation is 31.0 kcal mol^{−1}. The nucleophilic attack step for the alkyl anion released from **INT2A** must overcome an overall free energy of 35.7 kcal mol^{−1} through transition state **TS3B**, which is 4.7 kcal mol^{−1} higher than that of **TS3A**. This explains why the deborylative silylation can be promoted by KOMe but failed with sodium alkoxide or lithium alkoxide (Table 1, entries 3–5). Therefore, both the alkoxide anion and potassium cation are indispensable in this transformation, as these entities participate in the formation of a more stable intermediate species, lowering the energy barrier of C–B bond cleavage and C–Si bond formation.

For substrate germinal boronates, the pathways for C–B bond cleavage through transition **TS2A–B** and the subsequent C–Si bond formation *via* **TS3A–B** have the lower energies (Fig. 3). Comparing the energy barrier of transition state **TS3A–B** in the rate-determining step, the stepwise S_N2 pathway through **TS3B–B** was found to be disfavoured kinetically and ruled out. Overall, DFT analysis further reveals that the transition states **TS2A–B** and **TS3A–B** are substantially stabilized by the coordination between the potassium cation and oxygen of the reserved boronate (see ESI† for details).

Conclusions

In summary, we have established a general and practical method for synthesizing alkyl silane derivatives from readily available benzylic boronates, geminal bis(boronates) or

alkyltriboronates. This transition-metal-free transformation features excellent chemoselectivity, broad substrate scope, versatility, and scalability. These borylated silanes can readily be further functionalized by well-established organoboron chemistry to enhance the molecular complexity. In view of these features, this transformation should have high synthetic value in the field of materials and pharmaceuticals.

Data availability

The synthetic procedures, characterization, and spectral data supporting this article have been uploaded as part of the ESI.†

Author contributions

J. H. conceived the project and directed the research. S. J. and Z. S. supervised the mechanistic study. J. H. and Z. S. wrote the paper. M. T., W. Z., J. W. and H. S. performed the experiments. M. W. performed the DFT calculations.

Conflicts of interest

The authors declare no competing financial interest.

Acknowledgements

We are grateful for the financial support from the National Natural Science Foundation of China (21901114), the Start-up Grant from Nanjing Tech University (38274017102) and Tang Scholar. We are grateful to the High Performance Computing Center of Nanjing University for doing the numerical calculations in this paper on its blade cluster system.

Notes and references

- (a) A. K. Franz and S. O. Wilson, *J. Med. Chem.*, 2013, **56**, 388–405; (b) R. Ramesh and D. S. Reddy, *J. Med. Chem.*, 2018, **61**, 3779–3798.
- (a) R. Pietschnig and S. Spirk, *Coord. Chem. Rev.*, 2016, **323**, 87–106; (b) E. A. Marro and R. S. Klausen, *Chem. Mater.*, 2019, **31**, 2202–2211.
- (a) F. Zhao, S. Zhang and Z. Xi, *Chem. Commun.*, 2011, **47**, 4348–4357; (b) S. E. Denmark and A. Ambrosi, *Org. Process Res. Dev.*, 2015, **19**, 982–994; (c) T. Komiyama, Y. Minami and T. Hiyama, *ACS Catal.*, 2017, **7**, 631–651.
- S. Bähr, W. Xue and M. Oestreich, *ACS Catal.*, 2019, **9**, 16–24.
- (a) M. Lacout-Loustalet, J. Dupin, F. Metras and J. Valade, *J. Organomet. Chem.*, 1971, **31**, 187–204; (b) P. J. Lennon, D. P. Mack and Q. E. Thompson, *Organometallics*, 1989, **8**, 1121–1122; (c) A. P. Cinderella, B. Vulovic and D. A. Watson, *J. Am. Chem. Soc.*, 2017, **139**, 7741–7744; (d) B. Vulovic, A. P. Cinderella and D. A. Watson, *ACS Catal.*, 2017, **7**, 8113–8117; (e) S. Mallick, E.-U. Würthwein and A. Studer, *Org. Lett.*, 2020, **22**, 6568–6572.
- (a) K. Tamao, A. Kawachi and K. Ito, *J. Am. Chem. Soc.*, 1992, **114**, 3989–3990; (b) K. Tamao and A. Kawachi, *Adv. Organomet. Chem.*, 1995, **38**, 1–58; (c) C. K. Chu, Y. Liang



- and G. C. Fu, *J. Am. Chem. Soc.*, 2016, **138**, 6404–6407; (d) W. Xue, R. Shishido and M. Oestreich, *Angew. Chem., Int. Ed.*, 2018, **57**, 12141–12145; (e) R. Chandrasekaran, F. T. Pulikkottil and K. S. Elama, *Chem. Sci.*, 2021, **12**, 15719–15726.
- 7 (a) W. Xue, Z.-W. Qu, S. Grimme and M. Oestreich, *J. Am. Chem. Soc.*, 2016, **138**, 14222–14225; (b) W. Xue and M. Oestreich, *Angew. Chem., Int. Ed.*, 2017, **56**, 11649–11652; (c) Y. Takeda, K. Shibuta, S. Aoki, N. Tohnai and S. Minakata, *Chem. Sci.*, 2019, **10**, 8642–8647; (d) J. Scharfbier, B. M. Gross and M. Oestreich, *Angew. Chem., Int. Ed.*, 2020, **59**, 1577–1580.
- 8 (a) J. Duan, K. Wang, G.-L. Xu, S. Kang, L. Qi, X.-Y. Liu and X.-Z. Shu, *Angew. Chem., Int. Ed.*, 2020, **59**, 23083–23088; (b) L. Zhang and M. Oestreich, *Angew. Chem., Int. Ed.*, 2021, **60**, 18587–18590.
- 9 (a) D. A. Petrone, J. Ye and M. Lautens, *Chem. Rev.*, 2016, **116**, 8003–8104; (b) K. Ouyang, W. Hao, W.-X. Zhang and Z. Xi, *Chem. Rev.*, 2015, **115**, 12045–12090; (c) L.-J. Cheng and N. P. Mankad, *Chem. Soc. Rev.*, 2020, **49**, 8036–8064; (d) R. Sharma and M. R. Yadav, *Org. Biomol. Chem.*, 2021, **19**, 5476–5500; (e) F. Mo, D. Qiu, L. Zhang and J. Wang, *Chem. Rev.*, 2021, **121**, 5741–5829; (f) A. Varenikov, E. Shapiro and M. Gandelman, *Chem. Rev.*, 2021, **121**, 412–484; (g) S. Huang, M. Wang and X. Jiang, *Chem. Soc. Rev.*, 2022, **51**, 8351–8377.
- 10 (a) A. J. L. Lennox and G. C. Lloyd-Jones, *Chem. Soc. Rev.*, 2014, **43**, 412–443; (b) A. Fawcett, J. Pradeilles, Y. Wang, T. Mutsuga, E. L. Myers and V. K. Aggarwal, *Science*, 2017, **357**, 283–286; (c) J. Hu, Y. Zhao and Z. Shi, *Nat. Catal.*, 2018, **1**, 860–869; (d) X. Liu, W. Ming, Y. Zhang, A. Friedrich and T. B. Marder, *Angew. Chem., Int. Ed.*, 2019, **58**, 18923–18927; (e) J. Li, H. Wang, Z. Qiu, C. Y. Huang and C.-J. Li, *J. Am. Chem. Soc.*, 2020, **142**, 13011–13020.
- 11 From boron to other main group elements: (a) C. M. Crudden, B. W. Glasspoole and C. J. Lata, *Chem. Commun.*, 2009, 6704–6716; (b) D. Leonori and V. K. Aggarwal, *Angew. Chem., Int. Ed.*, 2015, **54**, 1082–1096; (c) T. C. Wilson, T. Cailly and V. Gouverneur, *Chem. Soc. Rev.*, 2018, **47**, 6990–7005.
- 12 From other main group elements to boron: (a) E. C. Neeve, S. J. Geier, I. A. I. Mkhaliid, S. A. Westcott and T. B. Marder, *Chem. Rev.*, 2016, **116**, 9091–9161; (b) M. Wang and Z. Shi, *Chem. Rev.*, 2020, **120**, 7348–7398; (c) J. Hu, M. Ferger, Z. Shi and T. B. Marder, *Chem. Soc. Rev.*, 2021, **50**, 13129–13188.
- 13 (a) W. Haubold, J. Herdtle, W. Gollinger and W. Einholz, *J. Organomet. Chem.*, 1986, **315**, 1–8; (b) A. A. Kolomeitsev, A. A. Kadyrov, J. Szczepkowska-Sztolcman, M. Milewska, H. Koroniak, G. Bissky, J. A. Bartene and G. Rösenthaller, *Tetrahedron Lett.*, 2003, **44**, 8273–8277; (c) D. L. Crossley, J. Cid, L. D. Curless, M. L. Turner and M. J. Ingleson, *Organometallics*, 2015, **34**, 5767–5774; (d) A. Lik, L. Fritze, L. Müller and H. Helten, *J. Am. Chem. Soc.*, 2017, **139**, 5692–5695; (e) C. You, M. Sakai, C. G. Daniliuc, K. Bergander, S. Yamaguchi and A. Studer, *Angew. Chem., Int. Ed.*, 2021, **60**, 21697–21701; (f) J. Yao, L. Yu, W. Duan and C.-J. Li, *Org. Chem. Front.*, 2023, **10**, 524–530.
- 14 C. Cheng and J. F. Hartwig, *Chem. Rev.*, 2015, **115**, 8946–8975.
- 15 A. M. Muzafarov, *Silicon Polymers*, Springer, Heidelberg, 2011.
- 16 (a) A. A. Szymaniak, C. Zhang, J. R. Coombs and J. P. Morken, *ACS Catal.*, 2018, **8**, 2897–2901; (b) Z. Cheng, J. Guo, Y. Sun, Y. Zheng, Z. Zhou and Z. Lu, *Angew. Chem., Int. Ed.*, 2021, **60**, 22454–22460; (c) Y. You and S. Ge, *Angew. Chem., Int. Ed.*, 2021, **60**, 20684–20688; (d) S. Jin, K. Liu, S. Wang and Q. Song, *J. Am. Chem. Soc.*, 2021, **143**, 13124–13134; (e) W. Sun, L. Xu, Y. Qin and C. Liu, *Nat. Synth.*, 2023, **2**, 413–422.
- 17 (a) S. N. Mlynarski, C. H. Schuster and J. P. Morken, *Nature*, 2014, **505**, 386–390; (b) R. D. Dewhurst and T. B. Marder, *Nat. Chem.*, 2014, **6**, 279–280; (c) L. Xu, S. Zhang and P. Li, *Chem. Soc. Rev.*, 2015, **44**, 8848–8858.
- 18 K. Hong, X. Liu and J. P. Morken, *J. Am. Chem. Soc.*, 2014, **136**, 10581–10584.
- 19 B. Lee and P. J. Chirik, *J. Am. Chem. Soc.*, 2020, **142**, 2429–2437.

