

## CORRECTION


[View Article Online](#)  
[View Journal](#) | [View Issue](#)Cite this: *Chem. Sci.*, 2023, 14, 9608**Correction: Structural tuning of organoruthenium compounds allows oxidative switch to control ER stress pathways and bypass multidrug resistance**Mun Juinn Chow,<sup>ab</sup> Cynthia Licona,<sup>cd</sup> Giorgia Pastorin,<sup>be</sup> Georg Mellitzer,<sup>cd</sup>  
Wee Han Ang<sup>\*ab</sup> and Christian Gaidon<sup>\*cd</sup>

DOI: 10.1039/d3sc90157b

[rsc.li/chemical-science](https://rsc.li/chemical-science)Correction for 'Structural tuning of organoruthenium compounds allows oxidative switch to control ER stress pathways and bypass multidrug resistance' by Mun Juinn Chow *et al.*, *Chem. Sci.*, 2016, 7, 4117–4124, <https://doi.org/10.1039/C6SC00268D>.

The authors regret that an incorrect version of **Fig. 3** was included in the original article, where two incorrect images were used, namely RAS-1H LD treatment and RAS-1H HD treatment, resulting in an unintentional duplication. The correct version of **Fig. 3** is presented here, which is now consistent with Fig. S5a from the ESI of the original publication.

<sup>a</sup>Department of Chemistry, National University of Singapore, 3 Science Drive 3, 117543, Singapore. E-mail: [chmawh@nus.edu.sg](mailto:chmawh@nus.edu.sg); Tel: +65 6516 5131<sup>b</sup>NUS Graduate School for Integrative Sciences and Engineering, Singapore<sup>c</sup>U1113 INSERM, 3 Avenue Molière, Strasbourg 67200, France. E-mail: [gaidon@unistra.fr](mailto:gaidon@unistra.fr); Tel: +33 68 52 53 56<sup>d</sup>Section Oncology, FMTS, Strasbourg University, Strasbourg, France<sup>e</sup>Department of Pharmacy, National University of Singapore, 18 Science Drive 4, 117543, Singapore



**Fig. 3** Complexes RAS-1H and RAS-1T induce early time-point ROS and activate cellular antioxidant defense mechanism. (a) Detection of ROS with carboxy-H<sub>2</sub>DCFDA (20 μM) after treatment with RAS-1H and RAS-1T for 3 h, 6 h and 9 h using a microplate assay. Mean ± s.e.m. (\**p* < 0.05, \*\**p* < 0.01, \*\*\**p* < 0.001, \*\*\*\**p* < 0.0001; Student's *t* test). (b) Detection of ROS with a fluorescence microscope after treatment for 6 h. (c) Western blot analysis of Nrf-2, a central protein in cellular antioxidant defence and (d) expression levels of Nrf-2 target gene in AGS cells after treatment with RAS-1H, RAS-1T and cisplatin at LD and HD for 6 h and 24 h. Homogeneous protein loading determined with reference to actin and gene expression normalized against *thp* levels.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

