



Cite this: *J. Mater. Chem. A*, 2023, **11**, 26821

DOI: 10.1039/d3ta90218h
rsc.li/materials-a

Correction: Designer hydrogenated wrinkled yolk@shell TiO_2 architectures towards advanced visible light photocatalysts for selective alcohol oxidation

Abolfazl Ziarati,^a Alireza Badiei,^{*a} Rafael Luque^{*bc} and Weiyi Ouyang^b

Correction for 'Designer hydrogenated wrinkled yolk@shell TiO_2 architectures towards advanced visible light photocatalysts for selective alcohol oxidation' by Abolfazl Ziarati *et al.*, *J. Mater. Chem. A*, 2018, **6**, 8962–8968, <https://doi.org/10.1039/C8TA02012D>.

The authors regret that in the original Fig. 2 of this article, it appeared as though there was a partial overlap between 2b and 2c. This occurred as two samples were mixed prior to measurements, and different sections were selected during high resolution SEM analysis to showcase two distinct structures, which resulted in partial overlap in the images.

An updated version of Fig. 2 is shown below.




Fig. 2 Alkaline hydrothermal partial etching process of the $\text{Y}@\text{WS}-\text{TiO}_2$ structure at different times. The scale bars are 1 μm .

An independent expert has viewed the original and new images and has concluded that they are consistent with the discussions and conclusions presented.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^aSchool of Chemistry, College of Science, University of Tehran, Tehran, Iran. E-mail: abadiei@khayam.ut.ac.ir; Fax: +98-21-61112614; Tel: +98-21-61112614

^bDepartamento de Química Organica, Universidad de Córdoba, Campus de Rabanales, Edificio Marie Curie, E-14014 Córdoba, Spain. E-mail: rafael.luque@uco.es

^cPeoples Friendship University of Russia (RUDN University), 6 Miklukho-Maklaya str., 117198, Moscow, Russia

