



Cite this: *Chem. Sci.*, 2024, **15**, 16169

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 31st July 2024
Accepted 5th September 2024

DOI: 10.1039/d4sc05111d
rsc.li/chemical-science

Introduction

Indole and indoline are the most common heterocycles widely found in nature, possessing unique biological activities.¹ Among them, ethylamino indole is involved in the entire metabolic process of the human body.² For example, tryptophan is one of the essential amino acids for humans, involved in protein synthesis.³ Melatonin is a key hormone regulating the body's biological clock, and serotonin is an important neurotransmitter closely related to feelings of happiness and well-being.⁴ Specifically, C4-ethylamino indole derivatives have also been proven to be dopamine receptor agonists. One such derivative, ropinirole, is a medication used to treat Parkinson's disease (PD) and restless legs syndrome (RLS). It is one of the most commonly prescribed medications in the United States.⁵ On the other hand, C7-aminoindoline is a type of microtubule protein inhibitor and is considered an important antitumor drug (Fig. 1a).⁶ Based on this, we envision that the one-step synthesis of C4-ethylamine indole and C7-aminoindoline using aziridine through C–N bond cleavage ring-opening reactions and C–H alkylation reactions is of significant importance.

Palladium/norbornene (Pd/NBE) chemistry, namely Catellani-type reactions,⁷ provides a strategy for the multi-functionalization of arenes. This reaction integrates the features of C–H

A switch strategy for the synthesis of C4-ethylamine indole and C7-aminoindoline *via* controllable carbon elimination[†]

Bo-Sheng Zhang, ^{‡*}a Bao-Jie Deng, [‡]a Yuan-Xin Zhi, ^a Tian-Jiao Guo, ^a Yi-Ming Wang, ^a Xue-Ya Gou, ^b Zheng-Jun Quan, ^{‡*}a Xi-Cun Wang ^{‡*}a and Yong-Min Liang ^b

Controllable β -carbon elimination to extrude norbornene remains a long-standing challenge in palladium and norbornene chemistry. Herein, this manuscript describes a switchable synthesis of biologically active C4-ethylaminoindole and C7-aminoindoline scaffolds by controlling β -carbon elimination, utilizing aziridine as a C–H ethylation reagent through a C–N bond cleavage reaction. Furthermore, the protecting groups of the product can be easily removed, offering an unusual method for the synthesis of dopamine receptor agonists.

functionalization and cross-coupling.⁸ In 2000, the cyclization reaction catalyzed by Pd/NBE cooperatively was first discovered by Lautens.⁹ The cyclization reaction has since been widely used in materials chemistry, natural product synthesis, and pharmaceutical synthesis.^{8,10} In 2009, Lautens discovered that using norbornadiene (NBD) instead of norbornene (NBE) in non-Catellani-type cascade cyclization reactions triggers a retro-Diels–Alder reaction, providing a method for synthesizing isoquinolinones and indoles.¹¹ In recent years, Liang,^{10d} Kwong,¹² Cheng,¹³ and our group^{10d,14} developed a series of tandem cyclization reactions combining Catellani-type reactions with retro-Diels–Alder reactions. Interestingly, Cheng discovered that oxanorbornadiene exhibits better retro-Diels–Alder reaction activity than NBD in 2018.^{13a} In recent years, the ring-opening C–H alkylation reaction of strained tricyclic heterocycles was achieved by Lautens,¹⁵ Dong,¹⁶ Zhou,¹⁷ and Liang^{10d,14,18} under Pd/NBE catalysis (Fig. 1c). However, C–H alkylation between *o*-idoanilines and aziridines under Pd/NBE cooperative catalysis is difficult due to the susceptibility of aziridine to nucleophilic attack leading to ring-opening reactions (Fig. 1b).¹⁹

β -carbon elimination to extrude norbornene has consistently been a focal point of interest in Pd/NBE research.^{8g,20} In 2018, the Dong group utilized the steric hindrance effect of C1 norbornene to effectively promote β -carbon elimination and achieve single C–H functionalization of aryl iodides without an *ortho*-substituent.^{20c} In 2019, Dong made a breakthrough in the halogenated olefin version by using norbornene amides.²¹ The Dong group further utilized this strategy to accelerate the extrusion of norbornene, inhibiting the formation of nitrene cyclization products and achieving the introduction of secondary amines in 2024.²² Interestingly, the Jiao group used hybrid cycloolefin ligands to achieve norbornene-like β -carbon elimination.²³ However, controllable β -carbon elimination to extrude


^aGansu International Scientific and Technological Cooperation Base of Water-Retention Chemical Functional Materials, College of Chemistry and Chemical Engineering, Northwest Normal University, Lanzhou, 730070, China. E-mail: zhangbs@nwnu.edu.cn; quanzhengjun@hotmail.com; wangxicun@nwnu.edu.cn

^bState Key Laboratory of Applied Organic Chemistry, Lanzhou University, Lanzhou 730000, China

[†] Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4sc05111d>

[‡] These authors contributed equally.





b) Ring-opening of aziridines





Fig. 1 Selective synthesis of C4-Ethylamine Indole and C7-Aminoindoline.

norbornene remains a long-standing challenge. Herein, this manuscript described a switchable synthesis of biologically active C4-Ethylaminoindole and C7-Aminoindoline scaffolds by controlling the β -carbon elimination, utilizing aziridine as a C–H ethylation reagent through a C–N bond cleavage reaction.


Results and discussion

Initially, we used sterically hindered *N*-*tert*-butyl-*o*-iodoaniline as the substrate, aziridine as the ring-opening C–H alkylating reagent, and norbornadiene (NBD) instead of norbornene (NBE) to attempt to achieve the synthesis of C4-Ethylaminoindole. After carefully studying various reaction parameters, $\text{Pd}(\text{OAc})_2$ and triphenylphosphine were chosen as the Pd/ligand combination, cesium carbonate (Cs_2CO_3) was used as the base, and a mixture of toluene and dioxane served as the solvent. Under an argon atmosphere, the reaction mixture was stirred initially at 90 °C for 12 hours, followed by an increase in temperature to 150 °C and further stirring for 24 hours. This procedure resulted in a 58% isolated yield of the C4-Ethylamino indole 3a, with no formation of the cyclized product 4a resulting from β -carbon elimination. It is worth mentioning that we found that heating to 150 °C in the later stage promoted the retro-Diels–Alder reaction to release cyclopentadiene. Next, deviation experiments from the standard conditions were explored. We first investigated the use of a single solvent instead of a mixed solvent. When toluene was used as the solvent, the yield of C4-Ethylamino indole 3a decreased to 31%, and 8% of C7-Aminoindoline 4a was detected. When dioxane was used as the solvent, the yield of 3a decreased to 40%. Notably, omitting CsBr as an additive also resulted in a yield of 48%. Subsequently, when we attempted to replace NBD with NBE, no target product 3a was detected, while the yield of indoline 4a increased to 16%. Potassium carbonate instead of cesium


c) Three-membered ring cleavage reaction in Pd/NBE chemistry



d) Selective Synthesis of C4-Ethylamine Indole and C7-Aminoindoline (this work)



carbonate as the base reduced the yield of the indole 3a to 30%, with 12% indoline 4a formed. Increasing the amount of cesium carbonate from 1.0 equivalent to 2.0 equivalents decreased the yield of the target product to 43%. Additionally, directly stirring at 140 °C for 24 hours resulted in a yield of 32% for 3a, and formation of 18% of 4a was detected. Finally, we investigated various protecting groups on the nitrogen of *o*-iodoaniline, such as methyl, cyclohexyl, *t*-butoxycarbonyl (Boc), and benzoyl, but none yielded the target product (Table 1).

Table 1 Optimization of reaction conditions for C4-Ethylamine indole^a

Entry	Deviation from the standard conditions	Yield (3a)	Yield (4a)
1	—	58	Trace
2	Toluene or dioxane as solvent	31/40	8/trace
3	Without CsBr	48	Trace
4	NBE instead of NBD	0	16
5	K_2CO_3 instead of Cs_2CO_3	30	12
6	2.0 equiv. of Cs_2CO_3	43	Trace
7	140 °C and 24 h	32	18

^a Standard conditions A: substrate 1a (0.2 mmol), 2a (0.5 mmol, 2.5 equiv.), $\text{Pd}(\text{OAc})_2$ (10 mmol%), PPh_3 (25 mmol%), CsBr (1.0 equiv.), Cs_2CO_3 (1.0 equiv.), NBD (0.6 mmol, 3.0 equiv.), toluene : dioxane (1 : 1, 2 mL) 90 °C 12 h, and then 150 °C, 24 h.



Table 2 Optimization of reaction conditions for C7-aminoindoline^a

Entry	Deviation from the standard conditions	Yield (4a)	Yield (3a)
		Yield (%)	
1	—	56	—
2	PPh ₃ instead of P(<i>p</i> -Cl-C ₆ H ₄) ₃	40	—
3	NBE instead of N1	42	—
4	NBD instead of N1	19	Trace
5	Without KI	39	—

^a Standard conditions B: substrate 1a (0.2 mmol), 2a (0.5 mmol, 2.5 equiv.), Pd(OAc)₂ (10 mmol%), P(*p*-Cl-C₆H₄)₃ (20 mmol%), KI (0.5 equiv.), K₂CO₃ (0.6 mmol, 3.0 equiv.), N1 (0.2 mmol, 1.0 equiv.), toluene (2 mL), 100 °C, 24 h.

Table 3 Investigation of substrate scope^a

		Synthesis of C4-Ethylamine Indole			
1	2	Pd	3 (Condition A) or 4 (Condition B)		
3a, 58%	3b, 51%	3c, 48%	3d, 39%	3e, 25%	3f, 30%
3h, 53%	3i, 29%	3k, 24% (R = Me) 3k', <5% (R = Et)	3j, 39%	3n, 26%	3l, 43%
3m, 38%					
		Synthesis of C7-Aminoindoline			
4a, 56%	4b, 52%	4c, 76%	4d, 78%	4e, 93%	4f, 61%
4g, 56%	4h, 51%	4i, 83%	4j, 91%	4k, 92%	4l, 88%
4m, 79%	4n, 82%	4o, 81%	4p, 73%	4q, 43%	4r, 87%

^a Standard conditions A: substrate 1 (0.2 mmol), 2 (0.5 mmol, 2.5 equiv.), Pd(OAc)₂ (10 mmol%), PPh₃ (25 mmol%), CsBr (1.0 equiv.), Cs₂CO₃ (1.0 equiv.), NBD (0.6 mmol, 3.0 equiv.), toluene : dioxane (1 : 1, 2 mL) 90 °C 12 h, and then 150 °C, 24 h. Standard conditions B: substrate 1 (0.2 mmol), 2 (0.5 mmol, 2.5 equiv.), Pd(OAc)₂ (10 mmol%), P(*p*-Cl-C₆H₄)₃ (20 mmol%), KI (0.5 equiv.), K₂CO₃ (0.6 mmol, 3.0 equiv.), N1 (0.2 mmol, 1.0 equiv.), toluene (2 mL), 100 °C, 24 h.

After achieving optimal conditions for generating C4-ethylamino indole, we aimed to direct the catalytic cycle towards β-carbon elimination to produce C7-aminoindoline 4a. After various conditional screenings, we found that when using *p*-chlorotriphenylphosphine as a ligand and 5-norbornene-2-carboxylate (N1) as a co-catalyst, indoline product 4a was obtained with a yield of 56%. After obtaining the optimal reaction conditions, simple control experiments were conducted. When triphenylphosphine was used instead of P(*p*-Cl-C₆H₄)₃ as the ligand, the yield decreased to 40%. Similarly, replacing N1 with NBE led to a reduced yield of 42%. However, when NBD is used instead of N1, the yield decreases to 19%, which may be due to the inhibition of β-carbon elimination. Lastly, omitting KI as an additive still yielded the target product with a 39% yield (Table 2).

After obtaining the optimal reaction conditions, we first studied the functional group tolerance of C4-ethylamine indole (Table 3). Both electron-donating groups ($-Me$, $-OMe$, tBu , and $-Ph$) and electron-withdrawing groups ($-F$ and $-Cl$) were suitable for this method. Notably, 2-methylaziridine underwent a selective ring-opening reaction to form the corresponding C4-ethylaminoindole product **3k**. More importantly, C4-ethylamino-aza-indole **3n** was successfully synthesized *via* this strategy.

Subsequently, we investigated the substrate scope of C7-aminoindoline. First, we studied the tolerance of functional groups on the indoline ring of the product. Both halogens ($-F$, $-Cl$, and $-Br$) and strong electron-withdrawing groups ($-NO_2$ and $-CO_2Me$) were suitable for the method, and the target products were obtained in 53–93% yield. It is noteworthy that the method can also achieve the biologically active C7-amino-aza-indoline scaffold (**4g**) with good yield. Specifically, 2-methylaziridine underwent a selective ring-opening reaction to produce the 2-methylindoline product **4h**. Additionally, aziridines with different benzenesulfonyl protecting groups on the nitrogen atom can be used to synthesize the corresponding C7-aminoindoline derivatives with antitumor activity.

Finally, we found that the *tert*-butyl group on the nitrogen atom of indole can be easily removed in hydrochloric acid, while the *p*-toluenesulfonyl group can be removed under basic conditions. Since pharmaceutically active indole or indoline molecules often have exposed N-H bonds, this further

enhances the application value of this synthetic method (Fig. 2). Additionally, the resulting C4-ethylaminoindole can be further converted into dopamine receptor agonists using established methods.²⁴

Based on the above experimental results and our previous mechanistic studies,^{10d,14a} we proposed a possible catalytic cycle (Fig. 3). First, *o*-iodoaniline **1a** undergoes oxidative addition with the Pd(0) complex to form intermediate **I**. Subsequently, it undergoes migratory insertion with norbornadiene (NBD) or norbornene (**N1**), followed by C–H bond activation and cyclization in the presence of carbonate, resulting in the formation of the aryl-norbornene-palladacycle ANP intermediate **II**. Among them, the common byproduct **II'**, which is detected in low-yielding cases as shown in Table 3, is generated from the reductive elimination reaction of intermediate **II**. Then, the intermediate **II** undergoes a ring-opening oxidative addition process with aziridine to generate the Pd(IV) intermediate **III**, and the C–H alkylation intermediate **IV** is obtained through reductive elimination. It is worth mentioning that intermediate **IV** can follow two distinct pathways, leading to the selective formation of C4-ethylaminoindole (Pathway A) and C7-aminoindoline (Pathway B), respectively. In Pathway A, intermediate **IV** is attacked by an anion, resulting in the cleavage of the N–Pd bond to form intermediate **V-1**. Its σ bond rotates and undergoes deprotonation to coordinate with the nitrogen atom of aniline, forming intermediate **VI-1**. Finally, the five-membered ring intermediate **VII-1** is obtained through reductive elimination, and it further forms C4-ethylaminoindole *via* a retro-Diels–Alder reaction. Density Functional Theory (DFT) calculations revealed that the retro-Diels–Alder reaction releases 18.2 kcal mol⁻¹, indicating that the process is irreversible (Fig. 2c). In Pathway B, intermediate **IV** selectively undergoes a β -carbon elimination to extrude norbornene, followed by reductive elimination to yield C7-aminoindoline.




Fig. 2 Removal of protecting groups and density functional theory (DFT) calculation.




Fig. 3 Proposed reaction mechanism.



Conclusions

In summary, we developed the first palladium-catalyzed regioselective synthesis of C4-ethylaminoindoles by utilizing the C–N bond ring-opening cleavage reaction of aziridine for the *ortho*-C–H ethylation of iodobenzene. Subsequently, by controlling β -carbon elimination to extrude norbornene, we further achieved the synthesis of the C7-aminoindoles. In addition, the reaction also effectively inhibited the nucleophilic addition of the amine group of *o*-iodoaniline to aziridine, making a smooth Pd/NBE catalytic cycle possible. Moreover, the *tert*-butyl or *p*-toluenesulfonyl protecting groups on C4-ethylaminoindole and C7-aminoindole were easily removed, providing a novel synthetic route to dopamine receptor agonists.

Data availability

All data associated with this study are available in the article and ESI.[†]

Author contributions

Conceptualization, B.-S. Z. and Y.-M. L.; methodology, B.-S. Z. and B.-J. D.; investigation, B.-J. D., Y.-X. Z., T.-J. G., and Y.-M. W.; writing – original draft, B.-S. Z. and B.-J. D.; writing – review & editing, B.-S. Z., B.-J. D., X.-Y. G. and Y.-M. L.; funding acquisition, B.-S. Z., X.-C. W., Z.-J. Q. and Y.-M. L.; resources, B.-S. Z., X.-C. W. and Z.-J. Q.; supervision, B.-S. Z., X.-C. W. and Z.-J. Q.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Financial support was received from the National Natural Science Foundation of China: 22101232 (B.-S. Z.), 22067018 (X.-C. W.), 22061038 (Z.-J. Q.) and 22171114 (Y.-M. L.) and the Long Yuan Youth Innovative Project of Gansu Province (2019–39).

Notes and references

- (a) R. D. Taylor, M. MacCoss and A. D. G. Lawson, Rings in Drugs, *J. Med. Chem.*, 2014, **57**, 5845–5859; (b) C. R. Jamison, J. J. Badillo, J. M. Lipshultz, R. J. Comito and D. W. C. MacMillan, Catalyst-controlled oligomerization for the collective synthesis of polypyridoindoline natural products, *Nat. Chem.*, 2017, **9**, 1165–1169; (c) H. Wei, B. Li, N. Wang, Y. Ma, J. Yu, X. Wang, J. Su and D. Liu, Development and Application of Indolines in Pharmaceuticals, *ChemistryOpen*, 2023, **12**, e202200235.
- (a) F. Tylš, T. Páleníček and J. Horáček, Psilocybin – Summary of knowledge and new perspectives, *Eur. Neuropsychopharmacol.*, 2014, **24**, 342–356; (b) R. Tittarelli, G. Mannocchi, F. Pantano and S. F. Romolo, Recreational Use, Analysis and Toxicity of Tryptamines, *Curr. Neuropharmacol.*, 2015, **13**, 26–46.
- A. Slominski, I. Semak, A. Pisarchik, T. Sweatman, A. Szczesniewski and J. Wortsman, Conversion of L-tryptophan to serotonin and melatonin in human melanoma cells, *FEBS Lett.*, 2002, **511**, 102–106.
- (a) L. F. Mohammad-Zadeh, L. Moses and S. M. Gwaltney-Brant, Serotonin: a review, *J. Vet. Pharmacol. Ther.*, 2008, **31**, 187–199; (b) F. Auld, E. L. Maschauer, I. Morrison, D. J. Skene and R. L. Riha, Evidence for the efficacy of melatonin in the treatment of primary adult sleep disorders, *Sleep Med. Rev.*, 2017, **34**, 10–22.
- (a) J. M. Cassady and J. A. Clemens, Use of 4-(2-di-n-propylaminoethyl)indole or a salt thereof as a presynaptic dopamine autoreceptor stimulant, *US pat.*, 4378368A, 1981; (b) H. Wikstroem, J. H. Lii and N. L. Allinger, The dopaminergic moiety of the ergots. A controversial topic studied with molecular mechanics, *J. Med. Chem.*, 1987, **30**, 1928–1934; (c) D. J. Tompson and D. Vearer, Steady-state pharmacokinetic properties of a 24-hour prolonged-release formulation of ropinirole: Results of two randomized studies in patients with Parkinson's disease, *Clin. Ther.*, 2007, **29**, 2654–2666.
- (a) S. Mehndiratta, Y.-F. Chiang, M.-J. Lai, H.-Y. Lee, M.-C. Chen, C.-C. Kuo, C.-Y. Chang, J.-Y. Chang and J.-P. Liou, Concise syntheses of 7-anilino-indoline-N-benzenesulfonamides as antimitotic and vascular disrupting agents, *Bioorg. Med. Chem.*, 2014, **22**, 4917–4923; (b) S.-Y. Wang, X. Liu, L.-W. Meng, M.-M. Li, Y.-R. Li, G.-X. Yu, J. Song, H.-Y. Zhang, P. Chen, S.-Y. Zhang and T. Hu, Discovery of indoline derivatives as anticancer agents via inhibition of tubulin polymerization, *Bioorg. Med. Chem. Lett.*, 2021, **43**, 128095.
- (a) M. Catellani, F. Frignani and A. Rangoni, A Complex Catalytic Cycle Leading to a Regioselective Synthesis of o,o'-Disubstituted Vinylarenes, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 119–122; (b) D. Dupommier and T. Basset, New trends for transition metal-catalyzed *ortho*/ipso difunctionalizations of arenes, *Chem.*, 2024, DOI: [10.1016/j.chempr.2024.07.003](https://doi.org/10.1016/j.chempr.2024.07.003).
- (a) J. Ye and M. Lautens, Palladium-catalysed norbornene-mediated C–H functionalization of arenes, *Nat. Chem.*, 2015, **7**, 863–870; (b) N. Della Ca', M. Fontana, E. Motti and M. Catellani, Pd/Norbornene: A Winning Combination for Selective Aromatic Functionalization via C–H Bond Activation, *Acc. Chem. Res.*, 2016, **49**, 1389–1400; (c) H.-G. Cheng, S. Chen, R. Chen and Q. Zhou, Palladium(II)-Initiated Catellani-Type Reactions, *Angew. Chem., Int. Ed.*, 2019, **58**, 5832–5844; (d) J. Wang and G. Dong, Palladium/Norbornene Cooperative Catalysis, *Chem. Rev.*, 2019, **119**, 7478–7528; (e) R. Li and G. Dong, Structurally Modified Norbornenes: A Key Factor to Modulate Reaction Selectivity in the Palladium/Norbornene Cooperative Catalysis, *J. Am. Chem. Soc.*, 2020, **142**, 17859–17875; (f) L.-Y. Liu, J. X. Qiao, K.-S. Yeung, W. R. Ewing and J.-Q. Yu, meta-Selective C–H Arylation of Fluoroarenes and Simple Arenes, *Angew. Chem., Int. Ed.*, 2020, **59**, 13831–13835; (g) H. Shi, Y. Lu,



J. Weng, K. L. Bay, X. Chen, K. Tanaka, P. Verma, K. N. Houk and J.-Q. Yu, Differentiation and functionalization of remote C–H bonds in adjacent positions, *Nat. Chem.*, 2020, **12**, 399–404; (h) L. Guillemand, N. Kaplaneris, L. Ackermann and M. J. Johansson, Late-stage C–H functionalization offers new opportunities in drug discovery, *Nat. Rev. Chem.*, 2021, **5**, 522–545; (i) H.-G. Cheng, S. Jia and Q. Zhou, Benzo-Fused-Ring Toolbox Based on Palladium/Norbornene Cooperative Catalysis: Methodology Development and Applications in Natural Product Synthesis, *Acc. Chem. Res.*, 2023, **56**, 573–591.

9 (a) M. Lautens and S. Piguel, A New Route to Fused Aromatic Compounds by Using a Palladium-Catalyzed Alkylation–Alkenylation Sequence, *Angew. Chem., Int. Ed.*, 2000, **39**, 1045–1046; (b) B. Mariampillai, J. Alliot, M. Li and M. Lautens, A Convergent Synthesis of Polysubstituted Aromatic Nitriles via Palladium-Catalyzed C–H Functionalization, *J. Am. Chem. Soc.*, 2007, **129**, 15372–15379; (c) A. Rudolph, N. Rackelmann and M. Lautens, Stereochemical and Mechanistic Investigations of a Palladium-Catalyzed Annulation of Secondary Alkyl Iodides, *Angew. Chem., Int. Ed.*, 2007, **46**, 1485–1488; (d) D. A. Candito and M. Lautens, Palladium-Catalyzed Domino Direct Arylation/N-Arylation: Convenient Synthesis of Phenanthridines, *Angew. Chem., Int. Ed.*, 2009, **48**, 6713–6716; (e) H. Weinstabl, M. Suhartono, Z. Qureshi and M. Lautens, Total Synthesis of (+)-Linoxepin by Utilizing the Catellani Reaction, *Angew. Chem., Int. Ed.*, 2013, **52**, 5305–5308; (f) A. Whyte, M. E. Olson and M. Lautens, Palladium-Catalyzed, Norbornene-Mediated, ortho-Amination ipso-Amidation: Sequential C–N Bond Formation, *Org. Lett.*, 2018, **20**, 345–348.

10 (a) P. Zhao, Y. Guo and X. Luan, Total Synthesis of Dalesconol A by Pd(0)/Norbornene-Catalyzed Three-Fold Domino Reaction and Pd(II)-Catalyzed Trihydroxylation, *J. Am. Chem. Soc.*, 2021, **143**, 21270–21274; (b) M. Bai, S. Jia, J. Zhang, H.-G. Cheng, H. Cong, S. Liu, Z. Huang, Y. Huang, X. Chen and Q. Zhou, A Modular Approach for Diversity-Oriented Synthesis of 1,3-trans-Disubstituted Tetrahydroisoquinolines: Seven-Step Asymmetric Synthesis of Michellamines B and C, *Angew. Chem., Int. Ed.*, 2022, **61**, e202205245; (c) B.-S. Zhang, Y.-X. Yang, J. C. A. Oliveira, Z.-Q. Zhang, S. Warratz, Y.-M. Wang, S.-X. Li, X.-C. Wang, X.-Y. Gou, Y.-M. Liang, Z.-J. Quan and L. Ackermann, Combined C–H amination and intermolecular alkyne insertion for a three-component cyclization, *Cell Rep. Phys. Sci.*, 2023, 101647; (d) B.-S. Zhang, Y. Li, Z. Zhang, Y. An, Y.-H. Wen, X.-Y. Gou, S.-Q. Quan, X.-G. Wang and Y.-M. Liang, Synthesis of C4-Aminated Indoles via a Catellani and Retro-Diels–Alder Strategy, *J. Am. Chem. Soc.*, 2019, **141**, 9731–9738.

11 P. Thansandote, D. G. Hulcoop, M. Langer and M. Lautens, Palladium-Catalyzed Annulation of Haloanilines and Halobenzamides Using Norbornadiene as an Acetylene Synthon: A Route to Functionalized Indolines, Isoquinolinones, and Indoles, *J. Org. Chem.*, 2009, **74**, 1673–1678.

12 W. C. Fu, Z. Wang, W. T. K. Chan, Z. Lin and F. Y. Kwong, Regioselective Synthesis of Polycyclic and Heptagon-embedded Aromatic Compounds through a Versatile π -Extension of Aryl Halides, *Angew. Chem., Int. Ed.*, 2017, **56**, 7166–7170.

13 (a) W. Lv, S. Wen, J. Yu and G. Cheng, Palladium-Catalyzed Ortho-Silylation of Aryl Iodides with Concomitant Arylsilylation of Oxanorbornadiene: Accessing Functionalized (Z)- β -Substituted Vinylsilanes and Their Analogues, *Org. Lett.*, 2018, **20**, 4984–4987; (b) W. Lv, J. Yu, B. Ge, S. Wen and G. Cheng, Palladium-Catalyzed Catellani-Type Bis-silylation and Bis-germanylation of Aryl Iodides and Norbornenes, *J. Org. Chem.*, 2018, **83**, 12683–12693; (c) Q. Tian, J. Ge, Y. Liu, X. Wu, Z. Li and G. Cheng, Solvent-Controlled Enantiodivergent Construction of P(V)-Stereogenic Molecules via Palladium-Catalyzed Annulation of Prochiral N-Aryl Phosphonamides with Aromatic Iodides, *Angew. Chem., Int. Ed.*, 2024, e202409366; (d) Y. Zhang, Y. Chen, Q. Tian, B. Wang and G. Cheng, Palladium-Catalyzed Multicomponent Assembly of (Z)-Alkenylborons via Carbopalladation/Boronation/Retro-Diels–Alder Cascade Reaction, *J. Org. Chem.*, 2023, **88**, 11793–11800.

14 (a) B.-S. Zhang, Z.-Q. Zhang, T.-J. Guo, J. C. A. Oliveira, S. Warratz, B.-J. Deng, Y.-M. Wang, J.-S. Zhou, X.-Y. Gou, X.-C. Wang, Z.-J. Quan and L. Ackermann, Direct Synthesis of C4-Acyl Indoles via C–H Acylation, *Org. Lett.*, 2024, **26**, 4998–5003; (b) Y. An, B.-S. Zhang, Y.-N. Ding, Z. Zhang, X.-Y. Gou, X.-S. Li, X. Wang, Y. Li and Y.-M. Liang, Palladium-catalyzed C–H glycosylation and retro Diels–Alder tandem reaction via structurally modified norbornadienes (smNBDs), *Chem. Sci.*, 2021, **12**, 13144–13150.

15 (a) D. A. Candito and M. Lautens, Exploiting the Chemistry of Strained Rings: Synthesis of Indoles via Domino Reaction of Aryl Iodides with 2H-Azirines, *Org. Lett.*, 2010, **12**, 3312–3315; (b) P. Thansandote, M. Raemy, A. Rudolph and M. Lautens, Synthesis of Benzannulated N-Heterocycles by a Palladium-Catalyzed C–C/C–N Coupling of Bromoalkylamines, *Org. Lett.*, 2007, **9**, 5255–5258.

16 (a) R. Li, F. Liu and G. Dong, Palladium-catalyzed asymmetric annulation between aryl iodides and racemic epoxides using a chiral norbornene cocatalyst, *Org. Chem. Front.*, 2018, **5**, 3108–3112; (b) R. Li and G. Dong, Direct Annulation between Aryl Iodides and Epoxides through Palladium/Norbornene Cooperative Catalysis, *Angew. Chem., Int. Ed.*, 2018, **57**, 1697–1701; (c) Z. Wang, Y. Kuninobu and M. Kanai, Palladium-Catalyzed Oxirane-Opening Reaction with Arenes via C–H Bond Activation, *J. Am. Chem. Soc.*, 2015, **137**, 6140–6143; (d) G. Cheng, T.-J. Li and J.-Q. Yu, Practical Pd(II)-Catalyzed C–H Alkylation with Epoxides: One-Step Syntheses of 3,4-Dihydroisocoumarins, *J. Am. Chem. Soc.*, 2015, **137**, 10950–10953.

17 (a) C. Wu, H.-G. Cheng, R. Chen, H. Chen, Z.-S. Liu, J. Zhang, Y. Zhang, Y. Zhu, Z. Geng and Q. Zhou, Convergent syntheses of 2,3-dihydrobenzofurans via a Catellani strategy, *Org. Chem. Front.*, 2018, **5**, 2533–2536; (b) G. Qian,



M. Bai, S. Gao, H. Chen, S. Zhou, H.-G. Cheng, W. Yan and Q. Zhou, Modular One-Step Three-Component Synthesis of Tetrahydroisoquinolines Using a Catellani Strategy, *Angew. Chem., Int. Ed.*, 2018, **57**, 10980–10984; (c) H.-G. Cheng, C. Wu, H. Chen, R. Chen, G. Qian, Z. Geng, Q. Wei, Y. Xia, J. Zhang, Y. Zhang and Q. Zhou, Epoxides as Alkylating Reagents for the Catellani Reaction, *Angew. Chem., Int. Ed.*, 2018, **57**, 3444–3448.

18 C. Liu, Y. Liang, N. Zheng, B.-S. Zhang, Y. Feng, S. Bi and Y.-M. Liang, Synthesis of indolines via a palladium/norbornene-catalyzed reaction of aziridines with aryl iodides, *Chem. Commun.*, 2018, **54**, 3407–3410.

19 (a) Z. Wang, Y.-T. Cui, Z.-B. Xu and J. Qu, Hot Water-Promoted Ring-Opening of Epoxides and Aziridines by Water and Other Nucleophiles, *J. Org. Chem.*, 2008, **73**, 2270–2274; (b) M. Zhu and B. Moasser, Aqueous ring opening of N-tosylaziridine with aniline derivatives, *Tetrahedron Lett.*, 2012, **53**, 2288–2291; (c) S. Sabir, G. Kumar, V. P. Verma and J. L. Jat, Aziridine Ring Opening: An Overview of Sustainable Methods, *ChemistrySelect*, 2018, **3**, 3702–3711.

20 (a) M. Catellani and E. Motti, Selective aryl coupling via palladacycles: a new route to m-alkylbiphenyls or m-terphenyls, *New J. Chem.*, 1998, **22**, 759–761; (b) Z. Dong, G. Lu, J. Wang, P. Liu and G. Dong, Modular ipso/ortho Difunctionalization of Aryl Bromides via Palladium/Norbornene Cooperative Catalysis, *J. Am. Chem. Soc.*, 2018, **140**, 8551–8562; (c) J. Wang, R. Li, Z. Dong, P. Liu and G. Dong, Complementary site-selectivity in arene functionalization enabled by overcoming the ortho constraint in palladium/norbornene catalysis, *Nat. Chem.*, 2018, **10**, 866–872; (d) H. Shi, A. N. Herron, Y. Shao, Q. Shao and J.-Q. Yu, Enantioselective remote meta-C–H arylation and alkylation via a chiral transient mediator, *Nature*, 2018, **558**, 581–585; (e) K.-J. Xiao, L. Chu, G. Chen and J.-Q. Yu, Kinetic Resolution of Benzylamines via Palladium(II)-Catalyzed C–H Cross-Coupling, *J. Am. Chem. Soc.*, 2016, **138**, 7796–7800.

21 J. Wang, Z. Dong, C. Yang and G. Dong, Modular and regioselective synthesis of all-carbon tetrasubstituted olefins enabled by an alkenyl Catellani reaction, *Nat. Chem.*, 2019, **11**, 1106–1112.

22 X. Liu, Q. Zhu and G. Dong, Beyond Tertiary Amines: Introducing Secondary Amines by Palladium/Norbornene-Catalyzed Ortho Amination, *Angew. Chem., Int. Ed.*, 2024, **63**, e202404042.

23 (a) Y.-X. Zheng and L. Jiao, Hybrid cycloolefin ligands for palladium-olefin cooperative catalysis, *Nat. Synth.*, 2022, **1**, 180–187; (b) F.-Y. Wang, Y.-X. Li and L. Jiao, Functionalized Cycloolefin Ligand as a Solution to Ortho-Constraint in the Catellani-Type Reaction, *J. Am. Chem. Soc.*, 2023, **145**, 4871–4881.

24 (a) T. Ankner and G. Hilmersson, Instantaneous Deprotection of Tosylamides and Esters with SmI₂/Amine/Water, *Org. Lett.*, 2009, **11**, 503–506; (b) X. Jiang, C. Zheng, L. Lei, K. Lin and C. Yu, Synthesis of 2-Oxindoles from Substituted Indoles by Hypervalent-Iodine Oxidation, *Eur. J. Org. Chem.*, 2018, **2018**, 1437–1442.

