



Cite this: *Analyst*, 2024, **149**, 2481

## Correction: Diamond nanowires modified with poly[3-(pyrrolyl)carboxylic acid] for the immobilization of histidine-tagged peptides

Palaniappan Subramanian,<sup>a</sup> Ievgen Mazurenko,<sup>b</sup> Vladimir Zaitsev,<sup>b,c</sup>  
 Yannick Coffinier,<sup>a</sup> Rabah Boukherroub<sup>a</sup> and Sabine Szunerits<sup>\*a</sup>

DOI: 10.1039/d4an90026j

[rsc.li/analyst](https://rsc.li/analyst)

Correction for 'Diamond nanowires modified with poly[3-(pyrrolyl)carboxylic acid] for the immobilization of histidine-tagged peptides' by Palaniappan Subramanian *et al.*, *Analyst*, 2014, **139**, 4343–4349, <https://doi.org/10.1039/C4AN00146J>.

In the original version of this article, Fig. 1C mistakenly shows the SEM image of BDD NWs after electrochemical deposition of pyrrole-3-carboxylic acid (100 mM) in TBATFB (0.1 M)/acetonitrile solution by electrochemical potentiodynamic cycling instead of the film prepared by applying a charge of 23 mC cm<sup>−2</sup>. In the revised graphic provided below, the correct morphology of BDD NWs coated with polypyrrole-3-carboxylic film formed by applying a charge of 23 mC cm<sup>−2</sup> is provided (right image). This correction does not change the conclusions of the work.


<sup>a</sup>Institut de Recherche Interdisciplinaire (IRI), CNRS USR 3078, Université Lille1, Parc de la Haute Borne, 50 avenue de Halley, BP 70478, 59658 Villeneuve d'Ascq, France.

E-mail: [Sabine.Szunerits@iri.univ-lille1.fr](mailto:Sabine.Szunerits@iri.univ-lille1.fr)

<sup>b</sup>Department of Analytical Chemistry, Taras Shevchenko National University of Kyiv, Volodymyrska str., 62a, 01033 Kyiv, Ukraine

<sup>c</sup>Chemistry Department, Pontifical Catholic University of Rio de Janeiro, 225 Rua Marquês de São Vicente, 22451 Rio de Janeiro, Brazil





**Fig. 1** (A) SEM images of BDD NWs before (left) and after cycling 4 times in pyrrole-3-carboxylic acid (100 mM in TBATFB (0.1 M)/acetonitrile) (right). (B) Successive cyclic voltammetric  $i$ - $E$  curves of BDD NWs in a solution of pyrrole-3-carboxylic acid (100 mM) in TBATFB (0.1 M)/acetonitrile solution, scan rate =  $0.05 \text{ V s}^{-1}$ ; first (black line), second (grey line), third (blue line), fourth (green line). (C) SEM images of BDD NWs after electrochemical deposition of pyrrole-3-carboxylic acid (100 mM) in TBATFB (0.1 M)/acetonitrile solution at  $E = +1.2 \text{ V}$  for different deposition charges:  $2 \text{ mC cm}^{-2}$  (left),  $11 \text{ mC cm}^{-2}$  (middle), and  $23 \text{ mC cm}^{-2}$  (right).

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

