

CORRECTION

[View Article Online](#)
[View Journal](#) | [View Issue](#)



Cite this: *RSC Chem. Biol.*, 2024, 5, 800

Correction: Virtual screening, identification and *in vitro* validation of small molecule GDP-mannose dehydrogenase inhibitors

Jonathan P. Dolan,^{ab} Sanaz Ahmadipour,^e Alice J. C. Wahart,^{ab} Aisling Ni Cheallaigh,^{ab} Suat Sari,^f Chatchakorn Eurtivong,^g Marcelo A. Lima,^{bd} Mark A. Skidmore,^{bd} Konstantin P. Volcho,^h Jóhannes Reynisson,^{bc} Robert A. Field^e and Gavin J. Miller^{*ab}

DOI: 10.1039/d4cb90026j

rsc.li/rsc-chembio

Correction for 'Virtual screening, identification and *in vitro* validation of small molecule GDP-mannose dehydrogenase inhibitors' by Jonathan P. Dolan *et al.*, *RSC Chem. Biol.*, 2023, **4**, 865–870, <https://doi.org/10.1039/D3CB00126A>.

The authors regret an error in the published article whereby the stereochemistry of one carbon within compound **13** in Fig. 2 and 3 was incorrect. All other designations of compound **13** in the article and supplementary information were correct. The corrected article figures are below.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Lennard-Jones Laboratory, School of Chemical & Physical Sciences, Keele University, Keele, Staffordshire, ST5 5BG, UK. E-mail: g.j.miller@keele.ac.uk

^b Centre for Glycoscience, Keele University, Keele, Staffordshire, ST5 5BG, UK

^c Hornbeam Building, School of Pharmacy & Bioengineering, Keele University, Keele, Staffordshire, ST5 5BG, UK

^d School of Life Sciences, Keele University, Keele, Staffordshire, ST5 5BG, UK

^e Department of Chemistry & Manchester Institute of Biotechnology, The University of Manchester, 131 Princess Street, Manchester, M1 7DN, UK

^f Hacettepe University, Faculty of Pharmacy, Department of Pharmaceutical Chemistry, 06100, Ankara, Turkey

^g Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Mahidol University, 447 Si Ayutthaya Road, Ratchathewi, Bangkok 10400, Thailand

^h N. Vorozhtsov Novosibirsk Institute of Organic Chemistry, Siberian Branch of the Russian Academy of Sciences, 630090 Novosibirsk, Russia






Fig. 2 Bar chart comparing percentage NADH production in the presence of each of 21 potential inhibitors without preincubation with GMD (solid bars) and with preincubation for 1 hour with GMD (hatched bars). Complete structure panel is shown in the ESI,† Section S1.2. The 6 best performing compounds are highlighted blue. Percentage NADH production was determined relative to a positive control containing no inhibitor and **1**.




Fig. 3 (A) Inhibition of GMD with hit **13**, determined by fluorescence of NADH. Error bars indicate the standard error of three measurements. (B) ESI-MS of GMD (47598.6 Da) before incubation with **13**. (C) ESI-MS of GMD after overnight incubation with **13**, showing the formation of a single covalent GMD-**13** adduct (49045.7 Da).