ChemComm



COMMUNICATION

View Article Online View Journal | View Issue

Check for updates

Cite this: Chem. Commun., 2024, 60, 1611

Facile synthesis of dibenzothiophene S-oxides from sulfinate esters[†]

Yukiko Kumagai.ª Akihiro Kobavashi.^{ab} Keisuke Nakamura^a and Suguru Yoshida 厄 *ª

Received 21st November 2023, Accepted 9th January 2024

DOI: 10.1039/d3cc05703h

rsc.li/chemcomm

An efficient method to prepare dibenzothiophene S-oxides is disclosed. Suzuki–Miyaura cross-coupling of 2-bromoaryl sulfinate esters with arylboronic acids selectively at the bromo group followed by electrophilic cyclization of the resulting sulfinate ester moiety provides diverse dibenzothiophene S-oxides. Further transformations including Pummerer-type C–H propargylation and aryne reactions realize to synthesize highly functionalized dibenzothiophene derivatives.

Dibenzothiophenes and their derivatives having S-oxide or S,Sdioxide moieties are of great significance in broad research fields such as pharmaceutical sciences, materials chemistry, and chemical biology (Fig. 1A).¹⁻³ Particularly, dibenzothiophene Soxides are gaining attention as precursors to generate atomic oxygen triggered by UV irradiation, which can facilitate DNA cleavage, oxidation of adenosine-S'-phosphosulfate kinase, and so on.¹ In addition, recent advances in synthetic methods through the activation of S-O bond have allowed us to prepare diverse functionalized dibenzothiophenes from dibenzothiophene Soxides (Fig. 1B).⁴ Despite the importance of dibenzothiophene Soxides, it is not always easy to synthesize highly functionalized dibenzothiophene S-oxides in a conventional manner, which involves thiophene ring formation of biaryls using S₈ and subsequent S-oxidation with oxidants such as mCPBA (Fig. 1C).^{5,6} Herein, we disclose a novel method to prepare dibenzothiophene S-oxides from 2-bromoaryl-substituted sulfinate esters by Brselective Suzuki-Miyaura coupling followed by intramolecular electrophilic sulfinylation (Fig. 1D). Since we recently found that sulfinate esters can be activated easily by triflic anhydride (Tf₂O)



Fig. 1 (A) Significant dibenzothiophene derivatives. (B) Transformations of dibenzothiophene *S*-oxide. (C) Representative synthesis of dibenzothiophene *S*-oxide. (D) This work.

realizing *S*-allylation with allylsilanes, we conceived the synthesis of dibenzothiophene *S*-oxides from 2-biarylyl sulfinate ester through the electrophilic activation.^{7,8}

We succeeded in the synthesis of dibenzothiophene *S*-oxide **4a** through Br-selective coupling and subsequent cyclization by electrophilic activation (Table 1). Treatment of 2-bromobenzenesulfinic acid methyl ester (**1a**)⁹ with 4-tolylboronic acid (**2a**) in the presence of potassium phosphate and a catalytic amount of $(amphos)_2PdCl_2$ afforded 2-(4-tolyl)benzenesulfinic acid methyl ester (**3a**) in an excellent yield. Of note, the coupling proceeded selectively at the bromo group, where palladium-catalyzed sulfoxide formation was not observed.¹⁰ Then, reaction conditions were screened for the cyclization of methyl 2-(4-tolyl)benzenesulfinate (**3a**) by electrophilic activation using acid anhydrides. As a result, we successfully synthesized dibenzothiophene *S*-oxide **4a** in high yield from **3a** with Tf₂O. Indeed, treatment of **3a** with Tf₂O in dichloromethane at room temperature followed by the addition of aqueous sodium

^a Department of Biological Science and Technology, Faculty of Advanced Engineering, Tokyo University of Science, 6-3-1 Niijuku, Katsushika-ku, Tokyo 125-8585, Japan. E-mail: s-yoshida@rs.tus.ac.jp

^b Laboratory of Chemical Bioscience, Institute of Biomaterials and Bioengineering, Tokyo Medical and Dental University (TMDU), 2-3-10 Kanda-Surugadai, Chiyoda-ku, Tokyo 101-0062, Japan

[†] Electronic supplementary information (ESI) available: Experimental procedures, characterization for new compounds including NMR spectra. See DOI: https://doi.org/10.1039/d3cc05703h

Table 1 Optimization of the reaction conditions

O S OMe (amphos) ₂ PdCl ₂ (1.0 mol %) K ₃ PO ₄ (2.0 equiv) + H ₂ O (v/v = 10/1) Me	O S OMe 3a	activator 2.0 equiv) solv. temp. 1 h	O S 4a
Activator	Solv.	Temp.	Yield (%) ^a
Tf_2O	CH_2Cl_2	rt	90
Tf_2O			
$2,6-(t-Bu)_2 py^b$	CH_2Cl_2	rt	88
$(CF_3CO)_2O$	CH_2Cl_2	rt	0
$(CH_3CO)_2O$	CH_2Cl_2	rt	0
TfOH	CH_2Cl_2	rt	51
Tf_2O	CH_2Cl_2	0 °C	97
Tf_2O	CH_2Cl_2	0 °C	83
Tf_2O	CH_2Cl_2	−40 °C	0
Tf_2O	Toluene	rt	3
Tf_2O	CH_3CN	rt	0
Tf_2O	CH_3NO_2	rt	0
Tf_2O	Et_2O	rt	97
Tf_2O	Et_2O	0 °C	Quant.
Tf_2O			-
$2,6-(t-Bu)_2 py^b$	Et_2O	0 °C	88
	$\begin{array}{c} 0\\ \textbf{S}\\ \textbf{OMe}\\ \textbf{Br 1a}\\ \textbf{K}_{3}\text{PO}_{4}\\ \textbf{(1.0 mol \%)}\\ \textbf{Fr 1a}\\ \textbf{K}_{3}\text{PO}_{4}\\ \textbf{(2.0 equiv)}\\ \textbf{H}_{2}O(\textbf{v}\textbf{V}=10/1)\\ \textbf{H}_{2}O(\textbf{v}V=10/1)\\ \textbf{H}_{2}O(\textbf{v}V=10/1$	$\begin{array}{c} 0\\ \textbf{S}\\ \textbf{OMe} \\ (1.0 \text{ mol} \%)_2 PdCl_2\\ (1.0 \text{ mol} \%)_6 \\ \textbf{Fr} 1a \\ (2.0 \text{ equiv})\\ +\\ \textbf{H}_2O (vV = 10/1)\\ 100 ^\circ\text{C}; 94\% \\ \textbf{Me} \end{array} \qquad \begin{array}{c} 0\\ \textbf{K}_3PO_4\\ \textbf{K}_4PO_4\\ $	$\begin{array}{c c} 0\\ S\\ OMe & (amphos)_2 PdCl_2\\ (1.0 \text{ mol } \%)\\ Br \ 1a & (2.0 \text{ equiv})\\ +\\ & (2.0 \text{ equiv})\\ & (2.0 \text{ equiv})\\ & (2.0 \text{ equiv})\\ +\\ & (2.0 \text{ equiv})\\ & (2.$

 a Isolated yields. b The reaction was conducted using 2,6-di(*tert*-butyl)-pyridine (4.0 equiv.) with Tf₂O (2.0 equiv.). c Gram-scale synthesis. See details in the ESI.

bicarbonate provided the desired dibenzothiophene S-oxide 4a in high yield, where side-products from further Pummerer-type reactions were not observed (entry 1). The reaction also took place smoothly when using Tf₂O with 2,6-di(t-butyl)pyridine as a base (entry 2). In contrast, trifluoroacetic anhydride (TFAA) or acetic anhydride did not activate sulfinate ester 3a (entries 3 and 4). Dibenzothiophene S-oxide 4a was also obtained in moderate vield using triflic acid (TfOH) (entry 5). The yield was slightly improved when the reaction was conducted at 0 °C to realize the synthesis of dibenzothiophene S-oxide 4a in an excellent yield (entry 6). We accomplished the gram-scale synthesis of 4a in good yield (entry 7). No reaction took place at -40 °C (entry 8). While the efficient synthesis of cyclized product 4a failed when the reaction was conducted in toluene, acetonitrile, or nitromethane (entries 9-11), we also succeeded in the facile synthesis of dibenzothiophene S-oxide in diethyl ether at room temperature or 0 °C (entries 12 and 13). The synthesis of 4a in the presence of 2,6-di(t-butyl)pyridine was achieved also in diethyl ether (entry 14). We also prepared dibenzothiophene Soxide 4a from 1a in high yield through a single silica-gel column chromatography, obviously indicating good practicality of the benzothiophene S-oxide synthesis from sulfinate esters (Fig. 2A).

A wide range of dibenzothiophene *S*-oxides having diverse functional groups were prepared from sulfinate ester **1a** and various arylboronic acids **2** (Fig. 2B and C). Due to the good solubility of dichloromethane for broad aromatic compounds, we chose the conditions using dichloromethane for synthesizing various dibenzothiophene *S*-oxides. For example, we succeeded in the synthesis of dibenzothiphene *S*-oxides **4b** and **4c** in good yields by the 2-step protocol without damaging fluoro and chloro groups. In the case of synthesis of dibenzothiophene *S*-oxide **4d** from **1a**



Fig. 2 (A) Synthesis of **4a** with a single silica-gel column chromatography purification. (B) General scheme for synthesizing various dibenzothiophene *S*-oxides. (C) Results using various arylboronic acids **2** with **1a**. (D) Results using various sulfinate esters **1** with **2a**. (E) Synthesis of benzothiophene *S*-oxide **6**. ^a Et₂O was used as a solvent instead of CH₂Cl₂. ^bThe reaction was conducted in the presence of 2,6-di(*tert*-butyl)pyridine (4.0 equiv.). See details in the ESI.† amphos = di(*tert*-butyl)(4-(dimethylamino)phenyl)phosphine.

and 4-methoxyphenylboronic acid, the electrophilic cyclization of methyl 2-(4-methoxyphenyl)benzenesulfinate bearing an electronrich aryl group took place efficiently when using diethyl ether as a solvent.¹¹ We succeeded in the preparation of 2-methyldibenzothiophene *S*-oxide (**4e**) along with minor isomer **4e'** from 3-methylboronic acid by efficient Suzuki–Miyaura coupling and following intramolecular arylation, in which cyclization at sterically unhindered site predominantly proceeded. The 2-step protocol using sulfinate ester **1a** and bulky 2-methylphenyl boronic acid allowed us to synthesize dibenzothiophene *S*-oxide **4f** in high yields. We successfully synthesized π -extended dibenzothiophene *S*-oxide **4g**, which was reported as a promising dibenzothiophene

ChemComm

S-oxide derivative having an extended chromophore to red-shift the absorption spectra for atomic oxygen precursors,¹² from 2naphthylboronic acid through regioselective cyclization, where the regioisomer was not detected. It is worth noting that a variety of heteroarylboronic acids participated in the coupling–cyclization protocol to furnish heteroaromatic ring-fused dibenzothiophene *S*oxides **4h**–**4k**. When using 3-thienyl- or 3-furylboronic acid, intramolecular *S*-arylation proceeded at 2-position selectively to afford dibenzothiophene *S*-oxide analog **4h** or **4j**, respectively. The synthesis of furan- or benzo[*b*]furan-fused benzo[*b*]thiophene **4j** or **4k** was realized by the addition of 2,6-di(*tert*-butyl)pyridine as a base. We also succeeded in the preparation of benzo[*b*]thiophene *S*oxide **6** from sulfinate ester **1a** and 2-styrylboronic acid (**5**) through the intramolecular *S*-alkenylation (Fig. 2E).

Various sulfinate esters served in synthesizing dibenzothiophene *S*-oxides having various electron-donating or electron-withdrawing functional groups (Fig. 2B and D). Indeed, 2,7-dimethyl- and 2,4,7-trimethyl-substituted unsymmetric dibenzothiophene *S*-oxides **41** and **4m** were prepared efficiently from 4-methyl- and 4,6-dimethyl-substituted 2-bromobenzenesulfinic acid esters, respectively. The synthesis of dibenzothiophene *S*-oxide **4n** bearing a methylenedioxy moiety was achieved efficiently through cross-coupling and cyclization in diethyl ether. We also accomplished the preparation of dibenzothiophene *S*-oxides **4o**–**4q** leaving electron-withdrawing chloro, trifluoromethyl, and ethoxycarbonyl groups untouched. These results obviously indicated that Br-selective coupling and following cyclization were not affected by a broad range of electron-donating or -withdrawing functional groups.

To gain insight into the reaction mechanism of the electrophilic cyclization of sulfinate esters, we then isolated sulfuran intermediate 7 by changing the protocol (Fig. 3A). Indeed, after treatment of sulfinate ester **3a** with Tf₂O, the resulting mixture was quenched with solid sodium bicarbonate instead of aqueous sodium bicarbonate. Then, filtrating the mixture to remove

sodium bicarbonate and washing the resulting filtrate with n-hexane furnished sulfuran 7 quantitatively. We achieved the synthesis of dibenzothiophene S-oxide 4a with the hydrolysis of sulfuran 7 with aqueous sodium bicarbonate. Based on this result, we proposed a plausible reaction mechanism for the dibenzothiophene S-oxide synthesis (Fig. 3B). First, Tf₂O activates S=O moiety to form sulfuran intermediate I. Second, electrophilic aromatic substitution at the electrophilic sulfur provides sulfuran 7. Finally, hydrolysis of sulfuran 7 with aqueous sodium bicarbonate furnishes dibenzothiophene S-oxide 4a via the nucleophilic attack of water or bicarbonate anion to the sulfur atom. When sulfoxide 8 was treated with Tf₂O followed by the addition of sodium bicarbonate, dibenzothiophene 9a was obtained in good yield through the demethylative hydrolysis of sulfonium intermediate III, where sulfoxide 4a was not obtained (Fig. 3C).¹³ This result shows that the difference between sulfinate esters and sulfoxides in the electrophilic cyclization lies in the reaction sites in the hydrolysis.

Good transformability of dibenzothiophene *S*-oxide enabled us to synthesize a wide variety of dibenzothiophene derivatives (Fig. 4A). For instance, *S*-oxidation of **4a** with *m*CPBA afforded dibenzothiophene *S*,*S*-dioxide **10a** in high yield.¹⁴ Rhodiumcatalyzed imidation of **4a** with trifluoroacetamide took place efficiently in the presence of iodobenzene diacetate and magnesium oxide to provide sulfoximine **11**.¹⁵ Reduction of sulfoxide **4a** with TFAA and sodium iodide proceeded smoothly to furnish dibenzothiophene **9a** in high yield.¹⁶ We also accomplished the preparation of thiophene- and furan-fused benzothiophenes **9b** and **9c** by the reduction of sulfoxides **4i** and **4j**. To our surprise, we found that the decomposition of furan-fused benzothiophene **9c** took place gradually in CDCl₃ at room temperature.¹⁷

We clarified that it is not easy to achieve the efficient synthesis of sulfoxides **4i** and **4j** by the conventional oxidation with *m*CPBA (Fig. 4B and C).⁵ Indeed, the oxidation of thienobenzothiophene **9b** with *m*CPBA (0.7 or 1.4 equiv.) afforded a



Fig. 3 (A) Stepwise synthesis of **4a** *via* isolation of **7**. (B) Plausible mechanism for the synthesis of **4a** from **3a**. (C) Electrophilic cyclization of sulfoxide **8**.



Fig. 4 (A) Transformations of 4a, 4i, and 4j. (B) Oxidation of 9b. (C) Oxidation of 9c.



Fig. 5 (A) C-H propargylation using **4a**. (B) Synthesis and transformations of o-silylaryl triflate **4r**.

mixture of sulfoxide **4i** and sulfone **10b**, clearly showing the problematic reactivity of sulfoxides under the oxidation conditions leading to sulfones (Fig. 4B). Furthermore, *m*CPBA oxidation of furanobenzothiophene **4j** provided a complex mixture of products (Fig. 4C). These results obviously indicate an advantage of the newly developed method to synthesize dibenzothiophene *S*-oxides.

Further transformations of dibenzothiophene S-oxides realized a wide variety of highly functionalized dibenzothiophene derivatives (Fig. 5). An extended Pummerer-type propargylation of dibenzothiophene S-oxide 4a with silane 12 took place efficiently to provide 13 and 13' in high combined yield, where C-C formation at the electron-rich but bulky site occurred smoothly to furnish 13 as a major product (Fig. 5A).^{4a} It is worth noting that the good functional group tolerance allowed us to prepare o-silylaryl triflate 4r efficiently by the coupling-cyclization approach without damaging reactive silvl and triflyloxy groups (Fig. 5B). This result shows a clear advantage over other dibenzothiophene S-oxide synthesis with nucleophilic carbanions or heating under basic conditions.⁶ Then, regioselective N-arylation of aniline with aryne IV took place smoothly to afford adduct 14 in moderate yield without forming the regioisomer, in which the sulfoxide moiety remained unreacted.¹⁸⁻²⁰ Moreover, we accomplished the synthesis of triazole-fused dibenzothiophene S-oxide 15 by regioselective cycloaddition of aryne IV with benzyl azide. Since o-silvlaryl triflate moiety serves to synthesize highly substituted arenes through aryne intermediates by a wide variety of transformations, a broad range of dibenzothiophene derivatives will be prepared by diverse aryne reactions.

In summary, we have developed an efficient method to prepare dibenzothiophene *S*-oxides from 2-bromoaryl sulfinate esters and arylboronic acids in 2 steps. Diverse dibenzothiophene *S*-oxides having various functionalities were synthesized by Br-selective cross-coupling and subsequent cyclization with Tf₂O without side reactions through the activation of sulfoxides. The good transformability

of dibenzothiophene *S*-oxides allowed us to access novel heterocyclic molecules such as highly ring-fused thiophenes. Further studies including applications for the construction of new organosulfur skeletons by transformations of dibenzothiophene *S*-oxides are ongoing in our laboratory.

The authors thank Central Glass Co., Ltd. for providing Tf_2O . This work was supported by JSPS KAKENHI Grant Number JP22H02086 (S. Y.); The Uehara Memorial Foundation (S. Y.); Tokuyama Science Foundation (S. Y.); The Ube Foundation (S. Y.); and Inamori Research Grants (S. Y.)

Conflicts of interest

There are no conflicts to declare.

Notes and references

- 1 M. Zhang, G. E. Ravilious, L. M. Hicks, J. M. Jez and R. D. McCulla, J. Am. Chem. Soc., 2012, 134, 16979.
- 2 G. C. M. Smith, A. J. Slade, C. J. Richardson and B. W. Durkacz, WO2006/126010 A2, 2006.
- 3 H. Wang, H. Wang, Z. Wang, L. Tang, G. Zeng, P. Xu, M. Chen, T. Xiong, C. Zhou, X. Li, D. Huang, Y. Zhu, Z. Wang and J. Tang, *Chem. Soc. Rev.*, 2020, **49**, 4135.
- 4 (a) J. A. Fernańdez-Salas, A. J. Eberhart and D. J. Procter, J. Am. Chem. Soc., 2016, **138**, 790; (b) X. Wang, X. Xun, H. Song, Y. Liu and Q. Wang, Org. Lett., 2022, **24**, 4580; (c) X. Li, C. Golz and M. Alcarazo, Angew. Chem., Int. Ed., 2021, **60**, 6943.
- 5 (a) D. Villemin and X. Vlieghe, *Sulfur Lett.*, 1998, 21, 199; (b) M. H. Aukland, M. Šiaučiulis, A. West, G. J. P. Perry and D. J. Procter, *Nat. Catal.*, 2020, 3, 163.
- 6 For other dibenzothiophene S-oxide synthesis, see: (a) T. Wesch,
 A. Berthelot-Bréhier, F. R. Leroux and F. Colobert, Org. Lett., 2013,
 15, 2490; (b) R. Wada, S. Kaga, Y. Kawai, K. Futamura, T. Murai and
 F. Shibahara, Tetrahedron, 2021, 83, 131978.
- 7 (a) A. Kobayashi, T. Matsuzawa, T. Hosoya and S. Yoshida, *Chem. Commun.*, 2020, **56**, 5429; (b) A. Kobayashi, T. Matsuzawa, T. Hosoya and S. Yoshida, *Chem. Lett.*, 2020, **49**, 813.
- 8 For arylation of sulfinate esters using AlCl₃, see: F. Yuste, A. H. Linares, V. M. Mastranzo, B. Ortiz, R. Sánchez-Obregón, A. Fraile and J. L. G. Ruano, *J. Org. Chem.*, 2011, **76**, 4635. In this report, it is not easy to perform *ortho*-sulfinylation of arenes.
- 9 K. Nakamura, Y. Kumagai, A. Kobayashi and S. Yoshida, *Org. Biomol. Chem.*, 2023, **21**, 6886.
- 10 M. Suzuki, K. Kanemoto, Y. Nakamura, T. Hosoya and S. Yoshida, Org. Lett., 2021, 23, 3793.
- 11 When the reactions were performed in CH_2Cl_2 at 0 °C, dibenzothiophene S-oxides **4d**, **4i**, **4j**, **4k**, and **4n** were obtained in 28, 76, 25, 0, and 0% yield, respectively.
- 12 A. Isor, B. V. Chartier, M. Abo, E. R. Currens, E. Weerapana and R. D. McCulla, *RSC Chem. Biol.*, 2021, 2, 577.
- 13 D. Vasu, J. N. Hausmann, H. Saito, T. Yanagi, H. Yorimitsu and A. Osuka, Asian J. Org. Chem., 2017, 6, 1390.
- 14 H. Hussain, A. Al-Harrasi, V. R. Green, I. Ahmed, G. Abbas and N. U. Rehman, RSC Adv., 2014, 4, 12882.
- 15 H. Okamura and C. Bolm, Org. Lett., 2004, 6, 1305.
- 16 M. de Greef and S. Z. Zard, Tetrahedron, 2004, 60, 7781.
- 17 When a time-course monitoring experiment of furan-fused benzothiophene 9c in $CDCl_3$ at room temperature was conducted, 85% of 9c was decomposed after 7 days.
- 18 Y. Nakamura, Y. Miyata, K. Uchida, S. Yoshida and T. Hosoya, Org. Lett., 2019, 21, 5252.
- (a) T. Matsuzawa, K. Uchida, S. Yoshida and T. Hosoya, Org. Lett., 2017, **19**, 5521; (b) S. Yoshida, H. Nakajima, K. Uchida, T. Yano, M. Kondo, T. Matsushita and T. Hosoya, Chem. Lett., 2017, **46**, 77.
- 20 S. Yoshida, T. Kuribara, T. Morita, T. Matsuzawa, K. Morimoto, T. Kobayashi and T. Hosoya, RSC Adv., 2018, 8, 21754.