



Cite this: *Chem. Commun.*, 2024, 60, 3154

Received 26th December 2023,
Accepted 15th February 2024

DOI: 10.1039/d3cc06249j

rsc.li/chemcomm

Diastereoselective access to C,C-glycosyl amino acids via iron-catalyzed, auxiliary-enabled MHAT coupling†

Mylène Lang,[†] Damien Tardieu, Benoit Pousse, Philippe Compain[†] * and Nicolas Kern[†] *

Access to C,C-glycosyl amino acids as a novel class of glycomimetics is reported by means of radical generation, intermolecular addition and stereoselective reduction via a metal-induced hydrogen atom transfer (MHAT) sequence. The 'matched' coupling of *exo*-D-glycals with an enantiopure dehydroalanine bearing a (*R*)-configured benzyl oxazolidinone enables a singular case of two-fold diastereocontrol under iron catalysis. In the common *exo*-D-glucal series, the nature of the C-2 substituent was found to play a key role from both reactivity and stereocontrol aspects.

Glycoproteins, as ubiquitous components of mammal tissues, are involved in multiple biological processes such as cell signaling, growth regulation and immunity.^{1,2} In the realm of pharmaceutical research, linking an amino acid chain to sugar(s) also stands as an established way to improve stability towards proteases and increase membrane permeability.³ As the overwhelming majority of natural glycopeptides comprise enzymatically labile C–O or C–N linkages (with C-mannosyl tryptophan as the sole exception⁴), unnatural C-glycopeptides both furnish useful probes to facilitate the elucidation of essential processes *in vivo*, as well as stable drug or vaccine candidates.⁵ This promising potential has therefore aroused intense synthetic efforts to construct C-linked sugar amino acid (SAA) building blocks.^{6,7} Formal "deletion" of the linking O- or N-atom ideally calls for the direct coupling of a pyranosyl unit activated at C-1⁸ with a functionalized two-carbon synthon bearing the (masked) AA motif. Such step-economic and convergent disconnection however adds challenges, especially regarding stereoselectivity and functional orthogonality. Notably, Ackermann and Li & Liu independently demonstrated the virtue of Pd-catalyzed C(sp²)-H and C(sp³)-H activation to

assemble a persilylated iodoglycal and auxiliary-tagged enantiopure amino acid (Fig. 1a).⁹ While excellent yields and diastereoselectivities (β to nitrogen, *i.e.* at C-1') could be achieved, efficient introduction of key O- or N-functionality at the C-2 site in such adducts can be nontrivial. Another mild approach involves the generation of glycosyl radicals¹⁰ and trapping with dehydroalanine-type acceptors (DHAs). Very recently, two photoinduced variants emanated from the groups of Di




Fig. 1 Recent convergent strategies to assemble C-glycosyl amino acids from pyranosides and (C₂) amino acid synthons (a) and (b) and aims of this work (c).

Laboratoire d'Innovation Moléculaire et Applications (LIMA), UMR 7042, Université de Strasbourg/Université de Haute-Alsace/CNRS, ECPM, 25 rue Becquerel, 67087, Strasbourg, France. E-mail: philippe.compain@unistra.fr, nicolas.kern@unistra.fr
† Electronic supplementary information (ESI) available: Experimental details including synthesis and characterization. See DOI: <https://doi.org/10.1039/d3cc06249j>




Carmine using pyranosyl bromides¹¹ and those from Xie, Walczak & Zhu who relied on furanosyl and pyranosyl dihydropyridyl esters (Fig. 1b).¹² Glycosyl radical additions to achiral and/or peptide-linked DHA derivatives typically proceed with axial selectivity but low control at the AA centre (α to nitrogen, *i.e.* at C-2'), emphasizing the poor level of induction imparted from the sugar and AA moieties. This trend, observed early by Kessler,¹³ appears as a longstanding issue. Therefore, Xie, Walczak & Zhu employed the (*S*)-enantiomer of the Karady-Beckwith acceptor and achieved high control in most cases through *cis*-selective reduction by hydrogen atom transfer (HAT), as documented.¹⁴

In connection with our search for *neo*-glycoconjugates and interest in novel transformations of *exo*-glycals,¹⁵ we questioned the viability of an MHAT (metal-induced HAT)^{16,17} coupling to form *C,C*-glycoaminoacids, unusual adducts embedding an atypical fully substituted pseudoanomeric carbon.¹⁸ Several intramolecular cyclization strategies were examined, all of which met with failure due to the low stability of DHA-acylated *exo*-glycals. While an intermolecular approach first appeared daunting in terms of two-fold stereocontrol (yet unreported for Fe-HAT Baran-Giese type coupling, and including an exocyclic centre), we were curious whether the inherent low stereoinduction inferred from the pyranoside (*vide supra*) could be enforced. We envisioned that such a task may be achievable through the identification of an appropriate “matched” nonracemic radical acceptor, along with careful choice of the protection patterns – especially for the C-2 hydroxyl moiety, in the vicinity of subsequently formed open-shell species (Fig. 1c).

To begin to answer this question, *O*-benzyl-*N*-phthaloyl DHA-based acceptor **2a** was selected (Table 1). Screening of coupling conditions (catalytic Fe(acac)₃, phenylsilane and Na₂HPO₄ buffer in EtOH) applied to a mixture of perbenzylated *exo*-glycal **1a** (1 equiv.) and **2a** (2 equiv.) first suggested that productive radical generation from *exo*-glucals calls for specific dilute conditions and moderate catalyst loading (entries 1–4). Along with targeted adduct **3a**, alanine imidoester **4a** was identified as the main side product. Competitive reduction of *C*-radicophiles has been scarcely mentioned in the Fe-HAT literature,¹⁹ and may be favored here considering the captodative character of the product radical,²⁰ partly overriding the usual cross-coupling chemoselectivity attributed to polar effects. This may also explain the use of excess electron-rich alkenes in related protocols,^{19,21} inconvenient with valuable *exo*-glycals. However, formal hydrogenation of **1a** and telomerization¹³ were fully avoided.²² Since chromatographical separation of **3a** and **4a** proved tedious, base-mediated hydrolysis of the benzyl ester resulted in the epimerization of the neoformed AA stereocentre. Substitution of **2a** for PMB ester **2b** (entry 5, conditions [A]) allowed quantitative and epimerization-free acid-mediated dealkylation with TFA in conjunction with 1,3-dimethoxybenzene (DMB), yielding free acid **5**.²³ Akin to **2a**, acceptor **2b** appeared almost insoluble at 25 °C and sparingly at 60 °C; therefore inclusion of cosolvents was evaluated. Use of minimal EtOH in THF (16 equiv. to reach full

Table 1 Optimization of the intermolecular diastereoselective catalytic MHAT coupling



Entry	Acceptor	Solvent(s)/conc. (M)	<i>t</i> (h)	Yield (%) / product	d.r. ^b
1	2a	EtOH (0.1)	1.5	60 (3a) ^a	69:31
2	2a	EtOH (0.01)	16	52 (3a) ^a	71:29
3	2a	EtOH (0.03)	4	77 (3a) ^a	73:27
4 ^c	2a	EtOH (0.03)	2	40 (3a) ^{ad}	nd
5 [A]	2b	EtOH (0.03)	4	76 (5) ^e	69:31
6 [B]	2b	THF (0.03)	16	66 (5) ^e	71:29
7	2c	EtOH (0.03)	4	50 (5) ^e	58:42
8	2d	EtOH (0.03)	4	Trace ^d	nd
9	2e	EtOH (0.03)	4	21 (5) ^e	70:30
10 [A]	2f	EtOH (0.03)	4	75 (5) ^e	88:12 ^g
11 [B]	2f	THF (0.03)	16	82	92:8 ^g
12 ^h	2f	THF (0.03)	16	45	96:4 ^g
13	2g	EtOH (0.03)	4	67	74:26
14	2h	EtOH (0.03)	4	31	93:7 ^g

^a Estimated yield by ¹H NMR (isolated material contaminated with **4a**).

^b Determined by ¹H NMR analysis of partly purified **3a** or pure **5**. ^c 20 mol% [Fe]. ^d Formation of unseparable side products. ^e Isolated yield after acidic hydrolysis to **5**. ^f 16 equiv. EtOH as additive. ^g Confirmed by HPLC. ^h At 25 °C. [A] & [B]: optimal conditions.

conversion; entry 6, conditions [B]) resulted in close efficiency at the cost of reaction rate, which may be rationalized from the involvement of the alcohol in the rate-determining iron hydride generation step.¹⁵ Overall, all experiments involving achiral acceptor **2b** furnished products **3** (or **5**) in moderate d.r. (close to 7:3). We thus moved forward by evaluating DHA-tagged chiral oxazolidinones **2c–2h**.^{21,24} Unlike esters, imide coupling adducts could be carefully hydrolyzed to **5** without epimerization using minimal LiOH and H₂O₂, enabling comparison of the stereochemical course with reactions of **2b**. First, (*R*)-configured isopropyl derivative **2c** underwent addition with lower diastereoselectivity (entry 7). Phenyl analog **2d** only led to traces of the coupling adduct, while *isobutyryl* analog **2e** reacted poorly with the glycosyl radical without impacting stereoinduction (entries 8 and 9). Gratifyingly, a ‘matched’ case was disclosed with benzyl surrogate **2f** (*i.e.* Evans’ auxiliary,²⁵ entry 10), which reacted efficiently and induced good diastereoselectivity (88:12) under conditions [A]. As the stereodetermining redox termination step involves the protic solvent,²⁶ system [B] with minimal EtOH was evaluated and induced a slight enhancement (entry 11). Since acceptor **2f** displayed better solubility in THF, decreasing the reaction temperature to 25 °C allowed even higher control but with lower efficiency (entry 12). Consecutively, the (*S*)-enantiomer **2g** was probed to be ‘mismatched’ (entry 13). The *gem*-dimethyl surrogate of the optimal auxiliary²⁷ acted as a potent inductor, but the apparent






Fig. 2 (a) Identification of the optimal *exo*-glucal protection pattern and derivatization. (b) Application to mannosyl and galactosyl surrogates. (c) Application to furanoside. (d) Main limitations observed through this study. ^a Reaction time: 3–4 h. ^b Reaction time: 12–16 h (full conv obs). ^c Isolated yields. ^d dr determined by ¹H NMR analysis of isolated products due to overlaps.

lower stability of parent acceptor **2h** only enabled moderate yields (entry 14). The observation that all auxiliaries and conditions afforded the same major diastereoisomer of **5** stands in line with previous reports involving pyranosides, where the degree but not the direction of stereoselectivity could be influenced.^{13,14} Furthermore, closer examination of related work¹² reveals that stereinduction in the termination step is usually lower for pyranosides bearing equatorial O- or N-functionality at C-2. In contrast to mannosides or 2-deoxy derivatives for example, their orientation may partly impede the population of a favorable conformer for stereoselective termination (*vide infra*). A series of *exo*-D-glucals diversely protected at C-2 were thus examined (Fig. 2a).[‡] With relatively bulky silyl groups (**1b** & **1c**) negligible conversions were observed while DHA reduction predominated. The solvolitically unstable free-hydroxyl derivative **1d** only led to the corresponding adduct **6** in low yield and d.r. Pleasingly, stable and less hindered methoxymethyl (MOM) derivative **1e** led to the highest yield and stereoselectivity (94 : 6) observed. Orthogonal removal of the auxiliary in **7** and acid-mediated lactonization to **8** were straightforward and allowed unambiguous assignment of the (*R*)-configuration of the AA by ¹H NOESY analysis. In view of the importance of 2-amino-*C*-glycosides,²⁸ coupling of the seldom (2-acetamido)*exo*-glycal²⁹ **1f** could be achieved in moderate yield and stereoselectivity, granting with adduct **9** a first entry to 2-amido-*C,C*-glycosidic motifs. Considering these

results, we speculate that (1) steric impediment of the olefin in **1** is detrimental to radical generation; (2) alongside competitive processes efficiency is also correlated to the chemical stability of the often sensitive *exo*-glycal; and (3) high stereoselectivities are obtained when the C-2 position of the sugar is relatively unhindered. As mentioned above, if the optimized acceptor and conditions enable enforcement of two-fold stereocontrol in the D-glucose series, extension to related important pyranosides should be feasible. This was probed with *exo*-D-mannal **1g** and *exo*-D-galactal **1h**, whose corresponding coupling adducts **10** and **11** were successfully isolated after direct hydrolysis (Fig. 2b). As notable, selective MOM protection at C-2 was not mandatory here to achieve high control. The (very) decent chemical yields obtained from these fragile, electron-rich terminal *exo*-glycals again testify of the relative mildness of the Fe-HAT conditions. Although optimized for pyranosides, the protocol also allowed the efficient conversion of furanoside **1i** into adduct **12** with significant control at the AA centre (Fig. 2c). The main limitations concern trisubstituted¹⁵ and electronically deactivated difluoroolefins,³⁰ which do not undergo MHAT at a sufficient rate with respect to DHA derivatives (Fig. 2d). On the other hand, sulfonylhydrazone acceptors used for reductive alkylation³¹ either seem to: (1) lack sufficient reactivity towards tertiary glycosyl radicals (alkylsulfonyl derivatives), whose reduction thus leads to formal alkene hydrogenation, or (2) undergo nonselective fragmentation (arylsulfonyl derivatives).

Based on our results and taking account of Poli & Holland's studies,²⁶ a tentative model for the stereocontrolled formation of the AA centre may be envisioned from two perspectives (Fig. 3). Following generation of an iron(III) enolate *via* inner-sphere electron transfer from Fe(II)(acac)₂ species (a), a non-chelated model – expectable considering a moderately Lewis acidic and coordinatively saturated Fe(III) complex – may explain the preference for *si*-face 'concerted' protonation from a metal-bound EtOH molecule leading to the (*R*)-configuration while accounting for the effect of substituents at C-2, in close vicinity of the aryl moiety presented by the auxiliary. Additional shielding of the *re*-face is imposed by the angular methyl group, while the flat phthalimide can partly rotate to minimize steric interactions with the former methyl and the auxiliary, also restricting the rotation of the pseudoanomeric C–C bond. Similar considerations may apply to a proton-coupled electron transfer process (b), in which a conjugated π -type radical in its




Fig. 3 Proposed models for stereoselective termination *via* SET/protonation (a) or proton-coupled electron transfer (b) in the D-glucose series.



ground state would adopt a similar planar geometry, poised to undergo stereoselective PCET from alcohol-bound Fe(II) species.

In conclusion, a novel entry to rare *C,C*-glycosyl amino acids was established. Guided by a systematic study of substituents and auxiliary effects, high two-fold stereocontrol was achieved in the direct assembly of *exo*-glycals with dehydroalanine acceptors, representing a first case using the MHAT intermolecular coupling pathway. Further development of new catalytic stereocontrolled reactions of *exo*-glycals is underway and will be reported in due course.

NK gratefully acknowledges support from Unistra (IDEX grant W19RAT27). ML and DT thank the French Ministry of Research (MRT fellowships). Dr E. Wasielewski and M. Chessé are thanked for their support. Dr X. Bugaut is thanked for insightful discussions.

Conflicts of interest

There are no conflicts to declare.

Notes and references

‡ Due to the varying polarities of adducts, from this stage the auxiliary was cleaved only if required for an easier purification.

- 1 E. Saxon and C. R. Bertozzi, *Science*, 2001, **291**, 2357; R. A. Dwek, *Chem. Rev.*, 1996, **96**, 683.
- 2 K. T. Schjoldager, Y. Narimatsu, H. J. Joshi and H. Clausen, *Nat. Rev. Mol. Cell Biol.*, 2020, **21**, 729; G. W. Hart, *J. Biol. Chem.*, 2019, **294**, 2211; A. Varki, *Glycobiology*, 2017, **27**, 3.
- 3 M. Muttenthaler, G. F. King, D. J. Adams and P. F. Alewood, *Nat. Rev. Drug Discovery*, 2021, **20**, 309; S. V. Moradi, W. M. Hussein, P. Varamini, P. Simerska and I. Toth, *Chem. Sci.*, 2016, **7**, 2492; C. R. Apostol, M. Hay and R. Polt, *Peptides*, 2020, **131**, 170369.
- 4 S. Minakata, S. Manabe, Y. Inai, M. Ikezaki, K. Nishitsuji, Y. Ito and Y. Ihara, *Molecules*, 2021, **26**, 5258.
- 5 D. P. Gamblin, E. M. Scanlan and B. G. Davis, *Chem. Rev.*, 2009, **109**, 131; S. A. W. Gruner, E. Locardi, E. Lohof and H. Kessler, *Chem. Rev.*, 2002, **102**, 491; T. P. Hogervorst, R. J. Eveline Li, L. Marino, S. C. M. Bruijns, N. J. Meeuwenoord, D. M. Filippov, H. S. Overkleeft, G. A. van der Marel, S. J. van Vliet, Y. van Kooyk and J. D. C. Codée, *ACS Chem. Biol.*, 2020, **15**, 728; U. Westerlind, *Beilstein J. Org. Chem.*, 2012, **8**, 804.
- 6 For reviews see: Z. Yuan, C. Zhu, Y. Zhang and Y. Rao, *Eur. J. Org. Chem.*, 2022, e202201036; A. Dondoni and A. Marra, *Chem. Rev.*, 2000, **100**, 4395; H. Herzner, T. Reipen, M. Schultz and A. Kunz, *Chem. Rev.*, 2000, **100**, 4495.
- 7 For selected recent as well as classical approaches see: Y.-H. Liu, Y.-N. Xia, T. Gulzar, B. Wei, H. Li, D. Zhu, Z. Hu, P. Xu and B. Yu, *Nat. Commun.*, 2021, **12**, 4924; Y. Wei, Q. Wang and M. J. Koo, *Angew. Chem., Int. Ed.*, 2023, **62**, e202214247; Y.-N. Ding, N. Li, Y.-C. Huang, Y. An and Y.-M. Liang, *Org. Lett.*, 2022, **24**, 4519; R. Mao, S. Xi, S. Shah, M. J. Roy, A. John, J. P. Lingford, G. Gäde, N. E. Scott and E. D. Goddard-Borger, *J. Am. Chem. Soc.*, 2021, **143**, 12699; P. Ji, Y. Zhang, Y. Wei, H. Huang, W. Hu, P. A. Mariano and W. Wang, *Org. Lett.*, 2019, **21**, 3086; C. Palomo, M. Oiarbide, A. Landa, M. C. González-Rego, J. M. García, A. González, J. M. Odriozola, M. Martín-Pastor and A. Linden, *J. Am. Chem. Soc.*, 2002, **124**, 8637; A. Nuzzi, A. Massi and A. Dondoni, *Org. Lett.*, 2008, **10**, 4485; G. J. McGarvey, T. E. Benedum and F. W. Schmidtmann, *Org. Lett.*, 2002, **21**, 3591; D. E. Paterson, F. K. Griffin, M.-L. Alcaraz and R. J. K. Taylor, *Eur. J. Org. Chem.*, 2002, 1323.
- 8 For complementary methods of anomeric C–C bond formation in C-diglycoside synthesis see: D. C. Koester, E. Kriemen and D. B. Werz, *Angew. Chem., Int. Ed.*, 2013, **52**, 2985; G. Hirai, T. Watanabe, K. Yamaguchi, T. Miyagi and M. Sodeoka, *J. Am. Chem. Soc.*, 2007, **129**, 15420; S. Palmier, B. Vauzeilles and J.-M. Beau, *Org. Biomol. Chem.*, 2003, **1**, 1097.
- 9 J. Wu, N. Kaplaneris, S. Ni, F. Kaltenhäuser and L. Ackermann, *Chem. Sci.*, 2020, **11**, 6521; Y. Liu, Y. Wang, W. Dai, W. Huang, Y. Li and H. Liu, *Angew. Chem., Int. Ed.*, 2020, **59**, 3491.
- 10 C. Wang, R. Qi and Z. Xu, *Chem. – Eur. J.*, 2023, **29**, e202203689; for reviews on radical glycosylation see: Y. Yang and B. Yu, *Chem. Rev.*, 2017, **117**, 12281; L.-Y. Xu, N.-L. Fan and X.-G. Hu, *Org. Biomol. Chem.*, 2020, **18**, 5095; A. Chen, L. Xu, Z. Zhou, S. Zhao, T. Yang and F. Zhu, *J. Carbohydr. Chem.*, 2021, **40**, 361.
- 11 L. Poletti, A. Massi, D. Ragno, F. Droghetti, M. Natali, C. De Risi, O. Bortolini and G. Di Carmine, *Org. Lett.*, 2023, **25**, 4862; for seminal work see: R. S. Andrews, J. J. Becker and M. R. Gagné, *Angew. Chem., Int. Ed.*, 2010, **49**, 7274.
- 12 A. Chen, S. Zhao, Y. Han, Z. Zhou, B. Yang, L.-G. Xie, M. A. Walczak and F. Zhu, *Chem. Sci.*, 2023, **14**, 7569.
- 13 H. Kessler, V. Wittmann, M. Köck and M. Kottenhahn, *Angew. Chem., Int. Ed. Engl.*, 1992, **31**, 902.
- 14 G. A. Adamson, A. L. J. Beckwith and C. L. L. Chai, *Aust. J. Chem.*, 2004, **57**, 629; J. R. Axon and A. L. J. Beckwith, *J. Chem. Soc., Chem. Commun.*, 1995, 549.
- 15 D. Tardieu, M. Desnoyers, C. Laye, D. Hazelard, N. Kern and P. Compain, *Org. Lett.*, 2019, **21**, 7262; H. Liu, A. Laporte, D. Tardieu, D. Hazelard and P. Compain, *J. Org. Chem.*, 2022, **87**, 13178.
- 16 J. C. Lo, D. Kim, C.-M. Pan, J. T. Edwards, Y. Yabe, J. Gui, T. Qin, S. Gutiérrez, J. Giacoboni, M. W. Smith, P. L. Holland and P. S. Baran, *J. Am. Chem. Soc.*, 2017, **139**, 2484; J. C. Lo, J. Gui, Y. Yabe, C.-M. Pan and P. S. Baran, *Nature*, 2014, **516**, 343.
- 17 J. L. M. Matos, S. A. Green and R. A. Shenvi, *Org. React.*, 2019, **100**, 383; S. W. M. Crossley, C. Obradors, R. M. Martinez and R. A. Shenvi, *Chem. Rev.*, 2016, **116**, 8912; S. A. Green, S. W. M. Crossley, J. L. M. Matos, S. Vásquez-Céspedes, S. L. Shevick and R. A. Shenvi, *Acc. Chem. Res.*, 2018, **51**, 2628; J. Wu and Z. Ma, *Org. Chem. Front.*, 2021, **8**, 7050.
- 18 M. Richard, A.-S. Felten, C. Didierjean, M. Ruiz-Lopez, Y. Chapleur and N. Pellegrini-Moïse, *Eur. J. Org. Chem.*, 2014, 7364; F. Schweizer and O. Hindsgaul, *Carbohydr. Res.*, 2006, **341**, 1730; S. Bera, B. Chatterjee and D. Mondal, *RSC Adv.*, 2016, **6**, 77212.
- 19 V. A. Larionov, N. V. Stoletova, V. I. Kovalev, A. F. Smol'yakov, T. F. Savel'yeva and V. I. Maleev, *Org. Chem. Front.*, 2019, **6**, 1094; B. Wu and R. Zhu, *ACS Catal.*, 2020, **10**, 510.
- 20 While additions forming captodative radicals witness increased exothermicity and lowered activation barriers, electron affinity is no more correlated to reaction enthalpy and thus polar stabilization decreases. See: R. Arnaud, N. Bugaut, V. Vetere and V. Barone, *J. Am. Chem. Soc.*, 1998, **120**, 5733.
- 21 H. Zhang, H. Li, H. Yang and H. Fu, *Org. Lett.*, 2016, **18**, 3362.
- 22 Presence of excess acceptor alongside the nontrivial stabilization conferred by the imide makes this noteworthy, see: S. Han and S. Z. Zard, *Org. Lett.*, 2014, **20**, 5386; Lowering the **1a**:**2** ratio to 1:1.5 led to partial conversion of **1a**, while increasing it to 1:3 resulted in lower yields.
- 23 M. E. Jung and P. Koch, *Tetrahedron Lett.*, 2011, **52**, 6051.
- 24 M. P. Sibi and J. Ji, *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 190.
- 25 J. R. Gage and D. A. Evans, *Org. Synth.*, 1990, **68**, 77.
- 26 D. Kim, S. M. W. Rahaman, B. Q. Mercado, R. Poli and P. L. Holland, *J. Am. Chem. Soc.*, 2019, **141**, 7473.
- 27 T. Herrmann, L. L. Smith and A. Zakarian, *J. Am. Chem. Soc.*, 2012, **134**, 6976; M. Pérez-Palau, N. Sanosa, P. Romea, F. Urpí, R. López, E. Gómez-Bengoa and M. Font-Bardia, *Org. Lett.*, 2021, **23**, 8852.
- 28 C. Nativi, F. Papi and S. Roelens, *Chem. Commun.*, 2019, **55**, 7729; C. H. Röhrig, M. Takhi and R. R. Schmidt, *Synlett*, 2001, 1170; D. Urban, T. Skydstrup, C. Riche, A. Chiaroni and J.-M. Beau, *Chem. Commun.*, 1996, 1883.
- 29 Y. Leshch, A. Jacobsen, J. Thimm and J. Thiem, *Org. Lett.*, 2013, **15**, 4948.
- 30 T. F. Herpin, W. B. Motherwell, B. P. Roberts, S. Roland and J.-M. Weibel, *Tetrahedron*, 1997, **53**, 15085.
- 31 H. T. Dao, C. Li, Q. Michaudel, B. D. Maxwell and P. S. Baran, *J. Am. Chem. Soc.*, 2015, **137**, 8046; M. Saladrigas, J. Bonjoch and B. Bradshaw, *Org. Lett.*, 2020, **22**, 684.

