



# Strain-induced carbon–carbon bond cleavage of bowl-shaped sumanenone†‡

 Mikey Nishimoto,<sup>a</sup> Yuta Uetake,<sup>ib</sup> Yumi Yakiyama<sup>ib</sup> and Hidehiro Sakurai<sup>ib</sup>\*

 Cite this: *Chem. Commun.*, 2024, 60, 3982

 Received 2nd January 2024,  
 Accepted 8th March 2024

DOI: 10.1039/d4cc00008k


rsc.li/chemcomm

**This study details a highly effective ring-opening reaction that involves acid-mediated carbon–carbon bond cleavage of the bucky bowl, sumanenone. The reaction of the bowl-shaped sumanenone with AcOH and TfOH results in the formation of a planar carboxylic acid. The examination of reactivity in comparison to planar analogues, along with theoretical calculations, suggests that the release of curved strain is a crucial factor for the success of this reaction.**

The inherent strain in a molecule has a significant influence on its reactivity. As a representative example, Bertozzi *et al.* reported that the azide–alkyne cycloaddition reaction proceeds without a copper catalyst using strained alkynes.<sup>1</sup> Curved  $\pi$ -conjugated molecules such as fullerenes<sup>2,3</sup> and buckybowls<sup>4–6</sup> are also regarded as destabilized substrates in comparison with planar  $\pi$ -conjugated molecules; therefore, they exhibit unique reactivities (Fig. 1a). Previous reports demonstrated that addition reactions involving dearomatization, which are challenging to initiate in planar molecules, successfully occur in the context of curved  $\pi$ -conjugated molecules.<sup>7–13</sup> For instance, the ring-opening of fullerenes arising from cycloaddition reactions has been employed to synthesize azafullerene dimer (C<sub>59</sub>N)<sub>2</sub>,<sup>14</sup> and diverse endohedral fullerenes by “molecular surgery” methods.<sup>15–17</sup> Although reports are more limited than for addition reactions, C–C bond cleavage reactions are also initiated by curved strain in several buckybowls (Scheme S1, ESI†). Shionoya *et al.* demonstrated an iridium-catalyzed C–C bond cleavage of corannulene bearing a 2-pyridyl group as a directing group.<sup>18</sup> Shao *et al.* observed cleavage at the peripheral C–C double bond of trithiasumanene with electron-donating substituents, achieved through

Oxone<sup>®</sup> oxidation, leading to the synthesis of planar  $\pi$ -conjugated molecules. These molecules were then employed to construct [5-6-7] fused poly-heterocycles, exhibiting unique photo-physical properties derived from a donor–acceptor system.<sup>19</sup> Consequently, the progression of reactions harnessing curved strain holds the potential to offer a pathway to access unique molecular skeletons that pose challenges in synthesis from planar molecules.

Sumanene,<sup>20,21</sup> a C<sub>60</sub> fragment bucky bowl as well as corannulene, possesses aromatic rings and reactive benzyl positions on its periphery. Investigations into both peripheral and internal functionalisation have been conducted. The sumanene bowl structure is more distorted than that of corannulene. Our group has reported osmylation to aromatic ring in sumanene<sup>13</sup> and the mild C–C bond cleavage of 2-hydroxyphenyl sumanenetrione.<sup>22</sup> In the latter case, the presence of the 2-hydroxyphenyl group is essential for the reaction to proceed.



**Fig. 1** (a) Energy diagram about the destabilization of the substrate by strain. (b) Strain-induced C–C bond cleavage of sumanenone (1).

<sup>a</sup> Division of Applied Chemistry, Graduate School of Engineering, Osaka University, 2-1 Yamadaoka, Suita, Osaka 565-0871, Japan.

E-mail: hsakurai@chem.eng.osaka-u.ac.jp

<sup>b</sup> Innovative Catalysis Science Division, Institute for Open and Transdisciplinary Research Initiatives (ICS-OTRI), Osaka University, 2-1 Yamadaoka, Suita, Osaka 565-0871, Japan

† Dedicated to Professor Koichi Narasaka in celebration of his 80th birthday.

‡ Electronic supplementary information (ESI) available. CCDC 2323143. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d4cc00008k>




Herein, we report the Brønsted acid-mediated five-membered ring-opening of sumanenone (**1**) *via* C–C bond cleavage (Fig. 1b). Experimental and theoretical approaches were employed to investigate the origin of the difference in reactivity between a bowl-shaped molecule and a planar compound having a similar  $\pi$ -conjugation. In contrast to prior examples, this reaction advances without the need for directing groups or electron-donating substituents, clearly demonstrating the influence of inherent strain in the molecule on reactivity.

We have previously reported that Baeyer–Villiger oxidation of sumanenone (**1**) was accelerated by the addition of Brønsted acids.<sup>23</sup> In the current study, we observed that the five-membered ring-opening reaction of **1** led to the formation of carboxylic acid (**2**) in the presence of trifluoromethanesulfonic acid (TfOH) (500 mol%) in acetic acid under heating (Scheme 1). The <sup>1</sup>H NMR spectrum of the crude mixture indicated the complete consumption of **1**, with only product peaks observed, suggesting the clean formation of **2**. After extraction and washing with *n*-hexane, the pure product was isolated in 91% yield. Notably, this reaction did not proceed without TfOH, and **1** was fully recovered, suggesting the essential role of a strong acid.

The thus-obtained ring-opened product **2** was characterized by high-resolution mass spectrometry (HRMS) and NMR spectroscopy. The HRMS displayed  $m/z = 296.0837$  ( $[M^+]$ ), indicating the formation of **2**. The DFT-optimised structure of **2** exhibited a planar structure due to the ring-opening process. The <sup>1</sup>H NMR spectra and their assignments for **1** and **2** are shown in Fig. 2, corroborated by DFT calculation (Table S1, ESI†). A comparison of the <sup>1</sup>H NMR spectra between **1** and **2** revealed an increase in signals in the aromatic region of **2**, accounting for seven protons (Fig. 2). The signal for H<sup>i</sup> in Fig. 2b appeared at a lower magnetic field than other aromatic protons, suggesting a deshielding effect from the adjacent carboxyl group. In the case of **1**, the proton signals at the benzyl position are observed as a doublet with the *J*-coupling constant of approx. 20 Hz, reflecting the geminal coupling between *endo*- and *exo*-protons due to the deep bowl structure.<sup>24</sup> In contrast, those of **2** appeared as a singlet peak (H<sup>c</sup>: 4.48 ppm, H<sup>f</sup>: 4.51 ppm) owing to the planarisation of the molecular framework. This planarised ring-opened framework was also supported by the single-crystal X-ray structure analysis by converting from **2** to the corresponding 4-bromophenyl ester (**3**) (Scheme 1 and Fig. S1, ESI†).

The ring-opening hydrolysis of fluorenone (**4**), a substructure of **1**, has been reported to occur under high-temperature



Scheme 1 Strain-induced ring-opening of **1**.





Fig. 2 <sup>1</sup>H NMR spectra of (a) **1** in CDCl<sub>3</sub> and (b) **2** in DMSO-*d*<sub>6</sub>.

conditions and with the addition of an excess amount of base.<sup>25</sup> Although ring-opening reactions of a particular fluorenone derivative have been reported to proceed with an excess amount of Lewis acid,<sup>26</sup> ring-opening reactions with simple Brønsted acids have not been reported. However, given our prior demonstration that the ring-opening reaction occurred at room temperature when employing a sumanenone derivative with significant strain, it is plausible that the acid-promoted ring-opening reaction also takes place influenced by steric or electronic effect. To test this hypothesis, we conducted the reaction under the same conditions using **4**, and triphenylene (**5**), another planar molecule with the same  $\pi$ -conjugated structure as **1**. However, in both cases, the ring-opening reaction did not proceed, and the starting materials were recovered (Scheme 2). Therefore, the curved distortion of the bowl skeleton in **1** should play a crucial role in facilitating this reaction.

Based on these results, we proposed a possible reaction mechanism for the acid-mediated ring opening reaction (Fig. 3). The carbonyl group of **1** undergoes protonation and reacts with acetic acid to form a corresponding hemiketal intermediate. The  $\alpha$ -carbon ( $sp^2$ ) at the quaternary carbon centre is protonated by the superacid TfOH, generating a cationic intermediate. A similar electrophilic addition of bromonium ions to the quaternary carbon on sumanene skeleton has been previously reported.<sup>27</sup> The cationic intermediate undergoes ring-opening involving the release of curved strain, followed by hydrolysis to give the target compound. In this proposed mechanism, the reaction is considered to proceed efficiently due to the irreversibility of the ring-opening step.

To understand the reaction mechanism and the impact of bowl structure on reactivity, the density functional theory (DFT) calculations were performed for **1**, **4**, and **5** (Fig. 4) at the  $\omega$ B97X-D/aug-cc-PVTZ/SMD(AcOH)// $\omega$ B97X-D/aug-cc-PVDZ/SMD(AcOH)



Scheme 2 Control experiments on the strain-induced ring-opening reaction.






Fig. 3 Possible reaction mechanism.

level of theory. There is no substantial difference in  $\Delta G^\ddagger$  in the formation of hemiketals **B** by nucleophilic addition of acetic acid at **1**, **4**, and **5** (**1**:  $\Delta G^\ddagger = 21.7 \text{ kcal mol}^{-1}$ ; **4**:  $\Delta G^\ddagger = 22.5 \text{ kcal mol}^{-1}$ ; **5**:  $\Delta G^\ddagger = 22.0 \text{ kcal mol}^{-1}$ ). In contrast, comparing the  $\Delta\Delta G$  values for the protonation step, all the compounds were significantly destabilized due to dearomatization. Nevertheless, the calculated  $\Delta\Delta G$  value for **1** was  $16.7 \text{ kcal mol}^{-1}$ , considerably lower than that of **4** and **5** ( $29.8 \text{ kcal mol}^{-1}$  for **4**,

$28.8 \text{ kcal mol}^{-1}$  for **5**). This can be attributed to the resonance effect and the hyperconjugation that exclusively benefits the carbocation intermediate **1-C** due to the presence of methylene bridges (Fig. S2, ESI $^\ddagger$ ). The maximum  $\Delta G^\ddagger$  for the reaction mechanism was determined to be  $+34.7 \text{ kcal mol}^{-1}$  for the ring-opening step (**TS2**), a value conducive to the reaction occurring at  $100 \text{ }^\circ\text{C}$ . In contrast, the corresponding values for **4** and **5** were evaluated to be  $+46.0$  and  $+46.6 \text{ kcal mol}^{-1}$ , respectively. This result confirms that the ring-opening reaction is unlikely to take place even at high temperatures in the case of **4** and **5**. For compound **1**, the  $\Delta G$  of the final product **1-D** was found to be a negative value ( $-9.4 \text{ kcal mol}^{-1}$ ), attributed to substantial stabilization by strain-release, rendering the overall reaction process exothermic. Meanwhile, those of **4** and **5** were  $+24.6$  and  $+24.1 \text{ kcal mol}^{-1}$ , respectively, indicating an endothermic nature and supporting the notion that the reaction does not proceed. In summary, these DFT calculations provide a good explanation of the validity of the reaction mechanism and are in good agreement with the experimental results. The key to this reaction is the capacity of compound **1** to stabilize the cationic intermediate using its molecular framework, mitigating the destabilizing impact of dearomatization caused by protonation. Furthermore, the substantial contribution to stabilization arising from the release of curved strain is considered crucial. Both factors are posited to play a pivotal role in facilitating the cleavage of a stable C–C bond.




Fig. 4 Energy profile of the acid-mediated ring-opening reaction of **1**, **4**, and **5**. The calculated Gibbs free energies ( $\Delta G$  and  $\Delta G^\ddagger$ , 298.15 K, 1 atm) are shown in  $\text{kcal mol}^{-1}$ . All calculations were performed at the  $\omega\text{B97X-D/aug-cc-PVTZ/SMD}(\text{AcOH})//\omega\text{B97X-D/aug-cc-PVDZ/SMD}(\text{AcOH})$  level of theory. To compare energies in all steps of the reaction, the energy of **A**, **B** and **TS1** is denoted by the sum of  $\Delta G$  of TfOH and **1-A**, **1-B**, and **TS1**, respectively.



In conclusion, we have demonstrated an effective ring-opening reaction involving the C–C bond cleavage of the sumanenone backbone. A comparative analysis of **1** and planar analogs *via* experimental and computational methods highlights the critical role of the release of curved strain. The findings illuminate the distinctive reactivity conferred by the strained structure of the curved  $\pi$ -conjugated system. Moreover, the resulting carboxylic acid holds promise as a precursor for constructing unique molecular skeletons with a bowl/planar  $\pi$ -conjugated configuration, such as heterobuckybowls.

This study was supported by a Grant-in-Aid for Transformative Research Areas “Science of 2.5 Dimensional Materials” (No. JP21H05233), “Hyper-Ordered Structures Sciences” (No. JP21H05563 and JP23H04112), and JSPS KAKENHI (JP19H00912, JP20H00400 and JP20K15279), and JST SPRING (JPMJSP2138). The theoretical calculations were performed at the Research Centre for Computational Science, Okazaki, Japan (Project: 22-IMSC068).

## Conflicts of interest

There are no conflicts to declare.

## Notes and references

- N. J. Agard, J. A. Prescher and C. R. Bertozzi, *J. Am. Chem. Soc.*, 2004, **126**, 15046–15047.
- H. W. Kroto, J. R. Heath, S. C. O'Brien, R. F. Curl and R. D. Smalley, *Nature*, 1985, **318**, 162–163.
- R. C. Haddon, *Science*, 1993, **261**, 1545–1550.
- Y.-T. Wu and J. S. Siegel, *Chem. Rev.*, 2006, **106**, 4843–4867.
- V. M. Tsefrikas and L. T. Scott, *Chem. Rev.*, 2006, **106**, 4868–4884.
- M. Saito, H. Shinokubo and H. Sakurai, *Mater. Chem. Front.*, 2018, **2**, 635–661.
- D. V. Preda and L. T. Scott, *Tetrahedron Lett.*, 2000, **41**, 9633–9637.
- A. V. Zabula, S. N. Spisak, A. S. Filatov, A. Y. Rogachev and M. A. Petrukhina, *Angew. Chem., Int. Ed.*, 2011, **50**, 2971–2974.
- M. Yanney, F. R. Fronczek and A. Sygula, *Org. Lett.*, 2012, **14**, 4942–4945.
- H. E. Bronstein and L. T. Scott, *J. Org. Chem.*, 2008, **73**, 88–93.
- J. A. Steckel and K. D. Jordan, *J. Phys. Chem. A*, 2002, **106**, 2572–2579.
- H. Y. Cho, R. B. M. Ansems and L. T. Scott, *Beilstein J. Org. Chem.*, 2014, **10**, 956–968.
- N. Ikuma, Y. Yoshida, Y. Yakiyama, N. Ngamsomprasert and H. Sakurai, *Chem. Lett.*, 2018, **47**, 736–739.
- J. C. Hummelen, B. Knight, J. Pavlovich, R. González and F. Wudl, *Science*, 1995, **269**, 1554–1556.
- K. Komatsu, M. Murata and Y. Murata, *Science*, 2005, **307**, 238–240.
- M. Murata, Y. Murata and K. Komatsu, *Chem. Commun.*, 2008, 6083–6094.
- K. Kurotobi and Y. Murata, *Science*, 2011, **333**, 613–616.
- S. Tashiro, M. Yamada and M. Shionoya, *Angew. Chem., Int. Ed.*, 2015, **54**, 5351–5354.
- X. Li, Y. Zhu, J. Shao, L. Chen, S. Zhao, B. Wang, S. Zhang, Y. Shao, H.-L. Zhang and X. Shao, *Angew. Chem., Int. Ed.*, 2015, **54**, 267–271.
- H. Sakurai, T. Daiko and T. Hirao, *Science*, 2003, **301**, 1878.
- H. Sakurai, *Bull. Chem. Soc. Jpn.*, 2021, **94**, 1579–1587.
- J. Han, Y. Uetake, Y. Yakiyama and H. Sakurai, *Chem. Commun.*, 2023, **59**, 4632–4635.
- M. Nishimoto, Y. Uetake, Y. Yakiyama, F. Ishiwari, A. Saeki and H. Sakurai, *J. Org. Chem.*, 2022, **87**, 2508–2519.
- T. Amaya, M. Hifumi, M. Okada, Y. Shimizu, T. Moriuchi, K. Segawa, Y. Ando and T. Hirao, *J. Org. Chem.*, 2011, **76**, 8049–8052.
- R. Fittig and E. Ostermayer, *Justus Liebigs Ann. Chem.*, 1873, **166**, 361–382.
- R. A. Haggam, *Synth. Commun.*, 2022, **52**, 1490–1499.
- N. Ngamsomprasert, J.-S. Dang, S. Higashibayashi, Y. Yakiyama and H. Sakurai, *Chem. Commun.*, 2017, **53**, 697–700.

