ChemComm

COMMUNICATION



View Article Online View Journal | View Issue

Check for updates

Cite this: Chem. Commun., 2024, 60, 3327

Received 31st January 2024, Accepted 27th February 2024

DOI: 10.1039/d4cc00492b

rsc.li/chemcomm

Designing a $\sigma^0 \pi^2$ singlet ground state carbene from dicationic carbones[†]

J. Philipp Wagner 问

Carbenes with a $\sigma^0 \pi^2$ singlet ground state are rare and little is known about their chemistry. Here, we study the potential formation of such carbenes by removal of two electrons from carbones/donor-substituted allenes. The desired electron configuration becomes favorable in the case of bis-diiminium substitution (CAAC motif).

The chemistry of carbenes:CR₂, can be understood from the distribution of their two non-bonding electrons over a σ and a π frontier orbital (Fig. 1A).¹ When each of these orbitals carries a single electron with parallel spin alignment as in the case of parent methylene (:CH₂), a triplet ground state ensues facilitating a radical-like reactivity. Long-lived versions of triplet carbenes can be realized by delocalization of the unpaired electrons and steric protection of the spin centres.² Likewise, the two nonbonding electrons can pair up in the low-lying σ orbital resulting in a $\sigma^2 \pi^0$ ground state displaying concerted reactivity. This situation gains favour when the σ orbital is lowered by introduction of electronegative substituents while the π orbital is elevated through conjugation with, for instance, lone pairs of electrons. Stable derivatives of such $\sigma^2 \pi^0$ singlet carbenes, especially N-heterocyclic ones, have created a tremendous impact on diverse areas of chemistry.³⁻⁷

The $\sigma^0 \pi^2$ configuration is located 59 kcal mol⁻¹ above the ground state in methylene making it much harder to stabilize this state.⁸ In principle, π -acceptors can be gainfully employed for this task, but this still often leaves the triplet as the ground state like in a recently reported diboryl carbene with a gap of more than 10 kcal mol⁻¹ to the $\sigma^0 \pi^2$ singlet.⁹ Hoffmann and Borden suggested that the experimentally known 2*H*-imidazol-2-ylidene can be characterized as a $\sigma^0 \pi^2$ singlet ground state carbene while it was previously rather thought of as a carbodiimide (Fig. 1B).^{10,11} The uncommon ground state becomes

possible because the carbene's lone pair is part of an aromatic sextet and the vacant σ orbital raises in energy through interaction with the in-plane lone pairs of the nitrogen atoms. Our lab has shown that this results in unusual chemistry, such as a preference for least-motion reactions with H₂ and alkenes¹² as opposed to other ordinary carbenes^{13,14} as well as a swift reaction with triplet oxygen at cryogenic temperatures¹⁵ despite the commonly encountered spin state selectivity of carbenes.^{16,17}

A major breakthrough in the chemistry of $\sigma^0 \pi^2$ carbenes has recently been accomplished with the isolation of rhodium-



Fig. 1 The usually high-lying $\sigma^0 \pi^2$ configuration of carbenes (A) becomes the ground state in **2** (B) and **4** (C) facilitating their unusual chemistry. The removal of two electrons from carbones can lead to the potential formation of dicationic carbenes (C).

Institut für Organische Chemie, Eberhard Karls Universität Tübingen,

Auf der Morgenstelle 18, 72076, Tübingen, Germany.

E-mail: philipp.wagner@orgchem.uni-tuebingen.de

[†] Electronic supplementary information (ESI) available: Optimized geometries and energies. See DOI: https://doi.org/10.1039/d4cc00492b

coordinated diphosphinocarbene 4 building on previous work on ephemeral metalated carbenes (Fig. 1C).¹⁸⁻²⁰ While donation of electron density from the metal centre into the σ orbital stabilizes this carbene, negative hyperconjugation of the π electrons was also discussed to play a role. The importance of the latter interaction suggests that a similar electronic structure could be achieved in carbodiphosphoranes by removal of two electrons to obtain a dicationic carbene (Fig. 1C). Carbodiphosphoranes can be considered as carbones - singlet carbon atoms formally coordinated by two σ -donor ligands bestowing the carbon atom with two lone pairs of electrons.²¹⁻²⁴ Carbodicarbenes and carbon suboxide are further experimentally known ('hidden') congeners of this compound class^{25,26} prompting us to explore their viability as carbene precursors through formal removal of one lone pair from the central carbon atom. Therefore, we optimized the dications in Fig. 2 in their respective singlet and triplet states at the B3LYP-D3/ def2-TZVPP level of theory and further computed the singlettriplet gap with the more accurate G4MP2 method where possible.27,28

The simplest dicationic carbone derivative, diphosphonium ion 5, indeed appears to adopt the electronic structure of a carbene displaying carbon-centred frontier orbitals with a doubly occupied σ orbital and vacant π orbital in its singlet state (see ESI† for all orbitals).‡ The C–P bonds are slightly shorter than the expected single bond length²⁹ and the Wiberg bond index (WBI) comes out near unity. The P–C–P angle of 139.7° is rather large for a singlet carbene which becomes understandable from the repelling substituents of like charge. The wide angle favours the triplet making it the ground state by a substantial 23.1 kcal mol⁻¹ (G4MP2). Since the lowest singlet corresponds to the $\sigma^2 \pi^0$ configuration, we performed a state averaged NEVPT2 calculation with a CAS(2,2) reference to study the $\sigma^0 \pi^2$ state, locating it at 73.8 kcal mol⁻¹ above the ground state. This result reveals that negative hyperconjugation to phosphonium groups is far from sufficient to stabilize the $\sigma^0 \pi^2$ state and re-emphasizes the role of the metal in 4.¹⁸ A comparable geometric and electronic structure is observed for the methylated diphosphonium ion **6**.

We reasoned that a more powerful electron acceptor is required, turning to carbon suboxide dication 7 in the following. The bent singlet state again assumes a $\sigma^2 \pi^0$ configuration with a large singlet-triplet gap of 22.3 kcal mol⁻¹. Natural resonance theory (NRT) predicts a dominant carbene structure with a weight of 63.6% in which two acylium ions carry the positive charges. The bending of the molecule is in line with the presence of a lone pair at the central carbon atom. The molecule is stabilized by in-plane delocalization of the carbene's nonbonding electrons producing mesomeric structures with a ketene unit. This might explain the Wiberg bond index of 1.328 and the rather short C-C bond distance of 1.38 Å, which is closer in length to a typical double than a single bond.

The dicationic carbodicarbene **8** displays a much smaller singlet triplet gap of 3.5 kcal mol⁻¹. The outermost, carboncentred orbitals are of σ - and π -type with the former being doubly occupied. Although the NRT analysis predicts a rather



Fig. 2 Structures of the herein considered singlet dications and their singlet-triplet gaps at the B3LYP-D3/def2-TZVPP and G4MP2 levels of theory.

complex set of structures, drawing a carbene appears reasonable since the imidazolium substituents can well support the positive charges.

The short C–C bond distances of 1.37 Å and the Wiberg bond index well above one cast doubt on the assignment of a carbene structure featuring two single bonds to the central atom in the case of **8**. Close inspection of the molecular orbitals reveals the presence of a doubly occupied π -type orbital that is all-bonding along the central C–C–C unit and not counteracted by a higher-lying filled orbital of strongly antibonding character. While the structure of the bis-benzimidazolium carbene **9** is comparable to **8**, dication **10** with its saturated backbone should not be regarded as a carbene. The latter molecule's LUMO still corresponds to a carbon-centred π orbital, but the two highest occupied orbitals are associated with the nitrogen atoms' lone pairs; the carbene σ orbital only represents the HOMO–2. In addition, the central C–C–C unit of dication **10** does not linearize in its triplet state as seen for molecules **5–9**.

So far, some of the considered carbone dications can well be considered as carbenes, but their singlet states pertain to the conventional $\sigma^2 \pi^0$ configuration. Hence, we reckoned that the carbone's substituents (ligand L) are required to be stronger π acceptors which is seen realized in the cyclic alkyl amino carbene (CAAC) motif.³⁰ Although the neutral precursor might not necessarily correspond to a carbone but rather to a donorsubstituted, linear allene, the singlet dication 11 indeed displays the desired $\sigma^0 \pi^2$ configuration (Fig. 3, top). The π orbital presents a significant coefficient of the atomic 2p orbital at the carbene carbon atom and indicates resonance with the neighbouring iminium ion groups acting as powerful electron sinks to stabilize the unconventional singlet state. Natural resonance theory confirms this analysis assigning the carbene as the dominant mesomeric structure (weight 30.2%) with delocalization of the lone pair facilitated by the iminium ions.

Despite this initial success, carbene **11** still prefers a triplet ground state by 10.9 kcal mol⁻¹. This motivated us to employ a second mode of stabilization targeting the carbene's σ orbital. A look at the LUMO in Fig. 3 shows that the vacant orbital is perfectly aligned for hyperconjugation with the five-membered rings' vicinal C–C bonds. Hence, we decided to bring the silicon



Fig. 3 Frontier orbitals of **11** and its dominant resonance structures.



Fig. 4 Transition state of the hydrogen insertion reaction of 12.

 β effect to use and placed SiMe₂ groups at the relevant position yielding dication **12**. This additional mode of stabilization helped to tip the balance finally predicting a $\sigma^0 \pi^2$ singlet ground state for **12**. The singlet-triplet gap amounts to -5.7 kcal mol⁻¹ at the G4MP2 level of theory.

Natural resonance theory produces a number of structures featuring silylium ion centres which is understandable from the elongated carbon-silicon bonds in **12** (2.06 Å). The central C–C bonds assume the length of a typical double bond bringing the carbene character of the dication once again into question. Therefore, we decided to verify the nature of the molecule by studying its single bond insertion reactivity. The relevant transition state for the insertion of the central carbon atom into the dihydrogen molecule is depicted in Fig. 4 and comes with a barrier of 21.9 kcal mol⁻¹. As expected for $\sigma^0 \pi^2$ carbenes,¹² the reaction mode corresponds to a σ -approach but deviates from the least-motion trajectory because the reaction centre is blocked by the two *N*-methyl substituents. The favourable bond insertion reactivity corroborates the nature of **12** as a $\sigma^0 \pi^2$ singlet ground state carbene.

In summary, we have demonstrated that some carbone dications can be viewed as carbenes. The unconventional $\sigma^0 \pi^2$ state becomes energetically accessible by stabilizing the π orbital through conjugation with iminium ions (CAAC scaffold) and destabilizing the σ orbital *via* hyperconjugation with carbon–silicon bonds. The experimental realization of such carbenes appears to be a worthwhile but challenging endeavour.

This work was generously funded by the Fonds der Chemischen Industrie in the form of a Liebig fellowship. Computations were performed on the BwForCluster JUSTUS 2 facilitated by bwHPC and the DFG (INST 40/575-1 FUGG). The author thanks Prof. Holger Bettinger for his valuable support.

Conflicts of interest

There are no conflicts to declare.

Notes and references

‡ We note that the singlet dications considered herein do not display a plane of symmetry with the exception of 7 rendering a rigorous definition of σ and π orbitals unattainable.

Communication

- 1 K. B. Eisenthal, R. A. Moss and N. J. Turro, Science, 1984, 225, 1439–1445.
- 2 H. Tomioka, E. Iwamoto, H. Itakura and K. Hirai, *Nature*, 2001, **412**, 626–628.
- 3 A. J. Arduengo, R. L. Harlow and M. Kline, *J. Am. Chem. Soc.*, 1991, **113**, 361–363.
- 4 A. Igau, A. Baceiredo, G. Trinquier and G. Bertrand, Angew. Chem., Int. Ed. Engl., 1989, 28, 621–622.
- 5 D. Bourissou, O. Guerret, F. P. Gabbaï and G. Bertrand, *Chem. Rev.*, 2000, **100**, 39–92.
- 6 M. N. Hopkinson, C. Richter, M. Schedler and F. Glorius, *Nature*, 2014, **510**, 485–496.
- 7 D. Munz, Organometallics, 2018, 37, 275-289.
- 8 A. Kalemos, T. H. DunningJr., A. Mavridis and J. F. Harrison, *Can. J. Chem.*, 2004, **82**, 684–693.
- 9 Y. Shibutani, S. Kusumoto and K. Nozaki, J. Am. Chem. Soc., 2023, 145, 16186-16192.
- 10 G. Maier and J. Endres, Chem. Eur. J., 1999, 5, 1590-1597.
- 11 B. Chen, A. Y. Rogachev, D. A. Hrovat, R. Hoffmann and W. T. Borden, J. Am. Chem. Soc., 2013, 135, 13954–13964.
- 12 S. F. Clewing and J. P. Wagner, *J. Org. Chem.*, 2021, **86**, 15247–15252. 13 R. Hoffmann, *J. Am. Chem. Soc.*, 1968, **90**, 1475–1485.
- 14 A. E. Keating, S. R. Merrigan, D. A. Singleton and K. N. Houk, J. Am. Chem. Soc., 1999, 121, 3933-3938.
- 15 J. P. Wagner, J. Am. Chem. Soc., 2022, 144, 5937-5944.
- 16 W. Sander, Angew. Chem., Int. Ed. Engl., 1990, 29, 344-354.
- 17 J. P. Wagner, Phys. Chem. Chem. Phys., 2022, 24, 25834-25841.

- 18 C. Hu, X.-F. Wang, J. Li, X.-Y. Chang and L. L. Liu, *Science*, 2024, **383**, 81–85.
- 19 J. Vignolle, H. Gornitzka, L. Maron, W. W. Schoeller, D. Bourissou and G. Bertrand, J. Am. Chem. Soc., 2007, **129**, 978–985.
- 20 J. Ruiz, M. E. G. Mosquera, G. García, E. Patrón, V. Riera, S. García-Granda and F. Van der Maelen, *Angew. Chem., Int. Ed.*, 2003, 42, 4767–4771.
- 21 F. Ramirez, N. B. Desai, B. Hansen and N. McKelvie, J. Am. Chem. Soc., 1961, 83, 3539-3540.
- 22 R. Tonner, F. Öxler, B. Neumüller, W. Petz and G. Frenking, *Angew. Chem., Int. Ed.*, 2006, **45**, 8038–8042.
- 23 G. Frenking, R. Tonner, S. Klein, N. Takagi, T. Shimizu, A. Krapp, K. K. Pandey and P. Parameswaran, *Chem. Soc. Rev.*, 2014, 43, 5106–5139.
- 24 G. Frenking, M. Hermann, D. M. Andrada and N. Holzmann, *Chem. Soc. Rev.*, 2016, **45**, 1129–1144.
- 25 R. Tonner and G. Frenking, Angew. Chem., Int. Ed., 2007, 46, 8695–8698.
- 26 C. A. Dyker, V. Lavallo, B. Donnadieu and G. Bertrand, Angew. Chem., Int. Ed., 2008, 47, 3206–3209.
- 27 P. J. Stephens, F. J. Devlin, C. F. Chabalowski and M. J. Frisch, J. Chem. Phys., 1994, 98, 11623–11627.
- 28 L. A. Curtiss, P. C. Redfern and K. Raghavachari, J. Chem. Phys., 2007, 127, 124105.
- 29 P. Pyykkö and M. Atsumi, Chem. Eur. J., 2009, 15, 186-197.
- 30 V. Lavallo, Y. Canac, C. Präsang, B. Donnadieu and G. Bertrand, Angew. Chem., Int. Ed., 2005, 44, 5705–5709.