


ChemComm

Chemical Communications

rsc.li/chemcomm



ISSN 1359-7345

COMMUNICATION

Mingji Dai *et al.*

Divergent synthesis of δ -valerolactones and furanones
via palladium or copper-catalyzed α -hydroxycyclopropanol
ring opening cyclizations



Cite this: *Chem. Commun.*, 2024, 60, 10112

Received 30th June 2024,
Accepted 31st July 2024

DOI: 10.1039/d4cc03255a

rsc.li/chemcomm

Divergent synthesis of δ -valerolactones and furanones *via* palladium or copper-catalyzed α -hydroxycyclopropanol ring opening cyclizations†

Pedro de Andrade Horn,^{‡a} Michael J. E. Collins,^{‡b} Cyrus C. Gudeman,^b Alexandra A. Fresh^{ib} ^a and Mingji Dai^{ib} ^{*bc}

Cyclopropanols are versatile starting materials which can undergo various ring opening reactions due to their intrinsic ring strain. Herein, we report two transition metal-catalyzed α -hydroxycyclopropanol ring opening cyclizations to divergently transform the same α -hydroxycyclopropanol substrate into two different products of enhanced value. One is a palladium-catalyzed α -hydroxycyclopropanol ring opening carbonylative lactonization to synthesize δ -valerolactones. The other one is a copper-catalyzed α -hydroxycyclopropanol ring opening cyclization to access furanones.

Cyclopropanols have been widely utilized in organic synthesis because of their unique and versatile reactivity profiles.^{1–3} Due to the intrinsic ring strain of the cyclopropane and the reactivity of the hydroxyl functional group, cyclopropanols can undergo various ring opening reactions. In general, a one-electron cyclopropanol ring opening pathway would generate a β -keto radical and a two-electron cyclopropanol ring opening pathway would give a homoenolate. If a transition metal is involved in the ring opening process, a metal homoenolate would be produced.⁴ The resulting β -keto radical and (metal) homoenolate can then engage in different downstream reactivity modes to give various products.^{5–7} We have been developing transition metal-catalyzed cyclopropanol ring opening reactions to facilitate chemical synthesis of biologically active natural products and pharmaceutically important compounds.^{8–11} In 2016 and 2020, we reported palladium-catalyzed hydroxycyclopropanol ring opening carbonylative lactonizations to synthesize either oxaspirolactones (Fig. 1A, **1** \rightarrow **2** \rightarrow **3**)¹² or fused bicyclic lactones (Fig. 1B, **4** \rightarrow **5** \rightarrow **6**)¹³ and have applied these enabling

carbonylation chemistry to synthesize a collection of natural products.^{14–16} While we were developing the fused bicyclic lactone synthesis, we discovered that the same cyclopropanol starting material **4** can undergo a copper-catalyzed cyclopropanol ring opening cyclization to give THF/THP-containing product **8** presumably *via* intermediate **7** (Fig. 1C).¹⁷ We then wondered if α -hydroxycyclopropanol **9** can undergo similar divergent pathways to synthesize δ -valerolactone **11** *via* the palladium-catalyzed ring opening carbonylative lactonization process and furanone **13** *via* the copper-catalyzed ring opening cyclization process (Fig. 1D). To realize these transformations, a few challenges need to be addressed. First, the 1,2-diol moiety of **9** can chelate on the transition metal and may deactivate the catalyst. Second, such α -hydroxycyclopropanol is prone to Pinacol rearrangement to partially release the ring strain and produce a cyclobutanone.¹⁸ Third, the lactonization process would generate a six-membered lactone, which has not been reported from a cyclopropanol ring opening carbonylation process. Fourth, once the Pd or Cu homoenolate is generated, β -hydride elimination, dimerization, and over oxidation are always competing pathways which need to be suppressed. Despite these challenges, the proposed divergent synthesis of δ -valerolactone **11** and furanone **13** from α -hydroxycyclopropanol **9** would further expand cyclopropanol chemistry and offer alternative methods to prepare the target structures, which are frequently found in biologically active molecules, especially natural products.^{19–23} Herein, we report the details of our exploration which led to divergent approaches to either δ -valerolactone **11** or furanone **13** from the same α -hydroxycyclopropanol **9** *via* a Pd-catalyzed ring opening carbonylative lactonization or a Cu-catalyzed ring opening cyclization, respectively.

We started our investigation with cyclopropanol **9a**, which was prepared *via* a modified Kulinkovich reaction (see the ESI†)²⁴ and used as a model compound for optimization of the reaction conditions (Table 1). When it was treated with Pd(OAc)₂ (10 mol%) and DDQ (2.0 eq.) in benzene at room temperature (the conditions we developed for the bicyclic

^a Department of Chemistry, Purdue University, West Lafayette, IN 47907, USA

^b Department of Chemistry, Emory University, Atlanta, GA 30322, USA.

E-mail: mingji.dai@emory.edu; Tel: 001-404-727-4299

^c Department of Pharmacology and Chemical Biology, Emory University, Atlanta, GA 30022, USA

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4cc03255a>

‡ Contributed equally.






Fig. 1 Our previous development in Pd and Cu-catalyzed cyclopropanol ring opening (carbonylative) cyclization and this work.

lactone synthesis),¹³ desired product **11a** was only produced in 13% yield (entry 1). Changing the palladium catalyst from Pd(OAc)₂ to Pd(PPh₃)₄ (entry 2), Pd(TFA)₂ (entry 3), and Pd(PPh₃)₂Cl₂ (entry 4) only slightly affected the reaction yield (11–19%). Switching the solvent from benzene to THF increased the yield more significantly (entries 5 and 6). When the combination of Pd(OAc)₂ and THF was used, the yield of the desired product **11a** increased to 55% yield (entry 6). When the amount of DDQ was reduced from 2.0 equivalents to 1.2 equivalents, **11a** was isolated in 59% yield (entry 7). THF was also found to be superior to MTBE (methyl *tert*-butyl ether), 1,4-dioxane, and DMSO (entries 8–10). Interestingly, when Cu(OTf)₂ (2.0 eq.) was used as oxidant, instead of **11a**, furanone **13a** was obtained in 70% yield (entry 11), which encouraged us to further develop a general cyclopropanol ring opening cyclization to synthesize furanones.

We then started to further optimize the furanone synthesis conditions and learned that palladium catalyst is not necessary. With 10 mol% of Cu(OTf)₂ as catalyst and DDQ (1.0 eq.) as oxidant, **13a** was obtained in 74% yield (entry 12). While CuCl₂ (entry 13), CuBr₂ (entry 14), CuSO₄ (entry 15), and CuTC (copper thiophene-2-carboxylate, entry 16) are less effective than Cu(OTf)₂, the more economical Cu(OAc)₂ (entry 17) is superior to Cu(OTf)₂ and **13a** was produced in 93% isolated yield.

Table 1 Optimization of reaction conditions

Entry	Conditions for 9a to 11a	Yield ^a (%) (11a)
1	Pd(OAc) ₂ (10 mol%), DDQ (2.0 eq.), PhH	13 ^b
2	Pd(PPh ₃) ₄ (10 mol%), DDQ (2.0 eq.), PhH	18
3	Pd(TFA) ₂ (10 mol%), DDQ (2.0 eq.), PhH	11
4	Pd(PPh ₃) ₂ Cl ₂ (10 mol%), DDQ (2.0 eq.), PhH	19
5	Pd(PPh ₃) ₂ Cl ₂ (10 mol%), DDQ (2.0 eq.), THF	28
6	Pd(OAc) ₂ (10 mol%), DDQ (2.0 eq.), THF	55
7	Pd(OAc) ₂ (10 mol%), DDQ (1.2 eq.), THF	59 ^b
8	Pd(OAc) ₂ (10 mol%), DDQ (1.2 eq.), MTBE	25
9	Pd(OAc) ₂ (10 mol%), DDQ (1.2 eq.), 1,4-dioxane	35
10	Pd(OAc) ₂ (10 mol%), DDQ (2.0 eq.), DMSO	0
11	Pd(OAc) ₂ (10 mol%), Cu(OTf) ₂ (2.0 eq.), PhH	70 (13a)

Entry	Conditions for 9a to 13a	Yield ^a (%) (13a)
12	Cu(OTf) ₂ (10 mol%), DDQ (1.0 eq.), THF	74
13	CuCl ₂ (10 mol%), DDQ (1.0 eq.), THF	66
14	CuBr ₂ (10 mol%), DDQ (1.0 eq.), THF	49
15	CuSO ₄ (10 mol%), DDQ (1.0 eq.), THF	0
16	CuTC (10 mol%), DDQ (1.0 eq.), THF	82
17	Cu(OAc) ₂ (10 mol%), DDQ (1.0 eq.), THF	93 ^b
18	Cu(OAc) ₂ (5.0 mol%), DDQ (1.0 eq.), THF	84
19	Cu(OAc) ₂ (1.0 mol%), DDQ (1.0 eq.), THF	58
20	Cu(OAc) ₂ (10 mol%), TFBQ (1.0 eq.), THF	74
21	Cu(OAc) ₂ (10 mol%), Chloranil (1.0 eq.), THF	69
22	Cu(OAc) ₂ (10 mol%), 2,5-DCBQ (1.0 eq.), THF	37
23	Cu(OAc) ₂ (10 mol%), 2,6-DCBQ (1.0 eq.), THF	19
24	Cu(OAc) ₂ (1.0 eq.), THF	79
25	DDQ (1.0 eq.), THF	0

^a NMR yield. ^b Isolated yield.

Further reducing the catalyst loading to 5 mol% (entry 18) and 1.0 mol% (entry 19) resulted in lower yields (84% and 58%, respectively). Other oxidants including TFBQ (tetrafluoro-1,4-benzoquinone, entry 20), Chloranil (entry 21), 2,5-DCBQ (2,5-dichloro-1,4-benzoquinone, entry 22) and 2,6-DCBQ (2,6-dichloro-1,4-benzoquinone, entry 23) gave reduced yield. In addition, if DDQ or its byproduct complicates the reaction or purification, the reaction can be conducted with a stoichiometric amount of Cu(OAc)₂ to produce **13a** in 79% yield (entry 24). Given the similar price of DDQ and Cu(OAc)₂, the stoichiometric condition doesn't significantly affect the overall cost. In addition, no product **13a** was obtained without the copper catalyst (entry 25).²⁵

With both the δ -valerolactone and furanone synthesis conditions established, we started to probe the substrate scope of both transformations (Fig. 2). Twenty different α -hydroxycyclopropanols were prepared and subjected to the optimized reaction conditions. In general, the Cu-catalyzed furanone synthesis tends to give higher yield than the corresponding Pd-catalyzed δ -valerolactone synthesis. Both transformations can be conducted on gram scale with only a slight drop of the reaction yields. Alkyl (**11/13a-c**) and aryl groups (**11/13d-s**) are tolerated. The electronic properties of the aryl groups don't influence the reaction yield significantly. Halides such as fluoride, chloride, and bromide are not affected under both reaction conditions. The tolerance of chloride and bromide






Fig. 2 Substrate scope.

offers opportunity for cross coupling reactions to further functionalize the corresponding products.

While both 1- or 2-naphthyl group containing substrates are effective, no desired products (**11t** and **13t**) were obtained for substrate **9t** with a 3-pyridyl under both reaction conditions. Finally, we prepared enantio-enriched starting material **9a** (94% ee) from L-phenylalanine and evaluated if both reaction conditions would erode the stereochemistry at the α -position of the newly formed ketone. When **9a** (94% ee) was subjected to the carbonylation conditions, product **11a** was obtained in 55% yield and 85% ee. When **9a** (94% ee) was subjected to the furanone synthesis conditions, product **13a** was obtained in 80% yield and 94% ee. These results indicate the mildness of the reaction conditions and the potential application of these two methods to prepare enantio-enriched δ -valerolactones and furanones.

In summary, two novel cyclopropanol ring opening cyclization reactions were developed to divergently transform the same α -hydroxycyclopropanol substrate to either a δ -valerolactone or a furanone. The δ -valerolactone synthesis was catalyzed by $\text{Pd}(\text{OAc})_2$ under carbon monoxide atmosphere and the furanone synthesis was catalyzed by $\text{Cu}(\text{OAc})_2$. DDQ was used as external oxidant in both reactions. An array of

δ -valerolactones and furanones were prepared. The reactions can be scaled up to gram scale. Overall, these two cyclopropanol ring opening cyclization reactions provide mild alternatives to synthesize δ -valerolactones and furanones, which are frequently found in biologically active molecules.

This work was financially supported by NSF 2349014.

Data availability

The ESI[†] includes the experimental procedures, compound characterization data, and NMR spectra. ESI[†] is included in the submission and will be available to the readers *via* the Chemical Communication online publication system.

Conflicts of interest

There are no conflicts to declare.

Notes and references

- X. Cai, W. Liang and M. Dai, *Tetrahedron*, 2019, **75**, 193–208.
- O. G. Kulinkovich, *Chem. Rev.*, 2003, **103**, 2597–2632.
- D. H. Gibson and C. H. DePuy, *Chem. Rev.*, 1974, **74**, 605–623.
- A. Nikolaev and A. Orellana, *Synthesis*, 2016, 1741–1768.



- 5 T. R. McDonald, L. R. Mills, M. S. West and S. A. L. Rousseaux, *Chem. Rev.*, 2021, **121**, 3–79.
- 6 L. R. Mills and S. A. L. Rousseaux, *Eur. J. Org. Chem.*, 2019, 8–26.
- 7 H. Yan, G. S. Smith and F.-E. Chen, *Green Syn. Catal.*, 2022, **3**, 219–226.
- 8 Y. Li, Z. Ye, T. M. Bellman, T. Chi and M. Dai, *Org. Lett.*, 2015, **17**, 2186–2189.
- 9 Z. Ye and M. Dai, *Org. Lett.*, 2015, **17**, 2190–2193.
- 10 Z. Ye, K. E. Gettys, X. Shen and M. Dai, *Org. Lett.*, 2015, **17**, 6074–6077.
- 11 Z. Ye, X. Cai, J. Li and M. Dai, *ACS Catal.*, 2018, **8**, 5907–5914.
- 12 D. C. Davis, K. L. Walker, C. Hu, R. N. Zare, R. M. Waymouth and M. Dai, *J. Am. Chem. Soc.*, 2016, **138**, 10693–10699.
- 13 X. Cai, W. Liang, M. Liu, X. Li and M. Dai, *J. Am. Chem. Soc.*, 2020, **142**, 13677–13682.
- 14 K. Ma, X. Yin and M. Dai, *Angew. Chem., Int. Ed.*, 2018, **57**, 15209–15212.
- 15 X. Yin, K. Ma, Y. Dong and M. Dai, *Org. Lett.*, 2020, **22**, 5001–5004.
- 16 H. S. Sims, P. de Andrade Horn, R. Isshiki, M. Lim, Y. Xu, R. H. Grubbs and M. Dai, *Angew. Chem., Int. Ed.*, 2022, **61**, e202115633.
- 17 W. Liang, X. Cai and M. Dai, *Chem. Sci.*, 2021, **12**, 1311–1316.
- 18 M. M. Hussain, H. Li, N. Hussain, M. Ureña, P. J. Carroll and P. J. Walsh, *J. Am. Chem. Soc.*, 2009, **131**, 6516–6524.
- 19 V. K. Omanakuttan, J. John and H. Hopf, *Eur. J. Org. Chem.*, 2021, 163–201.
- 20 A. Husain, S. A. Khan, F. Iram, M. A. Iqbal and M. Asif, *Eur. J. Med. Chem.*, 2019, **171**, 66–92.
- 21 R. Liu, S. Yang, Z. Chen, X. Kong, H. Ding and X. Fang, *Org. Lett.*, 2020, **22**, 6948–6953.
- 22 Y. Matsuda, T. Iwabuchi, T. Wakimoto, T. Awakawa and I. Abe, *J. Am. Chem. Soc.*, 2015, **137**, 3393–3401.
- 23 M. Dangelov, A. Fernández-Figueiras, M. A. Ravutsov, E. Vakarelska, M. K. Marinova, N. R. Candeias and S. P. Simeonov, *ACS Catal.*, 2023, **13**, 1916–1925.
- 24 J. S. Kingsbury and E. J. Corey, *J. Am. Chem. Soc.*, 2005, **127**, 13813–13815.
- 25 Note: with Cu(OTf)₂ (2.0 eq.) in toluene at room temperature, **9a** underwent a Pinnacol rearrangement to give the corresponding cyclobutanone in 80% yield. The cyclobutanone is a known compound: T. Nordvik and U. H. Brinker, *J. Org. Chem.*, 2003, **68**, 9394–9399.

