


 Cite this: *Phys. Chem. Chem. Phys.*,  
 2024, 26, 6646

# Kinetics of the reaction of OH with methyl nitrate (223–343 K)<sup>†</sup>

 Christin Fernholz,<sup>id</sup> Fabienne Baumann,<sup>id</sup> Jos Lelieveld and John N. Crowley<sup>id</sup>\*

Rate coefficients ( $k_4$ ) for the reaction of hydroxyl radicals (OH) with methyl nitrate ( $\text{CH}_3\text{ONO}_2$ ) were measured over the temperature range 232–343 K using pulsed laser photolysis to generate OH and pulsed laser-induced fluorescence to detect it in real-time and under pseudo-first-order conditions. In order to optimize the accuracy of the rate coefficients obtained, the concentration of  $\text{CH}_3\text{ONO}_2$  (the reactant in excess) was measured on-line by absorption spectroscopy at 213.86 nm for which the absorption cross-section was also measured ( $\sigma_{213.86} = 1.65 \pm 0.09 \times 10^{-18} \text{ cm}^2 \text{ molecule}^{-1}$ ). The temperature-dependent rate coefficient is described by  $k_4(T) = 7.5 \times 10^{-13} \exp[-(1034 \pm 40)/T] \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  with a room temperature rate coefficient of  $k_4(296 \pm 2 \text{ K}) = (2.32 \pm 0.12) \times 10^{-14} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  where the uncertainty includes the statistical error of  $2\sigma$  and an estimation of the potential systematic bias of 5%. This new dataset helps to consolidate the database for this rate coefficient and to reduce uncertainty in the atmospheric lifetime of  $\text{CH}_3\text{ONO}_2$ . As part of this study, an approximate rate coefficient for the reaction of H-atoms with  $\text{CH}_3\text{ONO}_2$  ( $k_9$ ) was also derived at room temperature:  $k_9(298 \text{ K}) = (1.68 \pm 0.45) \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ .


 Received 5th January 2024,  
 Accepted 1st February 2024

DOI: 10.1039/d4cp00054d

[rsc.li/pccp](http://rsc.li/pccp)


## 1 Introduction

Organic nitrates are formed in the atmosphere during the degradation of hydrocarbons in the presence of nitrogen oxides.<sup>1</sup> The atmospheric lifetimes of organic nitrates can vary from seconds (*e.g.* for thermally unstable peroxy-nitrates such as  $\text{CH}_3\text{OONO}_2$  at temperatures close to room temperature) to months (*e.g.* for  $\alpha$ -carbonyl peroxy-nitrates such as  $\text{CH}_3\text{C}(\text{O})\text{OONO}_2$  at temperatures close to 220 K typical for the upper troposphere). In contrast to peroxy-nitrates, alkyl nitrates (such as  $\text{CH}_3\text{ONO}_2$ ) are generally thermally stable at the temperatures of our atmosphere and can be transported over long distances before being oxidized *via* reaction with OH or photolyzed.<sup>1–3</sup> Following OH-induced oxidation of hydrocarbons, alkyl nitrates are formed in a usually minor (termolecular) channel of the reaction of organic peroxy radicals ( $\text{RO}_2$ ) with NO:



Alternatively,  $\text{RONO}_2$  can be formed in the  $\text{NO}_3$ -induced oxidation of unsaturated biogenic hydrocarbons in the presence of  $\text{O}_2$ , which involves electrophilic addition of  $\text{NO}_3$  to a C=C double bond followed by further reaction of

the nitro-peroxy radicals thus formed.<sup>4</sup> The target molecule of this study,  $\text{CH}_3\text{ONO}_2$  is formed in the lower atmosphere when  $\text{CH}_3\text{O}_2$  radicals (*e.g.* from methane oxidation) react with NO. The formation of  $\text{CH}_3\text{ONO}_2$  *via* this reaction is inefficient, with a yield of <1% at room temperature,<sup>5</sup> but with some evidence for an increase in the yield at lower temperatures, commensurate with its formation in a termolecular process:<sup>6</sup>



In the lower atmosphere, the fate of  $\text{CH}_3\text{O}$  is dominated by reaction with  $\text{O}_2$  to form  $\text{HO}_2$  and HCHO. However, in the stratosphere, when  $\text{O}_2$  concentrations are much lower, some  $\text{CH}_3\text{O}$  can react with  $\text{NO}_2$ <sup>7</sup> and this represents a further source of  $\text{CH}_3\text{ONO}_2$ :





The important chemical loss processes for  $\text{CH}_3\text{ONO}_2$  in the atmosphere are photodissociation and reaction with OH, with thermal decomposition and reaction with  $\text{O}_3$  or  $\text{NO}_3$  insignificant.<sup>1,8</sup> Previous measurements indicate that R4 has a small rate coefficient, of the order of  $10^{-14} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  (see below). The photolysis of  $\text{CH}_3\text{ONO}_2$  proceeds *via* O–N bond fission to form a methoxy radical and  $\text{NO}_2$ , whereas the slow reaction with OH (R4) proceeds *via* H-abstraction to form an unstable radical that dissociates to formaldehyde and  $\text{NO}_2$ (R5):

Atmospheric Chemistry Department, Max Planck Institute for Chemistry,  
 Mainz 55128, Germany. E-mail: john.crowley@mpic.de

<sup>†</sup> Electronic supplementary information (ESI) available. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d4cp00054d>





Both HCHO and CH<sub>3</sub>O react much more rapidly with OH than does CH<sub>3</sub>ONO<sub>2</sub> and secondary chemistry is a potential source of systematic bias in the determination of the rate coefficient ( $k_4$ ) for the title reaction when using absolute rate methods. A further source of systematic bias in an absolute rate coefficient study of this slow reaction (R4) is the presence of reactive impurities in the CH<sub>3</sub>ONO<sub>2</sub> sample. Such problems can be avoided by working at low OH concentrations (and thus low fractional conversion of CH<sub>3</sub>ONO<sub>2</sub> to products) and by carefully purifying the CH<sub>3</sub>ONO<sub>2</sub> sample, thus minimising the levels of reactive impurities (see Section 2).

The rate coefficient for reaction (R4) has been reported by Gaffney *et al.*,<sup>9</sup> Nielsen *et al.*,<sup>10</sup> Talukdar *et al.*,<sup>8</sup> and Shallcross *et al.*,<sup>11</sup> but there are large differences in the rate coefficient at room temperature (a factor of 10) and its temperature dependence which is reported to be both positive and negative (see later). The goal of this study is to generate highly accurate rate coefficients for the title reaction at different (atmospherically relevant) temperatures and to resolve these differences in the literature and thus better constrain the lifetime of CH<sub>3</sub>ONO<sub>2</sub> in the atmosphere.

## 2 Experimental

Rate coefficients for the reaction of OH with CH<sub>3</sub>ONO<sub>2</sub> were determined using pulsed-laser-photolysis (PLP) pulsed-laser-induced fluorescence (PLIF) under pseudo-first-order conditions ( $[\text{CH}_3\text{ONO}_2] \gg [\text{OH}]$ ) in (mainly) N<sub>2</sub> and He bath gases. The concentration of methyl nitrate was measured *in situ* by optical absorption. The experimental setup has been described in detail elsewhere<sup>12</sup> and is only briefly described here.

### 2.1 PLP-PLIF technique

The reaction took place in a cylindrical, jacketed quartz reactor with a volume of approximately 300 cm<sup>3</sup>. The temperature inside the reactor was controlled by circulating a 60:40 ethylene glycol/water mixture (343–251 K) or ethanol ( $T < 251$  K) through an outer-jacket. The gas mixture enters the reactor about 15 cm before the photolysis zone, ensuring thermal equilibrium with the walls of the reactor. The reaction temperature was measured by inserting a recently calibrated K-type thermocouple into the photolysis zone before each experiment at a different temperature. 100 and 1000 torr manometers were used to measure the pressure in the reactor.

OH was generated by excimer laser photolysis (Coherent COMPex 205 F or Pro 201 F, ~20 ns width) of a precursor gas molecule, either H<sub>2</sub>O<sub>2</sub> (248 nm), HONO (351 nm) or CH<sub>3</sub>ONO<sub>2</sub> itself (248 nm) (see below).

OH radicals were excited at 281.997 nm ( $A^2\Sigma(\nu' = 1) \leftarrow X^2\Pi(\nu'' = 0)$ ,  $Q_{11}$  (1) transition) using a Nd-YAG pumped (532 nm, ~6 ns pulse width) and frequency-doubled dye laser (Quantel Brilliant B, Lambda Physik Scanmate/Scanmate UV

with KPD crystal). The resulting fluorescence signal was detected by a photomultiplier tube screened by a long-pass filter ( $\lambda > 290$  nm). The photolysis and excitation lasers were mainly operated at a repetition rate of 10 Hz using a pulse/delay generator (Stanford Research Systems DG 535), which was also used to control the delay between the photolysis and excitation laser pulses. Prior to the PLP pulse, 20 pulses of the PLIF laser were used to measure a background signal that includes scattered light from the excitation laser and electronic noise. The averaged background signal was subtracted before further processing of the measured data. Each datapoint was averaged between 10 and 40 times, depending on the measurement.

The total flow and pressure in the reactor were chosen such that an average, linear gas-velocity of ~13–19 cm s<sup>-1</sup> was achieved, ensuring that the gas in the photolysis region was replenished between laser pulses. For measurements in helium, the linear velocity was 5 or 6.6 cm s<sup>-1</sup> and the laser repetition rate was reduced to 5 Hz, to ensure complete gas exchange.

### 2.2 *In situ* absorption spectroscopy: The absorption cross-section of CH<sub>3</sub>ONO<sub>2</sub> at 213.86 nm

As the experiments were performed under pseudo first order conditions, accurate rate coefficients depend on accurate measurement of the concentration of the excess reactant. The concentration of methyl nitrate was therefore determined on-line, downstream of the reactor by optical absorption using a dual-beam absorption cell (path length 34.8 cm) equipped with a low-pressure Zn-lamp and interference filter ( $214 \pm 5$  nm) to isolate the 213.86 nm line. The temperature (close to room temperature) and pressure (a few percent lower than in the reactor) in the absorption cell were also measured and used to make corrections for gas-density differences in the reaction and absorption cell. Optical density (OD) due to CH<sub>3</sub>ONO<sub>2</sub> was converted to a concentration using the Beer–Lambert expression (1) and an absorption cross-section that was determined as described below.

$$\text{OD} = \ln(I_0/I) = \sigma_{213.86} l [\text{CH}_3\text{ONO}_2] \quad (1)$$

$\sigma_{213.86}$  is the CH<sub>3</sub>ONO<sub>2</sub> absorption cross-section (cm<sup>2</sup> molecule<sup>-1</sup>),  $l$  is the absorption path-length and  $I_0$  and  $I$  are the transmitted light intensity without and with absorber, respectively and  $[\text{CH}_3\text{ONO}_2]$  is the concentration of methyl-nitrate (molecule cm<sup>-3</sup>).

To measure the absorption cross-section, the absorption cell was first filled with N<sub>2</sub> to determine  $I_0$ . Subsequently, static samples of CH<sub>3</sub>ONO<sub>2</sub> in N<sub>2</sub> at different pressures were introduced into the cell from a storage bulb and the optical absorption was recorded. Two different storage bulbs (0.482% and 0.728%) made up using samples of CH<sub>3</sub>ONO<sub>2</sub> from two different syntheses (~ one month apart in time) were used to check for reproducibility. The storage bulbs were prepared using standard manometric methods with 10 and 1000 torr capacitance manometers with quoted accuracies of better than 1%.






Fig. 1 Optical density (OD) plotted against concentration of methyl nitrate ( $\text{molecule cm}^{-3}$ ) multiplied by path length  $l$  (cm). The slope of the corresponding, weighted linear regression yields an effective absorption cross-section of  $\sigma_{\text{CH}_3\text{ONO}_2}^{213.86} = (1.65 \pm 0.09) \times 10^{-18} \text{ cm}^2 \text{ molecule}^{-1}$ . The error is  $2\sigma$  and includes an estimate of systematic uncertainty.

Fig. 1 presents the results from 17 determinations of optical density at room-temperature. The slope of the weighted, linear regression gives an absorption cross-section of  $\sigma_{213.86} = (1.65 \pm 0.09) \times 10^{-18} \text{ cm}^2 \text{ molecule}^{-1}$ , where the uncertainty is  $2\sigma$  and includes an assessment of systematic bias (2%) which stems predominantly from uncertainty in the  $\text{CH}_3\text{ONO}_2$  mixing ratio and thus  $[\text{CH}_3\text{ONO}_2]$ . In addition, minor contributions stem from uncertainty in the optical path-length ( $<1\%$ ) and in determination of the optical density (0.05–1.5%). As an interference filter with a finite FWHM was used to isolate the 213.86 nm Zn-line, our cross-section should strictly be regarded as an “effective” cross-section since weaker atomic lines may also be detected. Our cross-section is significantly larger than the value of  $3.04 \times 10^{-19} \text{ cm}^2 \text{ molecule}^{-1}$  at 213.99 nm reported by Taylor *et al.*<sup>13</sup> but is similar to a value of  $2.16 \times 10^{-18} \text{ cm}^2 \text{ molecule}^{-1}$  at 214.01 nm measured by McMillan *et al.* and reported by Calvert and Pitts.<sup>14</sup> More importantly, our value is in excellent agreement with the more recent determination of Talukdar *et al.*,<sup>15</sup> who obtained  $\sigma_{213.86} = (1.67 \pm 0.08) \times 10^{-18} \text{ cm}^2 \text{ molecule}^{-1}$  using a Zn-lamp. We conclude that our cross-section is appropriate for converting *in situ* optical density measurements to accurate concentrations of  $\text{CH}_3\text{ONO}_2$ .

### 2.3 Determination of impurities in the $\text{CH}_3\text{ONO}_2$ sample

As the reaction between  $\text{CH}_3\text{ONO}_2$  and OH is slow, the presence of reactive impurities can bias the rate coefficients obtained to higher values. Likely impurities are  $\text{CH}_3\text{OH}$  and  $\text{HNO}_3$  which are involved in the synthesis of  $\text{CH}_3\text{ONO}_2$  as well as  $\text{NO}_2$  and HONO. Initial measurements revealed the presence of low levels of  $\text{HNO}_3$  in the  $\text{CH}_3\text{ONO}_2$  samples. In order to eliminate this (and other acidic gases) the sample flow was passed through  $\approx 20$  cm of PFA tubing containing nylon before entering the reactor. In some experiments (in which methyl nitrate itself was used as OH source)  $\text{CaCO}_3$  was used to remove  $\text{HNO}_3$ .

Table 1 Impurities and their contribution to the OH decay coefficient

Impurity	Fraction <sup>a</sup>	$k_{\text{impurity}}^b$	Contribution to $k'$ (%)
$\text{HNO}_3$	$<0.01$	$1.5 \times 10^{-13}$	$<0.06$
$\text{CH}_3\text{OH}$	$<0.01$	$9.0 \times 10^{-13}$	$<0.38$
HONO	$<0.03$	$6.0 \times 10^{-12}$	$<6.80$
$\text{NO}_2$	$<0.03$	$4.2 \times 10^{-12c}$	$<4.78$






<sup>a</sup> Relative concentration of impurity (in %) of the  $\text{CH}_3\text{ONO}_2$  concentration. <sup>b</sup> Rate coefficient ( $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ ) at room temperature. <sup>c</sup> Calculated for a pressure of 135 torr.

Following each new synthesis of  $\text{CH}_3\text{ONO}_2$ , levels of  $\text{HNO}_3$ , HONO,  $\text{CH}_3\text{OH}$  and  $\text{NO}_2$  were determined using a combination of IR and UV/Vis absorption spectroscopy. An FTIR (Bruker Vector 22) spectrometer coupled to an absorption cell of 45 cm pathlength<sup>16</sup> and equipped with an external MCT detector was used to determine impurity levels of  $\text{HNO}_3$  and  $\text{CH}_3\text{OH}$  in a flowing mixture of  $\text{CH}_3\text{ONO}_2$  in  $\text{N}_2$ , as used in the kinetic experiments. Generally, 32 spectra at a resolution of  $1 \text{ cm}^{-1}$  (acquisition time  $\approx 1$  min) were co-added to measure absorption in the 700–3750  $\text{cm}^{-1}$  range. HONO and  $\text{NO}_2$  impurities were investigated using multi-pass ( $l = 880$  cm) UV/Vis absorption spectroscopy (346–518 nm) using a H4-lamp light source coupled to a CCD detector (Andor DU420A-OE) and a 0.5 m monochromator (BM50) with 300 lines  $\text{mm}^{-1}$  grating. The spectral resolution (FWHM of a Hg line from a low-pressure discharge lamp) was 0.5 nm.

The results of the impurity determinations (none were detected and upper limits were established) are summarized in Table 1, which shows that the rate coefficients obtained were not significantly influenced by impurities in the  $\text{CH}_3\text{ONO}_2$  sample.

### 2.4 OH sources

During the course of this study, three different sources of OH were tested. The photolysis of HONO at 351 nm (R6) and the photolysis of  $\text{H}_2\text{O}_2$  at 248 nm (R7) both generate OH in the ground vibrational state.<sup>17</sup> The photolysis of  $\text{CH}_3\text{ONO}_2$  at 248 nm is an indirect source of OH<sup>15</sup> via formation of H atoms (R8b) and (R9).



In Section 3.2 and 3.3, we describe how the OH-decays obtained using the 248 nm photolysis laser (*i.e.*, (R7) or (R8b) and (R9)) are strongly influenced by secondary reactions involving  $\text{CH}_3\text{O}$  and  $\text{NO}_2$  resulting from the photolysis of  $\text{CH}_3\text{ONO}_2$ . Only data obtained using the photolysis of HONO at 351 nm (a wavelength at which  $\text{CH}_3\text{ONO}_2$  no longer absorbs significantly) were used to derive the rate coefficient for the title reaction.



## 2.5 Chemicals

The synthesis of  $\text{CH}_3\text{ONO}_2$ , the nitration of methanol with  $\text{HNO}_3$ , was carried out according to Black and Babers<sup>18</sup> but with a modification of the purification step.  $\sim 62$  mL of Nitric acid ( $\text{HNO}_3$ ; Sigma-Aldrich,  $\geq 65\%$ ) was pre-cooled (ice bath) in a 250 ml Erlenmeyer flask.  $\sim 31$  mL of Methanol ( $\text{CH}_3\text{OH}$ ; Sigma-Aldrich,  $\geq 99.9\%$ ) was pre-cooled in a 100 mL Erlenmeyer flask.  $\sim 62$  mL of sulphuric acid (Roth, 98%) was added to the nitric acid. The remaining sulphuric acid was added to the methanol using a dropping funnel with vigorous stirring, whereby a temperature of  $14^\circ\text{C}$  was not exceeded. The  $\text{CH}_3\text{OH}/\text{H}_2\text{SO}_4$  mixture was then added. The temperature of the mixture increased to about  $45^\circ\text{C}$  and was subsequently kept between  $32^\circ\text{C}$  and  $35^\circ\text{C}$  for about 15 minutes. The mixture was then rapidly cooled in an ice-water bath for 5 minutes and then transferred to a separating funnel. The aqueous phase and part of the organic phase were discarded.

The unwashed methyl nitrate was transferred to a bubbler and  $\text{N}_2$  was bubbled through the liquid for about 15 min to remove any dissolved methanol. The liquid was then vacuum-distilled through a drying tube with  $\text{CaCl}_2$  (Roth,  $\geq 94\%$ , dehydrated) before the transmitted gas was trapped in a cold-finger at liquid nitrogen temperature. The methyl nitrate was then subjected to repeated freeze-pump-thaw cycles at  $-196^\circ\text{C}$ . Nitrogen ( $\text{N}_2$ , Air-Liquide, 99.999%) and helium (He, Westfalen, 99.999%) were used as buffer gases in the experiments. HONO was prepared *in situ* by adding 5–10 drops of 0.1 M  $\text{NaNO}_2$  solution (Roth,  $\geq 98\%$ ) into a 20 wt%  $\text{H}_2\text{SO}_4$  solution (Roth, 98%). Hydrogen peroxide ( $\text{H}_2\text{O}_2$ ; AppliChem, 35%) was vacuum distilled to  $>90\%$  purity.

## 3 Results and discussion

In all experiments, the methyl nitrate concentration was 5–6 orders of magnitude larger than that of OH (typically  $1 \times 10^{10}$  molecule  $\text{cm}^{-3}$ ) and pseudo-first order conditions prevailed. The decay of OH can therefore be described by

$$[\text{OH}]_t = [\text{OH}]_0 \exp(-k't) \quad (2)$$

where  $[\text{OH}]_t$  is the OH concentration at time  $t$  after the photolysis laser pulse,  $[\text{OH}]_0$  is the initial OH concentration and  $k'$  is the first-order decay coefficient. The latter is defined as:

$$k' = k_4[\text{CH}_3\text{ONO}_2] + k_d \quad (3)$$

where  $k_4$  is the bimolecular rate coefficient ( $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ ) of the reaction of methyl nitrate with OH and  $k_d$  ( $\text{s}^{-1}$ ) accounts for diffusion of OH from the detection zone and its reaction with  $\text{H}_2\text{O}_2$  or HONO.

### 3.1 Generation of OH from HONO photolysis at 351 nm

Fig. 2 shows an example of the OH fluorescence signal plotted against time after the excimer laser pulse. The typical HONO concentration in this experiment is approx.  $1.5 \times 10^{13}$  molecule  $\text{cm}^{-3}$ , which results in an initial OH concentration of  $\sim 1 \times 10^{10}$  molecule  $\text{cm}^{-3}$ . As expected (eqn (2)), the decay of




Fig. 2 OH-LIF signal (datapoints) with mono-exponential fits (solid lines) plotted against time after the photolysis laser pulse. Data were obtained using HONO as OH source at room temperature, the pressure was 134.0 torr ( $\text{N}_2$  bath-gas).

OH is exponential (over 2 orders of magnitude). Decay coefficients ( $k'$ ,  $\text{s}^{-1}$ ), obtained by least-squares-fitting of eqn (2) to the OH-decays, are plotted against the respective concentration of  $\text{CH}_3\text{ONO}_2$  in Fig. 3. A linear dependence of  $k'$  on the  $\text{CH}_3\text{ONO}_2$  concentration (eqn (3)) was observed under all experimental conditions, enabling derivation of  $k_4(T)$ . Variability in the y-axis intercept of  $\sim 200$ – $250 \text{ s}^{-1}$  reflects slight changes in the HONO concentration between experiments at different temperatures. Prior experiments indicate that the contribution of diffusion to  $k'$  is *circa*  $50$ – $100 \text{ s}^{-1}$ , depending mainly on the pressure (or bath gas) and the flow rate. As the rate coefficient for the reaction between OH and HONO is  $6.0 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$




Fig. 3 First-order OH decay-coefficient ( $k'$ ) plotted against the corresponding methyl nitrate concentration  $[\text{CH}_3\text{ONO}_2]$  (data obtained using HONO as the OH precursor). The slope of the linear regression (orange:  $k(298 \text{ K}) = (2.31 \pm 0.15) \times 10^{-14} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ , green:  $k(232 \text{ K}) = (7.78 \pm 0.92) \times 10^{-15} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ ,  $k(343 \text{ K}) = (3.62 \pm 0.22) \times 10^{-14} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ ) is the rate coefficient  $k_4$  under these experimental conditions. Error bars represent  $2\sigma$  statistical uncertainty.



Table 2 Experimental conditions and rate coefficients ( $k_4$ ) for the reaction OH + CH<sub>3</sub>ONO<sub>2</sub>

$T$ (K)	$p$ (torr)	$k_4$ ( $10^{-14}$ cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup> )	$\Delta k_4^a$ ( $10^{-14}$ cm <sup>3</sup> molecule <sup>-1</sup> s <sup>-1</sup> )	Bath gas
223.3	101.2	0.84	0.07	He
231.7	134.6	0.78	0.09	N <sub>2</sub>
239.6	134.3	0.87	0.09	N <sub>2</sub>
247.3	135.7	1.05	0.10	N <sub>2</sub>
251.5	134.8	1.28	0.05	N <sub>2</sub>
259.7	127.0	1.42	0.07	N <sub>2</sub>
266.0	136.3	1.49	0.10	N <sub>2</sub>
268.7	135.2	1.56	0.10	N <sub>2</sub>
276.2	134.8	1.69	0.09	N <sub>2</sub>
282.2	135.1	1.86	0.12	N <sub>2</sub>
289.0	135.1	2.07	0.15	N <sub>2</sub>
294.2	101.6	2.28	0.11	He
294.5	135.5	2.21	0.20	N <sub>2</sub>
295.2	98.8	2.14	0.13	N <sub>2</sub>
295.8	133.5	2.39	0.08	N <sub>2</sub>
296.2	99.1	2.41	0.15	N <sub>2</sub>
296.7	201.0	2.48	0.15	N <sub>2</sub>
297.5	135.1	2.34	0.17	N <sub>2</sub>
297.7	400.4	2.12	0.30	N <sub>2</sub>
298.0	134.0	2.31	0.15	N <sub>2</sub>
298.2	68.5	2.23	0.18	N <sub>2</sub>
317.8	135.2	2.78	0.19	N <sub>2</sub>
342.7	135.4	3.62	0.22	N <sub>2</sub>

Notes: All rate coefficients were obtained using the 351 nm photolysis of HONO as OH-source. <sup>a</sup> Uncertainty is 2 $\sigma$ , statistical only.

(at room temperature), the approximate HONO concentration can be calculated to be  $< 4 \times 10^{13}$  molecule cm<sup>-3</sup>. This is consistent with the concentration obtained using the weak absorption ( $OD < 1 \times 10^{-3}$ ) observed at 213.86 nm ( $\sigma = 1.8 \times 10^{-18}$  cm<sup>2</sup> molecule<sup>-1</sup>) when adding HONO in the absence of CH<sub>3</sub>ONO<sub>2</sub>.

Within experimental uncertainty, variation of the pressure or bath-gas (N<sub>2</sub> or He) (for a given temperature) had no effect on

the rate coefficient, which is thus independent of pressure or collision partner within the range of pressures (69–400 torr N<sub>2</sub>, 100 torr He) investigated in this study. Variation of the excimer-laser fluence over a factor 30 also had no significant influence ( $\pm 3\%$ ) on the OH decay coefficient, showing that reactions of OH with primary and secondary products of the title reaction do not contribute significantly to its loss. A complete list of the values of  $k_4$  obtained under various experimental conditions is given in Table 2 and plotted in Arrhenius form in Fig. 4. The regression through the data points is described by  $k_4(T) = 7.5 \times 10^{-13} \exp[(-1034 \pm 40)/T]$ . The average, room-temperature (296  $\pm$  2) K rate coefficient is  $2.32 \times 10^{-14}$  cm<sup>3</sup> molecule<sup>-1</sup> s<sup>-1</sup>. The total uncertainty associated with the rate coefficients obtained (and derived from the Arrhenius expression), is dominated by uncertainty in the concentration of CH<sub>3</sub>ONO<sub>2</sub> and is estimated to be  $\sim 5\%$  (see ESI,<sup>†</sup> Section S4).




Fig. 4 Arrhenius plot of  $k_4$  versus inverse temperature. In blue:  $k_4$  from generation of OH from HONO photolysis at 351 nm. In red, the exponential fit to the data with 2 $\sigma$  confidence limit. In green:  $k_4$  from generation of OH from CH<sub>3</sub>ONO<sub>2</sub> photolysis at 248 nm. In purple:  $k_4$  from generation of OH from CH<sub>3</sub>ONO<sub>2</sub> photolysis at 248 nm considering only data at  $t > 1.7$  ms.




Fig. 5 OH formation and decay when different concentrations of CH<sub>3</sub>ONO<sub>2</sub> were photolyzed at 248 nm.



### 3.2 Generation of OH from CH<sub>3</sub>ONO<sub>2</sub> or H<sub>2</sub>O<sub>2</sub> photolysis at 248 nm (298 K)

As described above, the photolysis of methyl nitrate at 248 nm indirectly produces OH. Based on the CH<sub>3</sub>ONO<sub>2</sub> concentration, the laser fluence and the photolysis quantum yields at 248 nm,<sup>15</sup> between  $8 \times 10^8$  and  $2 \times 10^9$  molecule cm<sup>-3</sup> of OH are generated *via* R8 and R9 when [CH<sub>3</sub>ONO<sub>2</sub>] is varied from  $7.4 \times 10^{15}$  to  $2.1 \times 10^{16}$  molecule cm<sup>-3</sup>. At the same time  $1.4 \times 10^{11}$  to  $3.9 \times 10^{11}$  molecule cm<sup>-3</sup> of NO<sub>2</sub> and CH<sub>3</sub>O are formed.

Fig. 5 displays OH time profiles from the photolysis of six different concentrations of CH<sub>3</sub>ONO<sub>2</sub>. The OH decays are bi-exponential, with well separated production and loss terms. The rapid production of OH (*via* R9) is followed by a slower decay as OH is lost (mainly *via* R4). The solid lines are fits to the data according to:

$$[\text{OH}]_t = [\text{CH}_3\text{ONO}_2]_0 \frac{k'_9}{k'_4 - k'_9} (\exp(-k'_9 t) - \exp(-k'_4 t)) \quad (4)$$

where  $k'_9$  is the pseudo-first order term dominated by H-atom loss *via* its reactions with CH<sub>3</sub>ONO<sub>2</sub> (R9). In principle, values of  $k_4$  and  $k_9$  can be extracted from the data by plotting  $k'_4$  and  $k'_9$




Fig. 6 First-order rate coefficients  $k'$  for OH generation and loss derived from the bi-exponential fits (eqn (4)) to OH profiles obtained by photolysis of CH<sub>3</sub>ONO<sub>2</sub> at 248 nm and plotted against the corresponding methyl nitrate concentration [CH<sub>3</sub>ONO<sub>2</sub>] in molecule cm<sup>-3</sup>. Error bars represent  $2\sigma$  statistical uncertainty. The solid lines are linear regressions where the slopes correspond to the rate coefficients  $k_4$  (upper panel) and  $k_9$  (lower panel).

*versus* the concentration of CH<sub>3</sub>ONO<sub>2</sub> as displayed in Fig. 6. From the regression slopes we derive  $k_4(298 \text{ K}) = (1.37 \pm 0.19) \times 10^{-14}$  and  $k_9(298 \text{ K}) = (1.68 \pm 0.45) \times 10^{-13}$  cm<sup>3</sup> molecule<sup>-1</sup> s<sup>-1</sup>. The y-axis intercept of  $75 \pm 22$  s<sup>-1</sup> for the plot of  $k'_4$  *versus* [CH<sub>3</sub>ONO<sub>2</sub>] is consistent with diffusive losses of OH. The much larger intercept for the plot of  $k'_9$  *versus* [CH<sub>3</sub>ONO<sub>2</sub>] stems from H-atom diffusion and possibly reaction of H atoms with O<sub>2</sub> present as impurity.

The value of  $k_4$  thus obtained is a factor  $\approx 1.6$  lower than the average room-temperature value ( $2.32 \times 10^{-14}$  cm<sup>3</sup> molecule<sup>-1</sup> s<sup>-1</sup>) obtained in the experiments in which OH was generated by the 351 nm photolysis of HONO. A different data treatment, in which only the latter part of the bi-exponential profile ( $t > 1700$   $\mu$ s) was analysed (*i.e.* after all H atoms have been converted to OH *via* reaction with CH<sub>3</sub>ONO<sub>2</sub>) results in a rate coefficient of  $(1.65 \pm 0.10) \times 10^{-14}$  cm<sup>3</sup> molecule<sup>-1</sup> s<sup>-1</sup>, which is also lower (factor 1.4) than that obtained using HONO photolysis. As the photolysis of CH<sub>3</sub>ONO<sub>2</sub> at 248 nm results mainly in the formation of CH<sub>3</sub>O and NO<sub>2</sub> and OH reacts rapidly with both ( $k(\text{OH} + \text{NO}_2) = 3.6 \times 10^{-12}$  cm<sup>3</sup> molecule<sup>-1</sup> s<sup>-1</sup><sup>5</sup> and  $k(\text{OH} + \text{CH}_3\text{O}) = 3 \times 10^{-11}$  cm<sup>3</sup> molecule<sup>-1</sup> s<sup>-1</sup><sup>19</sup> at 298 K and 100 torr), the unavoidable presence of CH<sub>3</sub>O and NO<sub>2</sub> would be expected to result in an overestimation of the rate coefficient *via* this method. In the ESI† we use numerical simulations to examine the impact of secondary reactions and the effect of simultaneous detection of both OH and CH<sub>3</sub>O (resulting in “OH” profiles that are not perfectly bi-exponential). Clearly, the photolysis of CH<sub>3</sub>ONO<sub>2</sub> with complex secondary chemistry does not lend itself to accurate measurement of the OH-rate coefficient. As shown in Fig. 6, the linear relationship between  $k'_9$  and [CH<sub>3</sub>ONO<sub>2</sub>] enables us to derive an approximate rate coefficient ( $k_9$ ) for the OH-forming reaction of H + CH<sub>3</sub>ONO<sub>2</sub>. Our result is  $k_9(298 \text{ K}) = (1.68 \pm 0.45) \times 10^{-13}$  cm<sup>3</sup> molecule<sup>-1</sup> s<sup>-1</sup>, where the uncertainty is  $2\sigma$ , statistical only. This result is in good agreement with that reported by Talukdar *et al.*<sup>15</sup> using an almost identical method and similar analysis.

The photolysis of H<sub>2</sub>O<sub>2</sub> is commonly used as a source of prompt OH radicals in PLP-PLIF experiments. The generation of OH *via* 248 nm photolysis of H<sub>2</sub>O<sub>2</sub> in the presence of CH<sub>3</sub>ONO<sub>2</sub> does however not result in exponential OH decays. The more complex behaviour compared to that observed when using HONO at 351 nm as OH precursor is related to the photolysis of CH<sub>3</sub>ONO<sub>2</sub> at 248 nm (see above). Values of  $k'$  obtained using this method were strongly dependent on the 248 nm pulse energy and no linear relationship with the concentration of methyl nitrate was found. Clearly, as described above for experiments in the absence of H<sub>2</sub>O<sub>2</sub>, the photolysis of methyl nitrate at 248 nm precludes a straightforward analysis of the OH-profiles thus obtained.

### 3.3 Comparison of $k_4$ with literature values and the CH<sub>3</sub>ONO<sub>2</sub> atmospheric lifetime

The low rate coefficient for the reaction of OH with CH<sub>3</sub>ONO<sub>2</sub> is consistent with the reaction proceeding *via* H-atom abstraction from the methyl group, with each C–H bond contributing about  $1 \times 10^{-14}$  cm<sup>3</sup> molecule<sup>-1</sup> s<sup>-1</sup> to the rate coefficient at room



Table 3 Comparison with previous determinations of  $k_4$ 

Source	$k_4$ ( $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ )	$T$ (K)	Method
This work	$(2.32 \pm 0.04) \times 10^{-14}$ $7.5 \times 10^{-13} \times \exp[(-1034 \pm 40)/T]$	$296 \pm 2$ 223–343	PLP–PLIF
Gaffney <i>et al.</i> <sup>9</sup>	$(3.4 \pm 0.4) \times 10^{-14}$	298	DF–RF
Nielsen <i>et al.</i> <sup>22</sup>	$(3.2 \pm 0.5) \times 10^{-13}$ $8.8 \times 10^{-15} \times \exp[(1050 \pm 180)/T]$	$298 \pm 2$ 298–393	PR–RA
Talukdar <i>et al.</i> <sup>8</sup>	$(3.1 \pm 0.7) \times 10^{-13}$ $(2.36 \pm 0.16) \times 10^{-14}$	$298 \pm 2$ 298	RR PLP–PLIF
Shallcross <i>et al.</i> <sup>23</sup>	$8.2 \times 10^{-13} \times \exp[(-1020 \pm 60)/T]$ $(4.7 \pm 1.0) \times 10^{-14}$	221–414 298	DF–RF
Kerr and Stocker <sup>20</sup>	$4.1 \times 10^{-13} \times \exp[(-604 \pm 121)/T]$ $(3.8 \pm 1.0) \times 10^{-13}$	298–423 303 $\pm$ 2	RR
Kakesu <i>et al.</i> <sup>3</sup>	$(3.0 \pm 0.7) \times 10^{-14}$	307 $\pm$ 3	RR

PLP–PLIF = Pulsed laser photolysis/pulsed laser induced fluorescence; DF–RF = discharge flow/resonance fluorescence; PR–RA = pulse radiolysis/resonant absorption; RR = relative rate.

temperature. Kinetic isotope effects<sup>8</sup> confirm that H-atom abstraction is the only significant pathway for the title reaction in the temperature range representative of our atmosphere.

In order to compare our rate coefficients and their temperature dependence with literature values, we consider only our data measured using the photolysis of HONO at 351 nm, which are not biased by complex secondary processes involving photo-fragments of  $\text{CH}_3\text{ONO}_2$ . Table 3 and Fig. 7 compare the room temperature rate coefficients and Arrhenius expression obtained in this work with those reported in the literature.

**Absolute rate coefficient measurements.** When considering only those studies that used absolute methods, our room-temperature rate coefficient is essentially identical to that of Talukdar *et al.*,<sup>8</sup> but factors of 13, 1.4 and 2 lower than those reported by Nielsen *et al.*,<sup>10</sup> Gaffney *et al.*<sup>9</sup> and Shallcross *et al.*,<sup>11</sup> respectively. For slow reactions such as  $\text{OH} + \text{CH}_3\text{ONO}_2$ ,

reactive impurities and secondary reactions of OH (*e.g.* with products or photo-fragments) can bias the rate coefficients derived from absolute methods to higher values. In the study of Nielsen *et al.* (the only one to report a negative dependence of  $k_4$  on temperature), the pulse-radiolysis technique generated not only very high concentrations of OH ( $\sim 10^{13} \text{ molecule cm}^{-3}$ ) but also would have caused substantial fragmentation of  $\text{CH}_3\text{ONO}_2$  (*e.g.* to  $\text{CH}_3\text{O}$  and  $\text{NO}_2$ ). This probably explains the very large rate coefficient and anomalous temperature dependence observed.

The other temperature-dependent studies report a positive dependence on temperature, with values of  $E/R$  (in K) varying from  $-1054$  (this work) to  $-1020$  (Talukdar *et al.*<sup>8</sup>) and  $-604$  (Shallcross *et al.*<sup>11</sup>). While the two studies using the “wall-free” method of PLP–PLIF agree extremely well, the flow tube study of Shallcross *et al.* has a substantially weaker dependence on temperature. Close examination of their data set (5 data points all at  $T > 298$  K) reveals considerable scatter in  $k_4$ , which is also seen in their plots of  $k'$  versus  $[\text{CH}_3\text{ONO}_2]$ . We speculate that the difference in slope reflects changes in the wall loss rates of OH, which were determined at room temperature only. As mentioned above, absolute rate studies of slow reactions require that very pure samples are used. In this study and that of Talukdar *et al.*,<sup>8</sup> spectroscopic checks were made to exclude the presence of specific reactive impurities. In the studies of Gaffney *et al.*<sup>9</sup> and Shallcross *et al.*,<sup>11</sup> the sample purity was determined to be  $> 99\%$ , but no attempt was made to detect impurities such as  $\text{HNO}_3$ ,  $\text{NO}_2$  or  $\text{CH}_3\text{OH}$ . Perhaps more pertinent however, is the fact that the two studies that utilized *in situ* (optical absorption) measurement of  $\text{CH}_3\text{ONO}_2$  are in excellent agreement. *In situ* measurement of the concentration of the excess reactant removes uncertainty associated with calculating  $[\text{CH}_3\text{ONO}_2]$  from partial flows, pressures and variations in the mixing-ratios in storage bulbs. The fact that both this work and Talukdar *et al.*<sup>8</sup> measured absorption cross-sections for  $\text{CH}_3\text{ONO}_2$  at 213.86 nm that are in almost perfect agreement strongly suggests that the excellent agreement in the values of  $k_4$  reported by these two groups is not fortuitous, but a result of multi-diagnostics and careful execution of the experiments, leading to derivation of the true rate coefficient.




Fig. 7 Arrhenius plot of  $k_4$  versus inverse temperature. The rate coefficients and experimental conditions of the present work can be found in Table 3.



**Relative rate coefficient measurements.** In principle, the relative rate measurements (all close to room temperature) should not be biased by reactive impurities in the  $\text{CH}_3\text{ONO}_2$  sample, yet the results of the three relative-rate studies are highly divergent. The very large rate coefficients measured by Nielsen *et al.*<sup>10</sup> and Kerr and Stocker<sup>20</sup> can be traced back to an inappropriate choice of reference reactants, which ideally should react with rate coefficients that are similar to the target molecule. The use of 2-methylpropane ( $k_{\text{OH}+2\text{-methylpropane}} = 2.1 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1.5}$ ) and ethene ( $k_{\text{OH}+\text{ethene}} = 7.8 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  at 1 bar<sup>5</sup>) by Nielsen *et al.*<sup>10</sup> and Kerr and Stocker,<sup>20</sup> respectively, would have made relative depletion rates of the reference and target molecule difficult to measure and also prone to other loss processes of  $\text{CH}_3\text{ONO}_2$ . In the relative rate study of Kakesu *et al.*,<sup>3</sup> the more slowly reacting reference trace-gases, methane ( $k_{\text{OH}+\text{methane}} = 6.4 \times 10^{-15} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1.5}$ ) and ethane ( $k_{\text{OH}+\text{ethane}} = 2.4 \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1.5}$ ) were used. Those authors derived a rate coefficient at 307 K of  $(3.0 \pm 0.7) \times 10^{-14} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ . Within combined uncertainty, this value agrees with the absolute rate coefficient derived in the present study for the same temperature: of  $k_4(307 \text{ K}) = (2.7 \pm 0.4) \times 10^{-14} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ .

**Atmospheric lifetime of  $\text{CH}_3\text{ONO}_2$  with respect to reaction with OH.** As the rate coefficient for the title reaction is independent of pressure under all conditions of the Earth's lower atmosphere (*i.e.* the troposphere and lower stratosphere) we use the Arrhenius expression derived from our experiments using HONO as OH source:  $k_4(T) = 7.5 \times 10^{-13} \exp(-1034/T)$  to calculate the lifetime of  $\text{CH}_3\text{ONO}_2$  with respect to OH loss. For this we assume an altitude independent, global mean OH concentration of  $1 \times 10^6 \text{ molecule cm}^{-3}$ .<sup>21</sup> The lifetime of  $\text{CH}_3\text{ONO}_2$  w.r.t. OH reaction at different altitudes is plotted in Fig. S5 (ESI<sup>†</sup>) and varies from *circa* 500 days at ground level to > 1000 days at 10 km. Photolysis rate coefficients for  $\text{CH}_3\text{ONO}_2$  vary from  $\sim 3 \times 10^{-7} \text{ s}^{-1}$  to  $5 \times 10^{-7} \text{ s}^{-1}$  in the boundary layer and lower stratosphere,<sup>15</sup> respectively, resulting in lifetimes with respect to photodissociation of close to 1 month. Our data thus confirms the conclusion of Talukdar *et al.*<sup>15</sup> showing that reaction with OH is a minor sink of  $\text{CH}_3\text{ONO}_2$  in the atmosphere. This is in contrast to the conclusions of Nielsen *et al.*<sup>10</sup> and Kerr and Stocker<sup>20</sup> who (based on too-high rate coefficients) suggested that reaction with OH is important.

## 4 Conclusions

Motivated by greatly divergent reports of the rate coefficient for the reaction of OH with  $\text{CH}_3\text{ONO}_2$ , we have determined the rate coefficient using pulsed laser photolysis-laser induced fluorescence. High-quality kinetic data were obtained when measures were taken to avoid the photolysis of  $\text{CH}_3\text{ONO}_2$ . Our result, a positive dependence of the rate coefficient on temperature described by  $k_4(T) = 7.5 \times 10^{-13} \exp[(-1034 \pm 40)/T]$ , is in close agreement with that reported by Talukdar *et al.*<sup>8</sup> Both studies used *in situ* measurement of  $[\text{CH}_3\text{ONO}_2]$  using

independently determined absorption cross-sections that agreed with 1.2%. The divergent results obtained by other groups are rationalised in terms of choice of OH source or reference reactant and data corrections.

## Author contributions

This work was carried out as part-fulfilment of CF's master's degree in chemistry at the Julius-Maximilians-University, Würzburg, Germany. CF and FB conducted the experiments, CF evaluated the data and prepared the manuscript. JC conceptualized the experiment, JC and JL contributed to writing the manuscript.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

Open Access funding provided by the Max Planck Society.

## Notes and references

- 1 J. M. Roberts, *Atmos. Environ., Part A*, 1990, **24**, 243.
- 2 E. Atlas, *Nature*, 1988, **331**, 426.
- 3 M. Kakesu, H. Bandow, N. Takenaka, Y. Maeda and N. Washida, *Int. J. Chem. Kinet.*, 1997, **29**, 933.
- 4 N. L. Ng, S. S. Brown, A. T. Archibald, E. Atlas, R. C. Cohen, J. N. Crowley, D. A. Day, N. M. Donahue, J. L. Fry, H. Fuchs, R. J. Griffin, M. I. Guzman, H. Herrmann, A. Hodzic, Y. Iinuma, J. L. Jimenez, A. Kiendler-Scharr, B. H. Lee, D. J. Luecken, J. Mao, R. McLaren, A. Mutzel, H. D. Osthoff, B. Ouyang, B. Picquet-Varrault, U. Platt, H. O. T. Pye, Y. Rudich, R. H. Schwantes, M. Shiraiwa, J. Stutz, J. A. Thornton, A. Tilgner, B. J. Williams and R. A. Zaveri, *Atmos. Chem. Phys.*, 2017, **17**, 2103.
- 5 IUPAC, Task Group on Atmospheric Chemical Kinetic Data Evaluation, (Ammann, M., Cox, R.A., Crowley, J.N., Herrmann, H., Jenkin, M.E., McNeill, V.F., Mellouki, A., Rossi, M. J., Troe, J. and Wallington, T. J.). Last access Sept. 2023, (<https://iupac.aeris-data.fr/>).
- 6 N. Butkovskaya, A. Kukui and G. Le Bras, *J. Phys. Chem. A*, 2012, **116**, 5972.
- 7 F. Flocke, E. Atlas, S. Madronich, S. M. Schauffler, K. Aikin, J. J. Margitan and T. P. Bui, *Geophys. Res. Lett.*, 1998, **25**, 1891.
- 8 R. K. Talukdar, S. C. Herndon, J. B. Burkholder, J. M. Roberts and A. R. Ravishankara, *J. Chem. Soc., Faraday Trans.*, 1997, **93**, 2787.
- 9 J. S. Gaffney, R. Fajer, G. I. Senum and J. H. Lee, *Int. J. Chem. Kinet.*, 1986, **18**, 399.
- 10 O. J. Nielsen, H. W. Sidebottom, M. Donlon and J. Treacy, *Chem. Phys. Lett.*, 1991, **178**, 163.



- 11 D. E. Shallcross, P. Biggs, C. E. Canosa-Mas, K. C. Clemitshaw, M. G. Harrison, M. Reyes López Alañón, J. A. Pyle, A. Vipond and R. P. Wayne, *J. Chem. Soc., Faraday Trans.*, 1997, **93**, 2807.
- 12 M. Wollenhaupt, S. A. Carl, A. Horowitz and J. N. Crowley, *J. Phys. Chem.*, 2000, **104**, 2695.
- 13 W. D. Taylor, T. D. Allston, M. J. Moscato, G. B. Fazekas, R. Kozlowski and G. A. Takacs, *Int. J. Chem. Kinet.*, 1980, **12**, 231.
- 14 J. G. Calvert and J. N. J. Pitts, *Photochemistry*, J. Wiley, New York, 1966.
- 15 K. R. Talukdar, J. B. Burkholder, M. Hunter, M. K. Gilles, J. M. Roberts and A. R. Ravishankara, *J. Chem. Soc., Faraday Trans.*, 1997, **93**, 2797.
- 16 C. B. M. Groß, T. J. Dillon, G. Schuster, J. Lelieveld and J. N. Crowley, *J. Phys. Chem. A*, 2014, **118**, 974.
- 17 G. L. Vaghjiani and A. R. Ravishankara, *J. Chem. Phys.*, 1990, **92**, 996.
- 18 A. P. Black and F. H. Babers, *Org. Synth.*, 1939, **19**, 64.
- 19 W. Tsang and R. F. Hampson, *J. Phys. Chem. Ref. Data*, 1986, **15**, 1087.
- 20 J. A. Kerr and D. W. Stocker, *J. Atmos. Chem.*, 1986, **4**, 253.
- 21 J. Lelieveld, S. Gromov, A. Pozzer and D. Taraborrelli, *Atmos. Chem. Phys.*, 2016, **16**, 12477.
- 22 O. J. Nielsen, H. W. Sidebottom, M. Donlon and J. Treacy, *Chem. Phys. Lett.*, 1991, **178**, 163.
- 23 D. Shallcross, C. Canosa-Mas, K. Clemitshaw, M. Harrison, J. Pyle and R. Wayne, *J. Chem. Soc., Faraday Trans.*, 1997, **93**, 2807.

