







 Cite this: *Phys. Chem. Chem. Phys.*,
 2024, 26, 25461

A multiphoton ionisation photoelectron imaging study of thiophene†

 Joseph J. Broughton,  Sarbani Patra,  Michael A. Parkes, 
 Graham A. Worth  and Helen H. Fielding *

Thiophene is a prototype for the excited state photophysics that lies at the heart of many technologies within the field of organic electronics. Here, we report a multiphoton ionisation photoelectron imaging study of gas-phase thiophene using a range of photon energies to excite transitions from the ground electronic state to the first two electronically excited singlet states, from the onset of absorption to the absorption maximum. Analysis of the photoelectron spectra and angular distributions reveal features arising from direct photoionisation from the ground electronic state, and resonance-enhanced photoionisation via the electronically excited singlet states. The first two ionisation energies from the ground electronic state were confirmed to be 8.8 eV (adiabatic) and 9.6 eV (vertical). The ionisation energies from the first two electronically excited singlet states were found to be 3.7 eV (adiabatic) and 4.4 eV (vertical).

 Received 22nd June 2024,
 Accepted 19th September 2024

DOI: 10.1039/d4cp02504k

rsc.li/pccp

1 Introduction

Thiophene is a small, heteroaromatic molecule that is a ubiquitous building block in many photonic materials, such as organic photovoltaics^{1–3} and non-linear optical devices.^{4,5} Although there have been many attempts to elucidate both the electronic structure and the excited-state dynamics of gas-phase thiophene,^{6–33} its UV photoabsorption and photoionisation are still not understood fully.

One of the difficulties in understanding the photoabsorption spectrum arises as a result of the first two electronically excited singlet states, S_1 ($\pi\pi^*$) and S_2 ($\pi\pi^*$), hereafter referred to as S_1 and S_2 , lying very close together in energy and having similar oscillator strengths. Theoretical studies by Holland *et al.*²⁶ and Prlj *et al.*²⁹ have highlighted how the precise energies and ordering of these states are highly dependent on the method and the basis sets used. Although there is still not a clear consensus on the origin of structure in the UV photoabsorption of thiophene, all assignments reported in the literature do agree that the intense feature at 5.157 eV is the origin of S_1 ^{6,8,10,11,13,14,17,22,26,33} (Fig. 1), with the peaks below the origin being variously attributed to hot bands, combination bands, or vibronic coupling between S_1 and S_2 .^{8,11,13}

There have been two recent vacuum ultraviolet (VUV) absorption spectroscopy studies of gas-phase thiophene.^{26,33}




Fig. 1 Left: Photoabsorption spectrum of thiophene recorded by Holland *et al.*²⁶ Right: Schematic energy level diagram for thiophene showing the vertical excitation energies of S_1 , S_2 , and S_3 , and the vertical ionisation energies to D_0 and D_1 continua. The adiabatic excitation energy (AEE) for S_1 is taken from measurements by Holland *et al.*,²⁶ vertical excitation energies (VEE) for S_1 and S_2 are taken from magnetic circular dichroism measurements from Hakansson *et al.*,³⁴ the AEE of S_2 is taken from DFT/MRCI calculations by Salzmann *et al.*,²³ the S_3 VEE is from Holland *et al.*²⁶ The ionisation energies D_0 and D_1 are taken from Weinkauff *et al.*²² The coloured arrows signify the different wavelengths used for one-colour measurements in the work reported here.

Both measurements found several progressions which were assigned to the ν_6 vibrational mode, corresponding to a CH in-plane bending vibration, with an average peak spacing of ~ 120 meV. This supported earlier work by Palmer *et al.*,¹⁷ in which the 0–0 transition was determined to be 5.157 eV, and structure below this transition was ascribed to hot bands and S_1 – S_2 vibronic coupling.

 Department of Chemistry, University College London, 20 Gordon Street, London
 WC1H 0AJ, UK. E-mail: h.h.fielding@ucl.ac.uk

 † Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4cp02504k>


Table 1 S_0 - D_0 and S_0 - D_1 vertical ionisation energies (VIEs) and adiabatic ionisation energies (AIEs), in eV, recorded using a range of ionisation and PES methods: ionisation using HeI or Ly- α radiation with a hemispherical electron energy analyser (HEA); Penning ionisation electron spectra (PIES) with He* (2^3S) and a hemispherical analyser; zero-kinetic energy (ZEKE) PES; time-resolved $1 + 1'$ PES (TRPES) with a magnetic bottle time-of-flight (TOF) (pump-probe delay = 50 fs); femtosecond $1 + 1$ multiphoton ionisation (MPI) with velocity-map imaging (VMI)

D_0	AIE/VIE	D_1	AIE/VIE	Method	Ref.
8.87 ± 0.05	AIE ^a	9.49 ± 0.1	VIE	HeI + HEA	Eland ³⁶
8.80 ± 0.05	AIE ^a	9.44 ± 0.05	AIE ^a	HeI + HEA	Baker <i>et al.</i> ⁷
8.872 ± 0.015	AIE ^a	9.52 ± 0.015	VIE	Ly- α + HEA	Derrick <i>et al.</i> ³⁷
8.90 ± 0.015	VIE	9.50 ± 0.015	VIE	HeI + HEA	Clark <i>et al.</i> ⁹
8.85 ± 0.05	VIE	9.49 ± 0.05	VIE	HeI + HEA	Klasinc <i>et al.</i> ¹²
8.96 ± 0.25	VIE	9.58 ± 0.25	VIE	He* PIES + HEA	Kishimoto <i>et al.</i> ³⁸
8.8742 ± 0.0002	AIE	—	—	ZEKE	Yang <i>et al.</i> ³⁹
8.87 ± 0.01	AIE	9.55 ± 0.01	VIE	TRPES + TOF	Weinkauf <i>et al.</i> ²²
8.8 ± 0.1	AIE	9.6 ± 0.1	VIE	MPI + VMI	This work

^a Adiabatic and vertical ionisation energies not distinguished.

Another way to probe electronic structure is to use photoelectron spectroscopy to determine electron binding energies, $eBE = h\nu - eKE$, where $h\nu$ is the photon energy and eKE is the measured electron kinetic energy. Ionisation energies of thiophene determined from such measurements are presented in Table 1. Measuring the photoelectron angular distribution (PAD) as well as the eKE provides additional information about the molecular orbital from which the electron is removed. For a two-photon ionisation process, the angular distribution $I(\theta) \propto 1 + \beta_2 P_2(\cos \theta) + \beta_4 P_4(\cos \theta)$, where $I(\theta)$ is the probability of photoelectron emission at angle θ , defined as the angle between the laser polarisation and the velocity vector of the photoelectron, β_2 and β_4 are the photoelectron anisotropy parameters, and $P_2(\cos \theta)$ and $P_4(\cos \theta)$ are the second and fourth-order Legendre polynomials in $\cos \theta$.³⁵ The two limiting values of β_2 are +2 and -1, corresponding to photoelectron emission predominantly parallel and perpendicular to the electric field vector of the laser, respectively. β_4 represents the alignment of the excited state and, generally, positive and negative values of β_4 correspond to the excited state being aligned parallel or perpendicular to the electric field vector of the laser, respectively.

Multiphoton ionisation (MPI) photoelectron spectroscopy studies of gas-phase thiophene using UV laser pulses have been reported by two groups.^{22,32} However, there has only been one photoelectron imaging study;³² this was a time-resolved study focussing on relaxation dynamics, and PADs were not reported. Here, we present a systematic one-colour MPI photoelectron imaging study of thiophene, employing wavelengths ranging from 243 nm, exciting just below the S_0 - S_1 0-0 transition, to 234 nm, which corresponds to the S_0 - S_1 VEE (Fig. 1), from which we gain further insight into the electronic structure!

2. Methods

2.1. Multiphoton photoelectron imaging

A molecular beam of thiophene was produced by passing 1 bar He carrier gas through liquid thiophene and expanding it through a 300 μm pulsed Amsterdam piezovalve^{40,41} operating at a repetition rate of 1 kHz and with a pulse duration of 8 μs .

The molecular beam was collimated by a 3 mm skimmer before passing into the interaction region of our velocity map imaging spectrometer that has been described before.⁴²⁻⁵¹ The molecular beam was intersected by femtosecond laser pulses with wavelengths in the range 234–243 nm, generated by frequency-doubling the sum frequency of the signal of the output of an optical parametric amplifier (TOPAS-Prime) pumped by a titanium: sapphire regenerative amplifier (Coherent Astrella-HE). To prevent multiphoton excitation, pulse energies were attenuated to $\approx 1 \mu\text{J}$ per pulse using a variable neutral density filter. The $1/e^2$ pulse duration was measured as 300 fs at 243 nm and spectral full-width at half maxima were $\sim 1 \text{ nm}$ (0.02 eV). Photoelectron images were recorded for 1200 s at each wavelength. Background images (without thiophene) were also recorded for 1200 s and subtracted from the photoelectron images. Photoelectron spectra were recovered from the background-subtracted data using the pBASEX image inversion algorithm.⁵² The energy scale was calibrated by recording $2 + 1$ resonance-enhanced MPI spectra of Xe at 249.63 nm.⁵³ The resolution was $\Delta E/E \leq 3\%$.

2.2. *Ab initio* calculations

The complete active space second-order perturbation theory (CASPT2)⁵⁴ method with the 6-31G* basis set was used for the computation of the ground and electronically excited states of thiophene. The ground electronic state has C_{2v} symmetry and the active space consisted of a 10-electron, 9-orbital space including all the p-orbitals of thiophene and the two pairs of σ -orbitals (σ/σ^*) for both C-S bonds, which can correctly account for the two equivalent C-S bonds, either of which might undergo cleavage after photoexcitation. The same theoretical description was employed by Schnappinger *et al.*³¹ to study the relaxation of photoexcited thiophene using surface hopping including arbitrary couplings (SHARC).⁵⁵ Calculations were performed using the MOLPRO 2022 program.^{56,57}

A number of other methods were also used to support the characterisation: TDDFT/B3LYP, ADC(2) and EOM-CCSD. These calculations used a 6-311+G** basis set with the QChem v5.4 program.⁵⁸ The calculated excitation energies of the four singlet states matching the CASPT2 calculated states, and the first four triplet states, are listed in Table 2, along with the known singlet state experimental values. All methods agree that the



Table 2 Calculated excitation and ionisation energies using various quantum chemistry methods. See text for details. Experimental values for the singlet states are vertical transition energies taken from ref. 26 experimental values for D_0 and D_1 ionisation are given in Table 1

State	Character	Symm.	TDDFT/ B3LYP	ADC(2)	EOM- CCSD	CASPT2	Expt.
S_1	$\pi_2 \rightarrow \pi_1^*$	A_1	5.865	5.773	5.842	5.641	5.64
S_2	$\pi_1 \rightarrow \pi_1^*$	B_2	5.915	6.184	6.320	6.070	5.97
S_3	$\pi_1 \rightarrow \sigma_1^*$	B_1	6.126	6.362	6.403	6.203	6.17
S_4	$\pi_2 \rightarrow \sigma_1^*$	A_2	6.022	6.272	6.357	6.249	6.33
T_1	$\pi_1 \rightarrow \pi_1^*$	B_2	3.778	4.031	3.866	3.788	
T_2	$\pi_2 \rightarrow \pi_1^*$	A_1	4.561	4.906	4.825	4.813	
T_3	$\pi_2 \rightarrow \sigma_1^*$	B_1	5.618	6.196	6.206	5.849	
T_4	$\pi_1 \rightarrow \sigma_1^*$	A_2	5.847	6.201	6.113	5.921	
D_0	π_1^1	B_2	EOM-IP- CCSD 8.896	EPT 8.967			
D_1	π_2^1	B_1	9.132	9.276			

singlet states are close in energy, in the range 5.7–6.5 eV. The S_3 and S_4 states are found to be near degenerate, with a change in order for all methods compared to the CASPT2. The triplets are spread more widely, but T_1 and T_2 are both predicted to lie below the S_1 state. Again the character of the close-lying T_3 and T_4 states changes order for all methods compared to CASPT2. TDDFT also reported a singlet state at 5.982 eV and a triplet at 5.805 eV involving Rydberg orbitals.

The VIEs were calculated using electron propagator theory (EPT)^{59–61} in Gaussian 09⁶² with the 6-311++G(3df,3pd) basis set, and EOM-IP-CCSD using QChem v5.4 with the 6-311+G** basis set. Dyson orbital norms (ionisation cross-sections) and β parameters (PADs) were calculated for transitions from S_n to D_0 and D_1 for $n = 1–4$ using the EOM-IP-CCSD/6-311++G** method in ezDyson.⁶³ The calculated D_0 and D_1 energies are also listed in Table 2 and agree well with the experimental values of Table 1. The next two cation states are found to be much higher in energy at around 12 eV and are ignored.

3. Results and discussion

3.1. Electronic structure

The electronic configurations and CASPT2(10,9)/6-31G* calculated VEEs of the first four singlet excited states of thiophene are presented in Fig. 2, along with the electronic configurations and EPT/6-311G(3df,3pd) calculated VIEs of the first two doublet states of the thiophene cation. To a first approximation, we can use Koopmans' theorem to determine ionisation propensities. From the electronic configurations shown in Fig. 2 it is clear that the S_0 state can ionise to D_0 and D_1 , that S_1 and S_4 are most likely to ionise to D_1 , and S_2 and S_3 are most likely to ionise to D_0 . These ionisation propensities are supported by calculations of photoionisation cross-sections for one photon ionisation from $S_0–S_4$ to the D_0 and D_1 states of the thiophene cation (Fig. 3).

3.2. Photoelectron spectra

The 243–234 nm one-colour MPI photoelectron spectra of thiophene are presented as a heat map plotted as a function




Fig. 2 Left: Electronic configurations of the five lowest singlet electronic states of thiophene and two lowest doublet electronic states of the corresponding radical cation. Numbers in brackets refer to CASPT2(10,9)/6-31G* calculated vertical excited energies and EPT/6-311++G(3df,3pd) vertical ionisation energies, in eV and relative to S_0 . Vertical arrows highlight transitions from S_0 to S_1 and S_2 . Dashed arrows highlight most probable ionisation processes based on Koopmans' correlations. Right: CASPT2(10,9)/6-31G* natural orbitals.




Fig. 3 One-photon photoionisation cross-sections from $S_0–S_4$ to the D_0 (left) and D_1 (right) states of thiophene. Dyson orbital norms were calculated using the EOM-IP-CCSD/6-311++G** method.

of eKE and photon energy (Fig. 4). The individual spectra are also plotted as a function of eKE in Fig. 5(a). Fig. 4 allows us to determine whether features in the spectra are a result of non-resonant or resonance-enhanced MPI processes. For non-resonant two-photon ionisation, electrons are emitted with $eKE = 2h\nu - IE$, where $h\nu$ is the photon energy and IE is the ionisation energy. In Fig. 4, the solid black lines highlight peaks in the photoelectron spectra whose eKEs increase at twice the rate of the photon energy, and represent non-resonant ionisation from S_0 to the two doublet cation states, D_0 and D_1 . Following resonant photoexcitation of an excited electronic state S_m , with excess vibrational energy $E_{\text{vib}} = h\nu - AEE(S_m)$ where $AEE(S_m)$ is the adiabatic excitation energy of S_m , ionisation by a second photon will result in the emission of






Fig. 4 One-colour MPI photoelectron spectra of thiophene plotted as a function of eKE and photon energy. Arrows mark the photon energies employed. The individual spectra were linearly interpolated and normalised to the most intense feature at 0.70 eV eKE, which corresponds to the S_0 - D_1 VIE. Direct ionisation to D_0 and D_1 were determined as 8.8 ± 0.1 eV and 9.6 ± 0.1 eV, as shown by the solid lines of gradient = 2. The most probable indirect ionisation processes are marked with dotted and dashed lines with a gradient of 1. These show ionisation via S_2 - D_0 and S_1 - D_1 , respectively (see text).

electrons with $eKE \leq 2h\nu - AIE(D_m) - E_{vib} = h\nu - [AIE(D_m) - AEE(S_n)]$, where $AIE(D_m)$ is the adiabatic ionisation energy of the cation doublet state D_m . In Fig. 4, the dotted and dashed black lines highlight peaks in the photoelectron spectra whose eKEs increase at the same rate as the photon energy, and represent resonance-enhanced ionisation to the two doublet cation states, D_0 and D_1 , respectively. The y-intercepts of the solid, dotted and dashed lines were obtained from plots of the MPI photoelectron spectra as functions of two- or one-photon electron binding energy, $eBE = nh\nu - eKE$, where $n = 2$ or 1 (Fig. 5).

In Fig. 5(b), the photoelectron spectra are plotted as a function of two-photon binding energy. There are two ionisation energies that can be identified in these spectra. The small bump at 8.8 ± 0.1 eV on the rising edge of the lowest energy peak can be assigned as the S_0 - D_0 AIE. This is in excellent agreement with the AIE obtained from PES^{7,9,12,22,36–38,64} and zero-kinetic energy (ZEKE) PES³⁹ measurements, several of which determined that the AIE was equivalent to the VIE (Table 1). The peak at 9.6 ± 0.1 eV is assigned as the S_0 - D_1 VIE, and this is also in good agreement with values reported previously.^{7,9,12,22,36–38,64} Note that the errors attributed to our assignments are derived from the uncertainty in the position rather than the energy resolution of our spectrometer.

It can be seen in Fig. 5(b) that the peak attributed to S_0 - D_1 does not have any clear vibrational structure (experimental resolution ~ 0.05 eV), in agreement with observations of Derrick *et al.* (experimental resolution ~ 0.02 eV).³⁷ Also, the peak broadens significantly as photon energy increases, losing the sharpness associated with non-resonant photoionisation from




Fig. 5 (a) Plot of photoelectron counts versus eKE following 1 + 1 photoionisation of thiophene for 234–243 nm. (b) Plot of photoelectron counts versus two-photon eBE following 1 + 1 photoionisation of thiophene for 234–243 nm. The solid and dashed lines show $D_0 = 8.8 \pm 0.1$ eV and $D_1 = 9.6 \pm 0.1$ eV. (c) Plot of photoelectron counts versus one-photon eBE following 1 + 1 ionisation of thiophene for 234–243 nm. The comb denotes ionisation from S_n - D_0 and S_n - D_1 using the S_1 AEE from Holland *et al.*,²⁶ the calculated S_2 AEE from Salzman *et al.*²³ and the ionisation energies from the work reported here. The solid and dashed vertical lines denote adiabatic ionisation from S_1 and S_2 , respectively. The individual spectra were normalised to their integrated area.



S_0 at photon energies lower than the S_0 - S_1 VEE. Although Rydberg states have been observed in the range 9.4–11.8 eV,²⁶ the structured features in our 5.10–5.19 eV photoelectron spectra do not remain at constant eKE with increasing photon energy (Fig. 5(a)) so cannot be attributed to autoionising Rydberg states. Also, the 5.20–5.30 eV photoelectron spectra do not have the identical profiles expected for an autoionising Rydberg states. This leaves two possible explanations for our experimental observations. First, as the photon energy increases above 5.157 eV, S_1 is excited. From Koopmans' correlations (Fig. 2) and calculation photoionisation cross-sections (Fig. 3), the S_1 - D_1 ionisation pathway is expected to dominate over S_1 - D_0 . There may be a large geometric change between S_1 and D_1 , which would result in a broader photoelectron spectrum. Moreover, the D_1 feature shifts to higher binding energies with increasing photon energy, which is a signature of resonance-enhanced ionisation from S_1 beginning to compete with, or dominate, non-resonant ionisation from S_0 . Second, D_1 could have a short lifetime. Calculations reported by Trofimov *et al.* identified strong vibronic coupling between D_1 and D_0 and a conical intersection between them,¹⁶ which would account for a short D_1 lifetime and the broad, vibrationally-unresolved photoelectron spectrum.

In Fig. 5(c), the photoelectron spectra are plotted as a function of one-photon binding energy. The comb marks S_n - D_m AIEs determined using AEEs from Holland *et al.*,²⁶ and Salzmann *et al.*,²³ and the D_0 and the D_1 IEs from this work. It is clear that S_1 - D_1 is the dominant resonance-enhanced ionisation pathway as the photoelectron counts are highest around the calculated S_1 - D_1 VIE (4.4 ± 0.1 eV). Nonetheless, there also appears to be a significant contribution from the S_1 - D_0 ionisation channel calculated at 3.7 ± 0.1 eV. For photon energies above the S_0 - S_1 AEE (5.157 eV), what looks like an isobestic point begins to appear. It lies at 3.7 ± 0.1 eV, which is very close to the calculated S_2 - D_0 AIE (3.8 ± 0.1 eV) and is illustrative of an increasing contribution from the population of, and subsequent ionisation from, S_2 . Based on Koopmans' correlations and photoionisation calculations, the S_2 - D_0 ionisation channel is expected to dominate over S_2 - D_1 ionisation, although it is difficult to determine whether there is any significant contribution from S_2 - D_1 ionisation from our spectra as they are very congested in the region where we would expect this channel to appear (4.5 ± 0.1 eV). Fig. 3 shows that the cross-sections for the dominant S_2 - D_0 and S_1 - D_1 ionisation channels are very similar. Population of, and ionisation from, S_2 is consistent with calculations performed by Köppel *et al.* which reported a rapid population of S_2 following photoexcitation of S_1 . At photon energies close to the S_0 - S_2 VEE (5.64 eV),³⁴ it is likely that the S_2 - D_0 and S_2 - D_1 channels would make a larger contribution to the measured photoelectron intensity due to the direct population of S_2 .

3.3. Photoelectron angular distributions

In Fig. 6(a) the measured anisotropy parameter β_2 is plotted as a function of eKE and photon energy. At higher eKEs, between the solid lines marking the upper limits of direct and indirect




Fig. 6 Plots of (a) β_2 and (b) β_4 parameters as a function of eKE and photon energy following 1 + 1 photoionisation of thiophene. The plots have been smoothed using a five-point average for eKE and β parameters. Arrows mark the photon energies employed. The solid lines with gradient = 2 mark the high eKE limits of direct photoionisation to D_0 and D_1 . The dotted and dashed lines mark the high eKE limits of indirect S_2 - D_0 and S_1 - D_1 photoionisation processes, respectively. The grey shaded region is energetically inaccessible via two-photon ionisation.

photoionisation to D_0 and D_1 , there is weak anisotropy with $\beta_2 \approx -0.05$. At lower eKEs, below the lines marking the upper limits of direct and indirect photoionisation to the D_1 continuum, the anisotropy changes sign and becomes weakly positive, with $\beta_2 \approx +0.05$. In Fig. 6(b) the measured β_4 anisotropy parameter, is plotted as a function of photon energy and eKE. At higher eKEs, $\beta_4 < 0$, whereas for $eKE < 0.4$, $\beta_4 \approx +0.1$.

The calculated one-photon β_2 parameters are plotted in Fig. 7. In the energy region of interest for our experiments, $\beta_2 < 0$ for one-photon photoionisation from S_0 , S_1 and S_2 , to D_0 and D_1 . The measured β_2 parameter is a superposition of direct and indirect ionisation processes and although it is negative for higher eKEs, it is positive for lower eKEs. This difference between the calculated and measured patterns could be due to the β_2 parameter for two-photon non-resonant ionisation from S_0 having the opposite sign to our calculated one-photon






Fig. 7 Plots of β_2 anisotropy parameters as a function of eKE following one-photon photoionisation from S_0 – S_4 to D_0 (left) and D_1 (right) continua, calculated using the EOM-IP-CCSD/6-311++G** method.

β_2 parameters for lower eKEs, or it could be the result of structural changes occurring on the timescale of our measurements. Interestingly, $\beta_2 > 0$ for photoionisation from S_3 and S_4 . Ionisation from S_0 – S_2 involve removing electrons from $\pi\pi^*$ molecular orbitals whereas ionisation from S_3 and S_4 involve removing electrons from $\pi\sigma^*$ molecular orbitals. (Fig. 2) This difference in orbital character could explain the difference in sign of the β_2 . Thus, another explanation could be that the higher lying S_3 and S_4 states are accessed on the timescale of the measurement, which would be consistent with ultrafast ring-opening. This is indeed found to be the case in our recent theoretical study of the photo-excited dynamics in the vibronically coupled S_1 – S_4 states of thiophene.⁶⁵

Thus, time-resolved photoelectron angular imaging experiments, with supporting calculations of the PADs, could prove valuable for investigating the electronic relaxation pathways following photoexcitation.

4. Conclusions

We have presented 243–234 nm multiphoton photoelectron imaging measurements of thiophene. Plots of the photoelectron spectra as a function of two-photon binding energy allowed us to identify spectral features with binding energies of 8.8 ± 0.1 eV and 9.6 ± 0.1 eV, corresponding to the S_0 – D_0 AIE and S_0 – D_1 VIE, respectively. These are in excellent agreement with ionisation energies determined using other methods.^{7,9,12,22,36–38,64} Plots of the photoelectron spectra as a function of one-photon binding energy revealed peaks around the S_1 – D_1 VIE (4.4 ± 0.1 eV) and S_2 – D_0 AIE (3.8 ± 0.1 eV). For spectra recorded with photon energies above the S_0 – S_1 AEE (5.157 eV), an isobestic point was observed at $h\nu$ -eKE = 3.7 ± 0.1 eV illustrating an increasing contribution from S_2 – D_0 ionisation, compared with S_1 – D_1 ionisation, with increasing photon energy.

Plots of β_2 and β_4 anisotropy parameters as a function of eKE and photon energy suggest that time-resolved photoelectron imaging measurements could provide a sensitive probe of electronic relaxation following UV photoexcitation of thiophene. This work serves as a benchmark for future time-resolved studies of thiophene and more complex molecular systems

containing thiophene as a building block that are ubiquitous in many photonic materials.

Author contributions

The project was conceived and supervised by H. H. F. The experimental data were recorded by J. J. B., with technical assistance from M. A. P., and processed, analysed and interpreted by J. J. B. Calculations were undertaken J. J. B., and by S. P. and M. A. P. under the guidance of G. A. W. The manuscript was written by J. J. B. and H. H. F., with contributions from S. P. and G. A. W. All authors read and approved the final version of the manuscript.

Data availability

Data in this article will be made available in the UCL Research Data Repository at <https://www.ucl.ac.uk/library/open-science-researchsupport/research-data-management/ucl-research-data-repository>.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The work was supported by EPSRC grants EP/V026690/1 and EP/L015862/1. The experiments were performed using the Ultrafast Laser Facility in the Department of Chemistry at UCL (EPSRC EP/T019182/1), with technical support from Dr Julia Davies. The authors thank Roisin Tapley for carrying out preliminary calculations of one-photon photoionisation cross-sections and β_2 anisotropy calculations.

Notes and references

- 1 F. Zhang, D. Wu, Y. Xu and X. Feng, *J. Mater. Chem.*, 2011, **21**, 17590–17600.
- 2 P. Kumaresan, S. Vegiraju, Y. Ezhumalai, S. L. Yau, C. Kim, W. H. Lee and M. C. Chen, *Polymers*, 2014, **6**, 2645–2669.
- 3 T. Taniguchi, K. Fukui, R. Asahi, Y. Urabe, A. Ikemoto, J. Nakamoto, Y. Inada, T. Yamao and S. Hotta, *Synth. Met.*, 2017, **227**, 156–162.
- 4 G. Kirsch, D. Prim, F. Leising and G. Mignani, *J. Heterocycl. Chem.*, 1994, **31**, 1005–1009.
- 5 H. L. Andriampanarivo, M. Köhler, J. L. Gejo, T. Betzwieser, B. C. Poon, P. L. Yue, S. D. Ravelomanantsoa and A. M. Braun, *Photochem. Photobiol. Sci.*, 2015, **14**, 1013–1024.
- 6 G. Horváth and A. I. Kiss, *Spectrochim. Acta, Part A*, 1967, **23**, 921–924.
- 7 A. D. Baker, D. Betteridge, N. R. Kemp and R. E. Kirby, *Anal. Chem.*, 1971, **43**, 375–381.
- 8 G. D. Lonardo, G. Galloni, A. Trombetti and C. Zauli, *J. Chem. Soc., Faraday Trans.*, 1972, **68**, 2009–2016.



- 9 P. A. Clark, R. Gleitter and E. Heilbronner, *Tetrahedron*, 1973, **29**, 3085–3089.
- 10 W. M. Flicker, O. A. Mosher and A. Kuppermann, *Chem. Phys. Lett.*, 1976, **38**, 489–492.
- 11 G. Varsanyi, L. Nyulaszi, T. Veszpremi and T. Narisawa, *J. Chem. Soc. Perkin Trans.*, 1982, 761–765.
- 12 B. L. Klasinc, A. Sabljic, G. Kluge, J. Rieger, K. Marx Universität Leipzig, S. Chemie and G. D. Republic, *J. Chem. Soc., Perkin Trans. 2*, 1982, 539–543.
- 13 E. J. Beiting, K. J. Zeringue and R. E. Stickel, *Spectrochim. Acta, Part A*, 1985, **41**, 1413–1418.
- 14 L. Nyulászi and T. Veszprémi, *J. Mol. Struct.*, 1986, **140**, 253–259.
- 15 F. Negri and M. Z. Zgierski, *J. Chem. Phys.*, 1994, **100**, 2571–2587.
- 16 A. B. Trofimov, H. Köppel and J. Schirmer, *J. Chem. Phys.*, 1998, **109**, 1025–1040.
- 17 M. H. Palmer, I. C. Walker and M. F. Guest, *Chem. Phys.*, 1999, **241**, 275–296.
- 18 F. Qi, O. Sorkhabi, A. H. Rizvi, A. G. Suits, C. Sciences, V. Di, E. Orlando and L. Berkeley, *J. Phys. Chem. A*, 1999, **103**, 8351–8358.
- 19 J. Wan, M. Hada, M. Ehara and H. Nakatsuji, *J. Chem. Phys.*, 2001, **114**, 842–850.
- 20 E. E. Rennie, D. M. Holland, D. A. Shaw, C. A. Johnson and J. E. Parker, *Chem. Phys.*, 2004, **306**, 295–308.
- 21 H. Köppel, E. V. Gromov and A. B. Trofimov, *Chem. Phys.*, 2004, **304**, 35–49.
- 22 R. Weinkauff, L. Lehr, E. W. Schlag, S. Salzmann and C. M. Marian, *Phys. Chem. Chem. Phys.*, 2008, **10**, 393–404.
- 23 S. Salzmann, M. Kleinschmidt, J. Tatchen, R. Weinkauff and C. M. Marian, *Phys. Chem. Chem. Phys.*, 2008, **10**, 380–392.
- 24 G. Cui and W. Fang, *J. Phys. Chem. A*, 2011, **115**, 11544–11550.
- 25 M. Stenrup, *Chem. Phys.*, 2012, **397**, 18–25.
- 26 D. M. Holland, A. B. Trofimov, E. A. Seddon, E. V. Gromov, T. Korona, N. de Oliveira, L. E. Archer, D. Joyeux and L. Nahon, *Phys. Chem. Chem. Phys.*, 2014, **16**, 21629–21644.
- 27 D. Fazzi, M. Barbatti and W. Thiel, *Phys. Chem. Chem. Phys.*, 2015, **17**, 7787–7799.
- 28 A. Prlj, B. F. Curchod and C. Corminboeuf, *Phys. Chem. Chem. Phys.*, 2015, **17**, 14719–14730.
- 29 A. Prlj, B. F. Curchod, A. Fabrizio, L. Floryan and C. Corminboeuf, *J. Phys. Chem. Lett.*, 2015, **6**, 13–21.
- 30 P. Kölle, T. Schnappinger and R. D. Vivie-Riedle, *Phys. Chem. Chem. Phys.*, 2016, **18**, 7903–7915.
- 31 T. Schnappinger, P. Kölle, M. Marazzi, A. Monari, L. González and R. D. Vivie-Riedle, *Phys. Chem. Chem. Phys.*, 2017, **19**, 25662–25670.
- 32 O. Schalk, M. A. Larsen, A. B. Skov, M. B. Liisberg, T. Geng, T. I. Sølling and R. D. Thomas, *J. Phys. Chem. A*, 2018, **122**, 8809–8818.
- 33 D. B. Jones, M. Mendes, P. Limão-Vieira, F. Ferreira da Silva, N. C. Jones, S. V. Hoffmann and M. J. Brunger, *J. Chem. Phys.*, 2019, **150**, 064303.
- 34 R. Hakansson, B. Nordén and E. W. Thulstrup, *Chem. Phys. Lett.*, 1977, **50**, 306–308.
- 35 K. L. Reid, *Annu. Rev. Phys. Chem.*, 2003, **54**, 397–424.
- 36 J. H. Eland, *Int. J. Mass Spectrom. Ion Phys.*, 1969, **2**, 471–484.
- 37 P. Derrick, L. Åsbrink, O. Edqvist, B.-O. Jonsson and E. Lindholm, *Int. J. Mass Spectrom. Ion Phys.*, 1971, **6**, 177–190.
- 38 N. Kishimoto, H. Yamakado and K. Ohno, *J. Phys. Chem.*, 1996, **100**, 8204–8211.
- 39 J. Yang, J. Li and Y. Mo, *J. Chem. Phys.*, 2006, **125**, 174313–174317.
- 40 D. Irimia, D. Dobrikov, R. Kortekaas, H. Voet, D. A. Van Den Ende, W. A. Groen and M. H. Janssen, *Rev. Sci. Instrum.*, 2009, **80**, 1–6.
- 41 D. Irimia, R. Kortekaas and M. H. Janssen, *Phys. Chem. Chem. Phys.*, 2009, **11**, 3958–3966.
- 42 G. A. Worth, R. E. Carley and H. H. Fielding, *Chem. Phys.*, 2007, **338**, 220–227.
- 43 D. S. Parker, R. S. Minns, T. J. Penfold, G. A. Worth and H. H. Fielding, *Chem. Phys. Lett.*, 2009, **469**, 43–47.
- 44 A. D. Nunn, R. S. Minns, R. Spesyvtsev, M. J. Bearpark, M. A. Robb and H. H. Fielding, *Phys. Chem. Chem. Phys.*, 2010, **12**, 15751–15759.
- 45 R. S. Minns, D. S. Parker, T. J. Penfold, G. A. Worth and H. H. Fielding, *Phys. Chem. Chem. Phys.*, 2010, **12**, 15607–15615.
- 46 R. Spesyvtsev, O. M. Kirkby, M. Vacher and H. H. Fielding, *Phys. Chem. Chem. Phys.*, 2012, **14**, 9942–9947.
- 47 R. Spesyvtsev, O. M. Kirkby and H. H. Fielding, *Faraday Discuss.*, 2012, **157**, 165–179.
- 48 O. M. Kirkby, M. Sala, G. Balardi, R. De Nalda, L. Bañares, S. Guérin and H. H. Fielding, *Phys. Chem. Chem. Phys.*, 2015, **17**, 16270–16276.
- 49 S. P. Neville, O. M. Kirkby, N. Kaltsoyannis, G. A. Worth and H. H. Fielding, *Nat. Commun.*, 2016, **7**, 11357.
- 50 O. M. Kirkby, M. A. Parkes, S. P. Neville, G. A. Worth and H. H. Fielding, *Chem. Phys. Lett.*, 2017, **683**, 179–185.
- 51 J. W. Riley, B. Wang, J. L. Woodhouse, M. Assmann, G. A. Worth and H. H. Fielding, *J. Phys. Chem. Lett.*, 2018, **9**, 678–682.
- 52 G. A. Garcia, L. Nahon and I. Powis, *Rev. Sci. Instrum.*, 2004, **75**, 4989–4996.
- 53 S. J. Bajic, R. N. Compton, X. Tang and P. Lambropoulos, *Phys. Rev. A*, 1991, **44**, 2102–2112.
- 54 P. Celani and H.-J. Werner, *J. Chem. Phys.*, 2000, **112**, 5546–5557.
- 55 M. Richter, P. Marquetand, J. González-Vázquez, I. Sola and L. González, *J. Chem. Theory Comput.*, 2011, **7**, 1253–1258.
- 56 H.-J. Werner, P. J. Knowles, G. Knizia, F. R. Manby and M. Schütz, *WIREs Comput. Mol. Sci.*, 2012, **2**, 242–253.
- 57 H.-J. Werner, P. J. Knowles, P. Celani, W. Györfy, A. Hesselmann, D. Kats, G. Knizia, A. Köhn, T. Korona, D. Kreplin, R. Lindh, Q. Ma, F. R. Manby, A. Mitrushenkov, G. Rauhut, M. Schütz, K. R. Shamasundar, T. B. Adler, R. D. Amos, S. J. Bennie, A. Bernhardsson, A. Berning, J. A. Black, P. J. Bygrave, R. Cimiraglia, D. L. Cooper, D. Coughtrie, M. J. O. Deegan, A. J. Dobbyn, K. Doll, M. Dornbach, F. Eckert, S. Erfort, E. Goll, C. Hampel, G. Hetzer, J. G. Hill, M. Hodges, T. Hrenar, G. Jansen, C. Köppl, C. Kollmar, S. J. R. Lee, Y. Liu, A. W. Lloyd, R. A. Mata, A. J. May, B. Mussard, S. J. McNicholas,



- W. Meyer, T. F. Miller III, M. E. Mura, A. Nicklass, D. P. O'Neill, P. Palmieri, D. Peng, K. A. Peterson, K. Pflüger, R. Pitzer, I. Polyak, M. Reiher, J. O. Richardson, J. B. Robinson, B. Schröder, M. Schwilk, T. Shiozaki, M. Sibaev, H. Stoll, A. J. Stone, R. Tarroni, T. Thorsteinsson, J. Toulouse, M. Wang, M. Welborn and B. Ziegler, *MOLPRO*, 2022.1, a package of ab initio programs, 2022, see <https://www.molpro.net/>.
- 58 E. Epifanovsky, A. T. B. Gilbert, X. Feng, J. Lee, Y. Mao, N. Mardirossian, P. Pokhilko, A. F. White, M. P. Coons, A. L. Dempwolff, Z. Gan, D. Hait, P. R. Horn, L. D. Jacobson, I. Kaliman, J. Kussmann, A. W. Lange, K. U. Lao, D. S. Levine, J. Liu, S. C. McKenzie, A. F. Morrison, K. D. Nanda, F. Plasser, D. R. Rehn, M. L. Vidal, Z.-Q. You, Y. Zhu, B. Alam, B. J. Albrecht, A. Aldossary, E. Alguire, J. H. Andersen, V. Athavale, D. Barton, K. Begam, A. Behn, N. Bellonzi, Y. A. Bernard, E. J. Berquist, H. G. A. Burton, A. Carreras, K. Carter-Fenk, R. Chakraborty, A. D. Chien, K. D. Closser, V. Cofer-Shabica, S. Dasgupta, M. de Wergifosse, J. Deng, M. Diedenhofen, H. Do, S. Ehlert, P.-T. Fang, S. Fatehi, Q. Feng, T. Friedhoff, J. Gayvert, Q. Ge, G. Gidofalvi, M. Goldey, J. Gomes, C. E. González-Espinoza, S. Gulania, A. O. Gunina, M. W. D. Hanson-Heine, P. H. P. Harbach, A. Hauser, M. F. Herbst, M. Hernández Vera, M. Hodecker, Z. C. Holden, S. Houck, X. Huang, K. Hui, B. C. Huynh, M. Ivanov, A. Jász, H. Ji, H. Jiang, B. Kaduk, S. Kähler, K. Khistyayev, J. Kim, G. Kis, P. Klunzinger, Z. Koczor-Benda, J. H. Koh, D. Kosenkov, L. Koulias, T. Kowalczyk, C. M. Krauter, K. Kue, A. Kunitsa, T. Kus, I. Ladjánszki, A. Landau, K. V. Lawler, D. Lefrancois, S. Lehtola, R. R. Li, Y.-P. Li, J. Liang, M. Liebenthal, H.-H. Lin, Y.-S. Lin, F. Liu, K.-Y. Liu, M. Loipersberger, A. Luenser, A. Manjanath, P. Manohar, E. Mansoor, S. F. Manzer, S.-P. Mao, A. V. Marenich, T. Markovich, S. Mason, S. A. Maurer, P. F. McLaughlin, M. F. S. J. Menger, J.-M. Mewes, S. A. Mewes, P. Morgante, J. W. Mullinax, K. J. Oosterbaan, G. Paran, A. C. Paul, S. K. Paul, F. Pavo, Z. Pei, S. Prager, E. I. Proynov, A. Rák, E. Ramos-Cordoba, B. Rana, A. E. Rask, A. Rettig, R. M. Richard, F. Rob, E. Rossomme, T. Scheele, M. Scheurer, M. Schneider, N. Sergueev, S. M. Sharada, W. Skomorowski, D. W. Small, C. J. Stein, Y.-C. Su, E. J. Sundstrom, Z. Tao, J. Thirman, G. J. Tornai, T. Tsuchimochi, N. M. Tubman, S. P. Veccham, O. Vydrov, J. Wenzel, J. Witte, A. Yamada, K. Yao, S. Yeganeh, S. R. Yost, A. Zech, I. Y. Zhang, X. Zhang, Y. Zhang, D. Zuev, A. Aspuru-Guzik, A. T. Bell, N. A. Besley, K. B. Bravaya, B. R. Brooks, D. Casanova, J.-D. Chai, S. Coriani, C. J. Cramer, G. Cserey, A. E. DePrince, R. A. DiStasio, A. Dreuw, B. D. Dunietz, T. R. Furlani, W. A. Goddard, S. Hammes-Schiffer, T. Head-Gordon, W. J. Hehre, C.-P. Hsu, T.-C. Jagau, Y. Jung, A. Klamt, J. Kong, D. S. Lambrecht, W. Liang, N. J. Mayhall, C. W. McCurdy, J. B. Neaton, C. Ochsenfeld, J. A. Parkhill, R. Peverati, V. A. Rassolov, Y. Shao, L. V. Slipchenko, T. Stauch, R. P. Steele, J. E. Subotnik, A. J. W. Thom, A. Tkatchenko, D. G. Truhlar, T. Van Voorhis, T. A. Wesolowski, K. B. Whaley, H. L. Woodcock, P. M. Zimmerman, S. Faraji, P. M. W. Gill, M. Head-Gordon, J. M. Herbert and A. I. Krylov, *J. Chem. Phys.*, 2021, **155**, 084801.
- 59 J. V. Ortiz, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2013, **3**, 123–142.
- 60 H. H. Corzo and J. V. Ortiz, *Adv. Quantum Chem.*, 2017, **74**, 267–298.
- 61 J. V. Ortiz, *J. Chem. Phys.*, 2020, **153**, 1–28.
- 62 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery Jr, J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, Ö. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski and D. J. Fox, *Gaussian09 Revision E.01*.
- 63 S. Gozem and A. I. Krylov, *WIREs Comput. Mol. Sci.*, 2022, **12**, e1546.
- 64 J. W. Rabalais, L. O. Werme, T. Bergmark, L. Karlsson and K. Siegbahn, *Int. J. Mass Spectrom. Ion Phys.*, 1972, **9**, 185–196.
- 65 M. A. Parkes and G. A. Worth, *J. Chem. Phys.*, 2024, **161**, 114305.

