



Cite this: *Phys. Chem. Chem. Phys.*,
2024, **26**, 12892

DOI: 10.1039/d4cp90065k

rsc.li/pccp

The authors would like to make changes to Table 1 and Fig. 4c in the published manuscript. The corrections pertain to the data for the Mg(TFSI)₂ systems, where the wrong concentration of TFSI ions was used for the original calculations. The revised versions of Fig. 4c and the relevant part of Table 1 are shown here.

Table 1 Extracted enthalpy, entropy and derived Gibbs free energy at 25 °C for the dissociation of Mg(TFSI)₂ in PEO, PCL and PTMC

Salt	Polymer	$\Delta H^\circ/\text{kJ mol}^{-1}$	$\Delta S^\circ/\text{J mol}^{-1}\text{ K}^{-1}$	$\Delta G^\circ/\text{kJ mol}^{-1}$
Mg(TFSI) ₂	PEO	-8.8	13.3	-12.7
	PCL	-12.6	-6.5	-10.7
	PTMC	-9.4	1.4	-9.8




Fig. 4 (c) Van't Hoff plots of the equilibrium constant K as a function of the inverse temperature for the dissociation equilibrium of Mg(TFSI)₂ in PCL, PTMC and PEO. Dashed lines represent the linear fit to the data.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

