## Dalton Transactions



**PAPER** 

View Article Online
View Journal | View Issue



**Cite this:** *Dalton Trans.*, 2024, **53**, 11410

Received 1st March 2024, Accepted 4th June 2024 DOI: 10.1039/d4dt00627e

rsc.li/dalton

# Preparation of a high-coordinated-silicon-centered spiro-cyclic compound†

Saroj Kumar Kushvaha, D<sup>a</sup> Sai Manoj N. V. T. Gorantla, D<sup>b</sup> Paula Kallenbach, Regine Herbst-Irmer, D<sup>a</sup> Dietmar Stalke D<sup>\*\*</sup> and Herbert W. Roesky D<sup>\*\*</sup>

Silicon compounds containing silicon–silicon bond with a variety of unusual oxidation states are quite important, because their high reactivity leads to the formation of a variety of silicon compounds. The isolation of such compounds with unusual oxidation states requires a resilient synthetic strategy. Herein, we report the synthesis of a silicon based spirocyclic compound containing a hyper-valent silicon atom and a silicon–silicon bond. The computational calculations employing natural bond orbital (NBO) analysis and energy decomposition analysis–natural orbitals for chemical valence (EDA–NOCV) reveal that the nature of bonding between the silicon atoms is of an electron sharing nature.

## Introduction

Generally, silylenes are highly reactive compounds that contain divalent silicon with a lone pair of electrons and a vacant p-orbital, similar to carbenes. 1-7 The diverse reactivity of silylenes may be attributed to their unique ambiphilic nature, wherein the nucleophilic property arises from the presence of a non-bonded electron pair on the silicon atom and the electrophilic property arises due to an empty p-orbital on the silicon atom. However, donor-stabilized silvlenes have also been reported wherein the electron pair present on the nitrogen atom of an amidinate moiety is donated to the empty p-orbital of the silicon atom. Such silvlenes are more stable than the usual silylenes. Since the first report of N-heterocyclic silylene (NHSi) in 1994,4 the chemistry of this class of molecules has progressed in different directions and it has found applications including activation of molecules, 8-10 stabilization of low-valent metal atoms 11a,b and as cross-linkers in elastomers. 11c Silylenes have also been used to prepare compounds with silicon-silicon bonds containing silicon atoms in unusual oxidation states.<sup>12</sup> In addition, silylenes have also provided access to heterocyclic/spirocyclic silicon compounds that are difficult to synthesize by conventional methods. As a result, organo-silicon heterocyclic compounds and silicon centred spirocyclic compounds have been

synthesised by reacting silylenes with unsaturated alkynes, alkenes, phosphorus etc. 13-17 For example, Sen et al. isolated silicon-centred spirocyclic compounds, A and B, in which two silicon atoms are bridged by two-coordinate phosphorus atoms (Fig. 1).18 Similarly, Chen et al. and Sen et al. synthesized an important silicon spirocyclic compound, C, that possesses a unique stable 1,4-disilabenzene ring. 13,15 Moreover, Imagawa et al. have reported a spirocyclic compound, D, wherein the spirocyclic silicon centre is bound to four carbon atoms.19 These spirocyclic compounds do not possess silicon-silicon bonds. However, Scheschkewitz and coworkers reported a spirocyclic silicon compound, E, in which one of the rings contains four silicon atoms connected to each other with silicon-silicon σ-bonds and a silicon-silicon  $\pi$ -bond.<sup>20</sup> This molecule represents the first example of a main group analogue of cyclobutane-1,3-diyls, wherein no other heteroatoms are present in the ring except silicon. Similarly, Matsumoto and coworkers have synthesized a unique cyclic compound, F, in which a single silicon-silicon  $\pi$ -bond is present, not accompanied by a  $\sigma$ -bond. This single  $\pi$ -bond happens to be the longest silicon-silicon bond reported to date. Moreover, Driess and coworkers were able to stabilize a silicon-silicon double bond in a geometrically compelled disilene, G, which was accessible by inserting a suitable organic linker between two amidinato-silylene molecules. 22 These spirocyclic compounds are expected to be more reactive compared to the usual silicon compounds due to ring strain and hence, reactivity studies of these compounds can give further impetus to silicon chemistry. Herein, we report preparation of a hypercoordinated silicon-centered spirocyclic compound containing a silicon-silicon bond.

<sup>&</sup>lt;sup>a</sup>Institut für Anorganische Chemie, Georg-August Universität, Göttingen, Germany. E-mail: hroesky@gwdg.de, dstalke@chemie.uni-goettingen.de

<sup>&</sup>lt;sup>b</sup>Hylleraas Centre for Quantum Molecular Sciences, Department of Chemistry,
University of Tromsø – The Arctic University of Norway, N-9037 Tromsø, Norway
†Electronic supplementary information (ESI) available. CCDC 2292820 and
2326834. For ESI and crystallographic data in CIF or other electronic format see
DOI: https://doi.org/10.1039/d4dt00627e

**Dalton Transactions** Paper

Fig. 1 Selected examples of silicon centered spirocyclic compounds: (top) spirocyclic compounds without silicon-silicon bonds and (bottom) spirocyclic compounds with silicon-silicon bonds.

## Results and discussion

#### Synthesis of compound 1

Amidinato-silylene (2 mmol) and Ph<sub>2</sub>SiCl<sub>2</sub> (2 mmol) were mixed at -77 °C in 100 ml of toluene to obtain a clear transparent solution (Scheme 1). The reaction mixture was stirred for 1 h at room temperature. Afterwards, it was cannulated to a flask containing KC<sub>8</sub> (4 mmol) at -77 °C and was stirred overnight. Afterwards, it was filtered using a Schlenk frit under a flow of nitrogen. The clear filtrate was concentrated to about 4 ml and kept undisturbed at room temperature for two

months. The block-shaped colorless crystals of compound 1 were picked from the flask with a 13% yield. Compound 1 was thoroughly characterized by single crystal X-ray diffraction (SC-XRD), and <sup>1</sup>H, <sup>13</sup>C and <sup>29</sup>Si NMR spectroscopy (Fig. S1-S3 in the ESI†). The aliphatic protons of the tert-butyl group appear at 0.95 ppm while aromatic protons appear in the range of 6.86-8.16 ppm with a well resolved splitting pattern. <sup>13</sup>C NMR also shows resonances in the expected region wherein the tert-butyl-C appears at 52.8 ppm and the methyl-C at 30.9 ppm, whereas carbon atoms from aromatic rings resonate in the range of 127-140 ppm. The <sup>29</sup>Si NMR spectrum

Scheme 1 Synthesis of compounds 1 and 2.

Fig. 2 Molecular structure of 1 with anisotropic displacement parameters at the 50% probability level. Hydrogen atoms and minor positions of the disordered molecules are omitted for clarity. Selected bond lengths [Å] and angles [°]: Si1-N2 1.8563(18), Si1-N1 1.8690(18), Si1-Si2 2.3884(8), Cl1-Si2 2.0947(8), N2-Si1-N1 69.65(8), Cl1-Si2-Si1 113.73




Fig. 3 Molecular structure of 2 with anisotropic displacement parameters at the 50% probability level. Hydrogen atoms are omitted for clarity. Selected bond lengths [Å] and angles [°]: Cl1-Si1 2.1218(6), Si1-N2 1.8339(12), Si1-C16 1.9622(14), Si1-N1 2.0022(13), Si1-Si2 2.3378(7), Si2-C23 1.8835(15), C16-C23 1.3669(19), N2-Si1-N1 68.16(5), C16-Si1-Si2 73.79(5), C23-Si2-Si1 76.96(5).

exhibits two resonances at 35.9 and 10.2 ppm originating from silylene-silicon and non-silylene silicon centers, respectively (Fig. S7-S9†).

#### Synthesis of compound 2

Compound 1 (0.5 mmol) and 1,2-diphenylethyne (0.5 mmol) were mixed in toluene at room temperature (Scheme 1). Subsequently, the reaction mixture was stirred for 24 h. It was then filtered and the solvent was removed. The compound was dissolved in hexane and the concentrated solution was kept for crystallization at room temperature for one month. Blockshaped crystals of compound 2 were isolated with 37% yield in a month. 2 was thoroughly characterized by SC-XRD, and <sup>1</sup>H, <sup>13</sup>C and <sup>29</sup>Si NMR spectroscopy (Fig. S4-S6 in the ESI†). The aliphatic protons of the tert-butyl group appear at 1.0 ppm while the aromatic protons resonate in the range of 8.04-6.78 ppm with well resolved splitting patterns. The <sup>13</sup>C NMR spectrum also shows resonances in the expected region wherein tert-butyl-C appears at 54.8 ppm and methyl-C at 32.7 ppm. The carbon atoms from aromatic rings are observed in the range of 125 to 168 ppm. The 29Si NMR spectrum exhibits two resonances at 17.6 and -71.2 ppm from the two nonequivalent silicon atoms (Fig. S10 and S11†).

## Structural description

1 crystallizes in the orthorhombic space group Pna2<sub>1</sub> with two molecules in the asymmetric unit (Fig. 2). One silicon atom coordinates to the bidentate amidinato ligand. The second silicon atom Si2 is substituted by two phenyl groups and by a chlorine atom in a tetrahedral environment. The Si-Si-Cl bond angle is widened to 113.74(3)° while the Si-Si bond distance is 2.3884(8) Å and the Si-Cl bond distance is 2.0947(8) Å.

2 crystallizes in the triclinic space group P1 with one molecule in the asymmetric unit. Beside the four-membered ring formed by the silicon atom and the bidentate amidinato ligand, another four-membered ring is formed from an




Fig. 4 Optimized geometries of compounds 1 and 2 at the BP86-D3BJ/Def2TZVPP level. Bond lengths are in angstroms.

**Dalton Transactions** 

Table 1 NBO results of the Si<sub>L</sub>-Si bonds (L = NHC) of LSi-Si(Cl)Ph<sub>2</sub> (1) and compound 2 at the BP86-D3(BJ)/def2-TZVPP level of theory.

Occupation number ON, polarization and hybridization of the Si-Si bonds and partial charges q

Compound	Bond	ON	Polarization and hybridization (%)		WBI	$q\mathrm{Si}(\mathrm{L})$	$q\mathrm{Si/C}$
1	$Si_L$ – $Si$	1.83	Si <sub>L</sub> : 42.1 s(10.5), p(88.5), d(1.0)	Si: 57.9 s(27.3), p(72.3), d(0.4)	0.86	0.635	1.153
2	$Si_L$ – $Si$	1.87	Si <sub>L</sub> : 49.1 s(33.6), p(65.7), d(0.7)	Si: 50.9 s(22.9), p(76.5), d(0.6)	0.81	1.402	1.235
	$\mathrm{Si_L}$ –C	1.90	Si <sub>L</sub> : 25.3 s(23.9), p(75.4), d(0.7)	C: 74.7 s(29.6), p(70.2), d(0.2)	0.65		-0.393




Fig. 5 Molecular orbital pictures of 1 (top) and 2 (bottom) at the BP86-D3(BJ)/def2-TZVPP level of theory.

**Table 2** EDA–NOCV results of the Si<sub>L</sub>–Si bond in LSi–Si(Cl)Ph<sub>2</sub> (1) using two different sets of fragments with different charges and electronic states (S = singlet, D = doublet) and associated bond types at the BP86-D3(BJ)/TZ2P level. Energies are in kcal mol<sup>-1</sup>. The most favourable fragmentation scheme and bond type are given by the smallest  $\Delta E_{orb}$  value written in bold. D = dative and E = electron sharing bonds

Molecule	Bond type	Fragments	$\Delta E_{ m int}$	$\Delta E_{ m Pauli}$	$\Delta E_{ m elstat}$	$\Delta E_{ m disp}$	$\Delta E_{ m orb}$
LSi-Si(Cl)Ph <sub>2</sub> (1)	E D D	$\begin{array}{l} LSi~(D) + Si(Cl)Ph_{2}~(D) \\ (LSi)^{+}~(S) + (Si(Cl)Ph_{2})^{-}~(S) \\ (LSi)^{-}~(S) + (Si(Cl)Ph_{2})^{+}~(S) \end{array}$	-75.5 -163.4 -237.6	154.3 188.1 187.7	-106.4 -188.2 -195.3	-20.3 -20.3 -20.3	-103.1 -142.9 -209.8

additional silicon atom and the two sp²-hybridized carbon atoms of the added 1,2-diphenylethyne substrate. Both spirocyclic four-membered rings are almost planar. Remarkably, in the course of the reaction the chlorine atom at the Si( $\pi$ ) atom in the starting material is shifted to the former Si( $\pi$ ), now the penta-coordinated spiro-centre of the molecule. The bond distance between the two sp²-hybridized carbon atoms (1.3669 (19) Å) confirms the double bond character (Fig. 3).

**Table 3** The EDA–NOCV results at the BP86-D3(BJ)/TZ2P level for the Si–Si bond of the LSi–Si(Cl)Ph<sub>2</sub> (1) compound using LSi and Si(Cl)Ph<sub>2</sub> in the electronic doublet (D) states as interacting fragments. Energies are in kcal  $mol^{-1}$ 

Energy	Interaction	LSi (D) + Si(Cl)Ph <sub>2</sub> (D)
$\begin{array}{c} \Delta E_{\rm int} \\ \Delta E_{\rm Pauli} \\ \Delta E_{\rm disp}{}^{a} \\ \Delta E_{\rm elstat}{}^{a} \\ \Delta E_{\rm orb}{}^{a} \\ \Delta E_{\rm orb(1)}{}^{b} \\ \Delta E_{\rm orb(2)}{}^{b} \\ \Delta E_{\rm orb(rest)}{}^{b} \end{array}$	LSi–Si(Cl)Ph <sub>2</sub> $\sigma$ e <sup>-</sup> sharing LSi $\rightarrow$ Si(Cl)Ph <sub>2</sub> $\sigma$ e <sup>-</sup> donation	-75.5 154.3 -20.3 (8.8%) -106.4 (46.3%) -103.1 (44.9%) -86.7 (84.1%) -6.0 (5.8%) -10.4 (10.1%)

<sup>a</sup> The values in the parentheses show the contribution to the total attractive interaction  $\Delta E_{\rm elstat} + \Delta E_{\rm orb} + \Delta E_{\rm dis}$ . <sup>b</sup> The values in parentheses show the contribution to the total orbital interaction  $\Delta E_{\rm orb}$ .

## Computational calculations

We have employed DFT methods to get insight into the electronic properties of compounds 1 and 2. Fig. 4 illustrates the optimized geometries of compounds 1 and 2 in the singlet state at the BP86-D3BI/Def2TZVPP level. The computed Si<sub>L</sub>-Si (Si<sub>L</sub> = Si bonded to amidinato ligands) bond length of compound 1 is slightly longer than the experimental value, whereas in compound 2 the computed value matches well with the experimental one. The Wiberg bond indices of 0.86 and 0.81 and occupation numbers of 1.83-1.87 indicate a single bond nature of the Si<sub>L</sub>-Si bonds in 1 and 2. NBO analyses suggest a relatively strong polarization of charge (Table 1) towards the coordinated Si atom (57.9%) of the Si<sub>L</sub>-Si bond in 1 and a slight polarization in 2 (50.9%). The Si<sub>L</sub>-C bond shows a substantial charge concentration on the C atom (74.7%) of C<sub>2</sub>Ph<sub>2</sub>, indicating the direction of electron flow, and the negative charge on the C atom corroborates the same (Table 1). Fig. 5 shows the molecular orbital pictures of compounds 1 and 2. The HOMOs of both compounds 1 and 2 indicate the lone pair on the Si atoms. Their HOMO-1 and HOMO-2 demonstrate the lone pair on the N atoms of amidinato ligands (Fig. 5), whereas the HOMO-1 of compound 2 suggests a C-C π-bond of C<sub>2</sub>Ph<sub>2</sub> and the HOMO-2 of compound 1 shows an  $Si_L$ -Si  $\sigma$  type orbital interaction.




Fig. 6 The shape of the deformation density  $\Delta \rho_{(1)}$  that corresponds to  $\Delta E_{\mathrm{orb}(1)}$  and the associated fragment MOs of the LSi–Si(Cl)Ph<sub>2</sub> (1) compound at the BP86-D3(BJ)/TZ2P level at an isosurface value of 0.001 au. The eigenvalues  $|\nu_n|$  give the size of the charge migration in e. The direction of the charge flow of the deformation densities is red  $\rightarrow$  blue.

We have evaluated the nature of the  $\mathrm{Si_L-Si}$  bond of LSi–Si (Cl)Ph<sub>2</sub> (1) using energy decomposition analysis with natural orbitals for chemical valence analysis (EDA–NOCV). To understand the best bonding scenario, we have explored three different bonding possibilities with varied charges and multiplicities, *i.e.*, a neutral LSi and a neutral  $\mathrm{Si}(\mathrm{Cl})\mathrm{Ph_2}$  fragment in the electronic doublet state forming an electron sharing bond and a singly charged LSi (-/+) and a  $\mathrm{Si}(\mathrm{Cl})\mathrm{Ph_2}$  (+/-) fragment in the electronic singlet state forming a dative bond (Table 2). The bonding possibility with the smallest change in orbital interactions ( $\Delta E_{\mathrm{orb}}$ ) between the fragments is rated as the best bonding option. <sup>22–24</sup> Accordingly, from Table 2, it is evident that the electron sharing interaction between neutral LSi and  $\mathrm{Si}(\mathrm{Cl})\mathrm{Ph_2}$  fragments in a doublet state with the least  $\Delta E_{\mathrm{orb}}$  is the best bonding model.

The  $Si_L$ –Si bond of compound 1 is characterized by strong contributions from electrostatic and orbital interactions but with slightly dominant electrostatic characteristics (Table 3). The dispersion interactions also contribute significantly (8.8%) to the stability of the  $Si_L$ –Si bond. The pair-wise breakdown of total orbital interactions (Table 3) suggests a major contribution (84.1%) from the  $\sigma$  electron-sharing interaction, which is depicted in the deformation density picture (Fig. 6). Additionally, a weak contribution from  $LSi \rightarrow Si(Cl)Ph_2 \ \sigma \ e^-$  donation (5.8%) can also be observed (Table 3). The oxidation states of the silicon atoms were calculated by the LOBA (localized orbital bonding analysis) method<sup>25</sup> using Multiwfn.  $^{26a}$  The silicon atoms in 1 display the +2 and +4 oxidation states, whereas in compound 2 both silicon atoms possess the +4 oxidation state.

### Conclusion

In conclusion, we have prepared a spirocyclic compound containing a hyper-coordinated silicon atom. This spirocycle contains a silicon–silicon bond which is rarely found in silicon-based chemistry. The theoretical studies using NBO and EDA–NOCV calculations revealed the nature of the silicon–silicon bond to be electron sharing.

## **Experimental section**

#### General procedures and instrumentation

The required chemicals for the preparation of starting materials were purchased from commercial sources and were used without further purification. The required solvents were also purchased from commercial sources and were dried by stirring over Na/K alloy for four days. The dried and distilled solvents were collected immediately before use in the reactions. All manipulations were performed under inert conditions using standard Schlenk and glovebox techniques under a flow of high purity dinitrogen/argon gas. Deuterated NMR solvents ( $C_6D_6/THF-D_8/toluene-D_8$ ) were dried by stirring them for several days over Na/K alloy followed by distil-

lation under vacuum and degassing several times.  $^1H$ ,  $^{13}C$  and  $^{29}Si$  NMR spectra were recorded on Bruker Avance 400/500 MHz, 125 MHz and 99 MHz NMR spectrometers, respectively, and referenced to the resonances of the solvent used. Amidinato-silylene chloride, LSiCl (L = PhC(N^tBu)<sub>2</sub>), was synthesized as per the procedure reported in the literature.  $^1$ 

Characterization of 1. <sup>1</sup>H NMR (400 MHz, 298K, C<sub>6</sub>D<sub>6</sub>, ppm):  $\delta$  = 8.15 (d, J = 7.3, 4H, Ar–CH), 7.54–7.48 (m, 1H, Ar–CH), 7.22 (t, J = 7.4, 4H, Ar–CH), 7.18–7.12 (m, 2H, Ar–CH), 7.06 (d, J = 7.8, 1H, AR–CH), 7.02–6.94 (m, 2H, Ar–CH), 6.94–6.86 (m, 1H, Ar–CH), 0.95 (s, 18H, CH<sub>3</sub>). <sup>13</sup>C NMR (100 MHz, 298K, C<sub>6</sub>D<sub>6</sub>):  $\delta$  = 156.4 (NCN), 139.3 (Ar–C), 134.3 (Ar–C), 133.9 (Ar–C), 131.4 (Ar–C), 129.4 (Ar–C), 128.9 (Ar–C), 128.6 (Ar–C), 127.8 (Ar–C), 127.2 (Ar–C), 52.8 (<sup>†</sup>Butyl–C), 30.9 (CH<sub>3</sub>). <sup>29</sup>Si NMR (99 MHz, C<sub>6</sub>D<sub>6</sub>, ppm):  $\delta$  = 35.9 and 10.2. Anal. (%) calcd for compound 1 (C<sub>27</sub>H<sub>33</sub>ClN<sub>2</sub>Si<sub>2</sub>): C, 67.96; H, 6.97; N, 5.87, Found: C, 67.82; H, 7.22; N, 5.68.

Characterization of 2.  $^{1}$ H NMR (500 MHz, 298K, toluene-D<sub>8</sub>, ppm):  $\delta$  = 8.04–8.02 (m, 4H, Ar–CH), 7.83–7.81 (m, 2H, Ar–CH), 7.26–7.24 (m, 6H, Ar–CH), 7.17–7.10 (m, 2H, Ar–CH), 7.01–6.98 (m, 2H, Ar–CH), 6.92–6.68 (m, 7H, Ar–CH), 6.86–6.78 (m, 2H, Ar–CH), 1.00 (s, 18H, CH<sub>3</sub>).  $^{13}$ C NMR (125 MHz, 298K, toluene-D<sub>8</sub>, ppm):  $\delta$  = 167.7 (NCN), 161.0 (Ar–C), 144.3 (Ar–C), 142.5 (Ar–C), 136.7 (Ar–C), 135.1 (Ar–C), 130.4 (Ar–C), 129.9 (Ar–CH), 129.5 (Ar–C), 128.3 (Ar–C), 126.6–126.5 (C=C), 54.8 (<sup>†</sup>Butyl-C), 32.7 (CH<sub>3</sub>).  $^{29}$ Si NMR (99 MHz, 298K, toluene-D<sub>8</sub>, ppm):  $\delta$  = 17.6 and –71.2.

Crystal data. The datasets were collected on a Bruker D8 three circle diffractometer, equipped with a SMART APEX II CCD detector and an INCOATEC microfocus source (Mo  $K_{\alpha}$  radiation) with INCOATEC Quazar mirror optics. The data were integrated with SAINT<sup>27</sup> and a multi-scan absorption correction was applied using SADABS. The structures were solved by SHELXT<sup>29</sup> and refined on  $F^2$  using SHELXL<sup>30</sup> in the graphical user interface ShelXle. The structures were solved by SHELXT<sup>29</sup> and refined on  $F^2$  using SHELXL<sup>30</sup> in the graphical user interface ShelXle.

Crystal data for 1 at 100(2) K: C<sub>27</sub>H<sub>33</sub>ClN<sub>2</sub>Si<sub>2</sub>,  $M_{\rm r}$  = 477.18 g mol<sup>-1</sup>, 0.34 × 0.175 × 0.13 mm, orthorhombic,  $Pna2_1$ , a = 25.543(4) Å, b = 9.225(2) Å, c = 22.485(3) Å, V = 5298.2(16) Å<sup>3</sup>, Z = 8,  $\mu$ (MoK $_{\alpha}$ ) = 0.252 mm<sup>-1</sup>,  $\theta_{\rm max}$  = 28.393°, 200 799 reflections measured, 13 285 independent ( $R_{\rm int}$  = 0.0488),  $R_{\rm 1}$  = 0.0268 [I > 2 $\sigma$ (I)], w $R_{\rm 2}$  = 0.0668 (all data),  $\Delta \rho_{\rm max}/\Delta \rho_{\rm min}$  = 0.218/–0.169 e Å<sup>-3</sup>, CCDC 2292820.†

Crystal data for 2 at 100(2) K:  $C_{41}H_{43}ClN_2Si_2$ ,  $M_r = 655.40$  g mol<sup>-1</sup>,  $0.183 \times 0.152 \times 0.124$  mm, triclinic,  $P\bar{1}$ , a = 10.498(2) Å, b = 10.599(2) Å, c = 17.493(4) Å,  $\alpha = 72.73(2)^\circ$ ,  $\beta = 85.07(3)^\circ$ ,  $\gamma = 75.62(2)^\circ$ , V = 1800.3(7) Å<sup>3</sup>, Z = 2,  $\mu(MoK_\alpha) = 0.204$  mm<sup>-1</sup>,  $\theta_{max} = 27.219^\circ$ , 80 583 reflections measured, 8015 independent ( $R_{int} = 0.0409$ ),  $R_1 = 0.0325$  [ $I > 2\sigma(I)$ ], w $R_2 = 0.0849$  (all data),  $\Delta \rho_{max}/\Delta \rho_{min} = 0.343/-0.271$  e Å<sup>-3</sup>, CCDC 2326834.†

### Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

S. M. N. V. T. G. thanks Prof. Kathrin Hopman, Department of Chemistry, UiT for computational facilities. D. St. thanks the BENCh, which is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - 389479699/GRK2455. Dedicated to Professor C. N. R. Rao on the occasion of his 90<sup>th</sup> birthday.

## References

Paper

- 1 S. S. Sen, H. W. Roesky, D. Stern, J. Henn and D. Stalke, J. Am. Chem. Soc., 2010, 132, 1123-1126.
- 2 S. Yao, Y. Xiong, A. Saddington and M. Driess, Chem. Commun., 2021, 57, 10139-10153.
- 3 J. Li, J. Li, Y. Liu, S. Kundu, H. Keil, H. Zhu, R. Herbst-Irmer, D. Stalke and H. W. Roesky, Inorg. Chem., 2020, 59, 7910-7914.
- 4 M. Denk, R. Lennon, R. Hayashi, R. West, A. V. Belyakov, H. P. Verne, A. Haaland, M. Wagner and N. Metzler, J. Am. Chem. Soc., 1994, 116, 2691-2692.
- 5 M. Kira, Chem. Commun., 2010, 46, 2893-2903.
- 6 M. Haaf, T. A. Schmedake and R. West, Acc. Chem. Res., 2000, 33, 704-714.
- 7 S. S. Sen, S. Khan, P. P. Samuel and H. W. Roesky, Chem. Sci., 2012, 3, 659-682.
- 8 C. Shan, S. Yao and M. Driess, Chem. Soc. Rev., 2020, 49, 6733-6754.
- 9 S. Fujimori and S. Inoue, Eur. J. Inorg. Chem., 2020, 2020, 3131-3142.
- 10 J. Keuter, A. Hepp, A. Massolle, J. Neugebauer, C. Mück-Lichtenfeld and F. Lips, Angew. Chem., Int. Ed., 2022, 61, e202114485.
- 11 (a) Y. Wang, M. Karni, S. Yao, A. Kaushansky, Y. Apeloig and M. Driess, J. Am. Chem. Soc., 2019, 141, 12916-12927; (b) S. Du, H. Jia, H. Rong, H. Song, C. Cui and Z. Mo, Angew. Chem., Int. Ed., 2022, 61, e202115570; (c) F. A. D. Herz, M. Nobis, D. Wendel, P. Pahl, P. J. Altmann, J. Tillmann, R. Weidner, S. Inoue and B. Rieger, Green Chem., 2020, 22, 4489-4497.
- 12 (a) A. V. Protchenko, A. D. Schwarz, M. P. Blake, C. Jones, N. Kaltsoyannis, P. Mountford and S. Aldridge, Angew. Chem., Int. Ed., 2013, 52, 568-571; (b) M. M. D. Roy, M. J. Ferguson, R. McDonald, Y. Zhou and E. Rivard, Chem. Sci., 2019, 10, 6476-6648; (c) D. Reiter, R. Holzner, A. Porzel, P. J. Altmann, P. Frisch and S. Inoue, J. Am. Chem. Soc., 2019, 141, 13536-13546; (d) M. K. Bisai, V. S. V. S. N. Swamy, T. Das, K. Vanka, R. G. Gonnade and S. S. Sen, *Inorg. Chem.*, 2019, 58(16), 10536–10542; (e) S. K. Kushvaha, P. Kallenbach, N. V. T. S. M. Gorantla, R. Herbst-Irmer, D. Stalke and H. W. Roesky, Chem. - Eur. J., 2023, e202303113.
- 13 (a) Y. Chen, J. Li, Y. Zhao, L. Zhang, G. Tan, H. Zhu and H. W. Roesky, J. Am. Chem. Soc., 2021, 143, 2212-2216;

- (b) S. S. Sen, S. Khan, S. Nagendran and H. W. Roesky, Acc. Chem. Res., 2012, 45, 578-587.
- 14 W. H. Atwell, Organometallics, 2009, 28, 3573-3586.
- 15 S. S. Sen, H. W. Roesky, K. Meindl, D. Stern, J. Henn, A. C. Stückl and D. Stalke, Chem. Commun., 2010, 46, 5873-5875.
- 16 J. Li, M. Zhong, H. Keil, H. Zhu, R. Herbst-Irmer, D. Stalke, S. De, D. Koley and H. W. Roesky, Chem. Commun., 2019, 55, 2360,
- 17 R. Azhakar, S. Pillai Sarish, G. Tavčar, H. W. Roesky, J. Hey, D. Stalke and D. Koley, *Inorg. Chem.*, 2011, **50**(7), 3028–3036.
- 18 S. S. Sen, S. Khan, H. W. Roesky, K. Meindl, J. Henn, D. Stalke, J.-P. Demers and A. Lange, Angew. Chem., Int. Ed., 2011, 50, 2322-2325.
- 19 T. Imagawa, L. Giarrana, D. M. Andrada, B. Morgenstern, M. Nakamoto and D. Scheschkewtz, J. Am. Chem. Soc., 2023, 145, 4757-4764.
- 20 C. B. Yildiz, K. I. Leszczynska, S. Gonzlez-Gallardo, M. Zimmer, A. Azizoglu, T. Biskup, C. W. M. Kay, V. Huch, H. S. Rzepa and D. Scheschkewitz, Angew. Chem., Int. Ed., 2020, 59, 15087-15092.
- 21 S. Kyushin, Y. Kurosaki, K. Otsuka, H. Imai, S. Ishida, T. Kyomen, M. Hanaya and H. Matsumoto, Nat. Commun., 2020, 11, 4009.
- 22 A. Kostenko and M. Driess, J. Am. Chem. Soc., 2018, 140, 16962-16966.
- 23 (a) Q. Zhang, W.-L. Li, C. Xu, M. Chen, M. Zhou, J. Li, D. M. Andrada and G. Frenking, Angew. Chem., Int. Ed., 2015, **54**, 11078; (b) D. M. Andrada and G. Frenking, *Angew*. Chem., Int. Ed., 2015, 54, 12319; (c) R. Saha, S. Pan, G. Merino and P. K. Chattaraj, Angew. Chem., Int. Ed., 2019, 58, 8372; (d) Q. Wang, S. Pan, Y. Wu, G. Deng, G. Wang, L. Zhao, M. Zhou and G. Frenking, Angew. Chem., Int. Ed., 2019, 58, 17365; (e) S. M. N. V. T. Gorantla, P. Parameswaran and K. C. Mondal, J. Comput. Chem., 2021, 42, 1159; (f) S. K. Kushvaha, S. M. N. V. T. Gorantla and K. C. Mondal, J. Phys. Chem. A, 2022, 126, 845-858; (g) G. Frenking, S. M. Gorantla, S. Pan and K. C. Mondal, Chem. - Eur. J., 2020, 26, 14211.
- 24 F. Engelhardt, C. Maaß, D. M. Andrada, R. Herbst-Irmer and D. Stalke, Chem. Sci., 2018, 9, 3111-3121.
- 25 A. J. W. Thom, E. J. Sundstrom and M. Head-Gordon, *Phys.* Chem. Chem. Phys., 2009, 11, 11297-11304.
- 26 (a) T. Lu and F. Chen, J. Comput. Chem., 2012, 33, 580-592; (b) S. K. Kushvaha, P. Kallenbach, N. V. T. S. M. Gorantla, R. Herbst-Irmer, D. Stalke, P. Parameswaran H. W. Roesky, Chem. - Eur. J., 2024, 30, e202303113.
- 27 Bruker AXS Inc., in Bruker Apex CCD, SAINT v8.40B, Bruker AXS Inst. Inc., WI, USA, Madison, 2019.
- 28 L. Krause, R. Herbst-Irmer, G. M. Sheldrick and D. Stalke, J. Appl. Crystallogr., 2015, 48, 3-10.
- 29 G. M. Sheldrick, Acta Crystallogr., Sect. A: Found. Adv., 2015, **71**, 3–8.
- 30 G. M. Sheldrick, Acta Crystallogr., Sect. C: Struct. Chem., 2015, 71, 3-8.
- 31 C. B. Hübschle, G. M. Sheldrick and B. Dittrich, J. Appl. Crystallogr., 2011, 44, 1281-1284.