

Faraday Discussions

Accepted Manuscript



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

This article can be cited before page numbers have been issued, to do this please use: R. Lipowsky, *Faraday Discuss.*, 2024, DOI: 10.1039/D4FD00184B.

Cite this: DOI: 00.0000/xxxxxxxxxx

The many faces of membrane tension for biomembranes and vesicles[†]

10 Dec 2024

Reinhard Lipowsky^{*a}Received Date
Accepted Date

DOI: 00.0000/xxxxxxxxxx

Membrane tension is generated by external forces and constraints that affect the membrane's surface area. Many studies have been pursued to elucidate this intuitive concept. Here, we focus on biomimetic model systems, for which the notion of membrane tension can be elaborated in a quantitative manner. Bottom-up, these membrane systems are planar bilayers, unilamellar nanovesicles, and giant unilamellar vesicles (GUVs). For planar bilayers and nanovesicles, we need to distinguish the mechanical bilayer tension within the whole bilayer from the mechanical leaflet tensions within the individual bilayer leaflets. In addition, the fluctuation spectrum of the bilayer defines the bilayer's fluctuation tension. These different tensions can be determined by molecular dynamics simulations. On the micron-scale, GUV membranes as observed by light microscopy experience both a mechanical and a curvature-elastic tension, as follows from the theory of curvature elasticity. The curvature-elastic tension represents a material parameter whereas the mechanical tension is a kind of hidden variable that depends on the size and shape of the membrane as demonstrated for multispherical shapes of GUVs. Furthermore, the excess membrane area stored in the shape fluctuations is again governed by the fluctuation tension. Many membrane systems exhibit characteristic tension thresholds. Examples include the fusion of nanovesicles, the shapes of tubular membranes, and the morphological responses of vesicle membranes to condensate droplets.

1 Introduction

Experimental procedures that generate membrane tension include osmotic inflation, adhesion, and micropipette aspiration of vesicles. In order to elucidate the concept of membrane tension, it is very useful to consider biomimetic model membranes. Here, we will study three such systems: planar lipid bilayers, unilamellar nanovesicles, and giant unilamellar vesicles (GUVs).

On the nanometer scale, the mechanical tension of the membrane is provided by the bilayer tension as determined by molecular dynamics simulations. This bilayer tension can be decomposed into two leaflet tensions, which control the spatio-temporal remodeling of lipid bilayers and nanovesicles.¹ On the micrometer scale, the total membrane tension of a biomembrane with transbilayer asymmetry is equal to the sum of its mechanical and its spontaneous or curvature-elastic tension.^{2,3}

This paper is organized as follows. The next Sect 2 emphasizes three fundamental aspects of biomembranes as provided by their molecular bilayer structure, their fluidity, and their shape fluctuations. In Sect 2.3, we remind the reader about the important differences between membrane tension and interfacial tension. The

subsequent Sects 3, 4, and 5 elucidate the different notions of membrane tension for planar bilayers, unilamellar nanovesicles, and giant unilamellar vesicles (GUVs), respectively. In Sect 6, we show that the remodeling of GUVs by condensate droplets is governed by the competition between the curvature-elastic tensions of the membrane segments and the interfacial tension of the liquid-liquid interface. The last Sect 7 describes the most important conclusions.

2 Fundamental aspects of biomembranes

2.1 Symmetric and asymmetric bilayer membranes

All biomembranes are based on lipid bilayers, consisting of two leaflets (or monolayers) of lipid molecules. In general, the two leaflets may differ in their lipid composition, in the packing densities of these lipids – as described by area per lipid or volume per lipid – and in the molecular composition of the two aqueous solutions adjacent to the two leaflets. All of these differences contribute to the transbilayer asymmetry and the associated spontaneous curvature.

For the closed membrane of a vesicle, the inner leaflet is in contact with the interior aqueous solution, the outer leaflet with the exterior aqueous solution. Furthermore, biomembranes typically contain membrane proteins which often exhibit an average orientation towards the interior or towards the exterior so-

^a Max Planck Institute of Colloids and Interfaces, 14424 Potsdam, Germany. Tel: +49 331 5679600; E-mail: lipowsky@mpikg.mpg.de

[†] Electronic Supplementary Information (ESI) available: Sects S1 – S9, Figs S1 – S15, and Table 1. See DOI: 00.0000/00000000.



lution. Therefore, a symmetric bilayer membrane is obtained if both leaflets have the same lipid composition and exhibit the same packing densities for all lipid components. Furthermore, a symmetric bilayer also requires that the exterior solution has the same molecular composition as the interior solution. Such a symmetric membrane represents a highly idealized bilayer state but also provides a reference states that is useful from a conceptual point of view. Furthermore, such a symmetric bilayer is easy to prepare in molecular dynamics simulations, provided one uses a planar bilayer with periodic boundary conditions. Experimentally, a planar membrane geometry is obtained for hole-spanning or black lipid membranes⁴ and for pore-spanning membranes⁵.




Fig. 1 Two examples for one-component lipid bilayers with transbilayer asymmetry: (a,b) Planar bilayer consisting of 920 lipids in its upper and 738 lipids in its lower leaflet. The lipids have blue headgroups and red chains; and (c) Planar bilayer with 832 lipids in the upper and 832 lipids in the lower leaflet but exposed to two different concentrations of solutes (grey beads) in the upper and lower solution. The solutes adsorb onto the bilayer and form different adsorption layers at the two leaflets. The bilayers have a thickness of about 4 nm.⁶

2.2 Fluidity of membranes

The most fundamental property of a biomembrane is its fluidity. On the molecular scale, the lipid molecules undergo fast lateral diffusion with a diffusion constant of the order of a few μm^2 per second. Thus, if we consider two lipids that are in contact to each other, it takes of the order of one or a few nanoseconds for them to swap their spatial positions. On the nanoscale, fluid membranes can exhibit two types of elastic deformations. First, they can be stretched and compressed, which leads to different areas per lipid or different volumes per lipid. Second, they can attain bent or curved states, which deviate from the essentially planar states in Fig 1. These two elastic deformations follow from classical elasticity theory⁷ for a liquid material with vanishing shear modulus.

Stretching and compression of membranes. First, let us consider the stretching and compression of a membrane in response to a mechanical tension, Σ , acting on this membrane. The fluidity of the membrane implies that this mechanical tension must be spatially uniform and isotropic for any (meta)stable state of the membrane. Indeed, tension gradients generate lipid flows within the membrane but these flows decay when the membrane relaxes

towards a stable state.

If the membrane is tensionless, it attains a certain optimal area $A = A_0$. In general, the membrane area A differs from this optimal value and is then subject to the mechanical membrane tension

$$\Sigma = K_A(A - A_0)/A_0 \equiv K_A\Delta a, \quad (1)$$

which is proportional to the area compressibility modulus K_A and to the relative area dilation $(A - A_0)/A_0 = \Delta a$. A positive mechanical tension, $\Sigma > 0$, leads to membrane stretching, a negative tension, $\Sigma < 0$, to membrane compression. The Hooke-type relationship between Σ and $(A - A_0)/A_0$ has been used, e.g., to analyse micropipette aspiration experiments.⁸

The corresponding elastic energy is obtained from Eq (1) by integration over the membrane area A , which leads to the elastic stretching energy

$$E_{st} = \int_{A_0}^A dA' \Sigma(A') = \frac{1}{2} K_A \frac{(A - A_0)^2}{A_0}. \quad (2)$$

This energy represents the reversible work that we have to perform in order to stretch or compress the membrane from its optimal value A_0 at zero tension to its actual value A . Note that the elastic energy E_{st} is always positive, both for stretching with $A > A_0$ and for compression with $A < A_0$.

Bending of membranes. So far, we have not considered bent or curved states of the membrane and the associated bending energies. On the nanoscale, the curvature of the membrane is described by its principal curvatures, C_1 and C_2 , or equivalently by the mean and Gaussian curvatures $M \equiv \frac{1}{2}(C_1 + C_2)$ and $G \equiv C_1 C_2$. As explained in Sect S1[†], the shape of a vesicle with volume V and membrane area A is computed by minimizing the elastic shape functional

$$F_{el} = -\Delta P V + \Sigma' A + 2\kappa \int dA (M - m)^2 \quad (3)$$

which depends on the two Lagrange multipliers ΔP and Σ' as well as on the area integral over the bending energy density $2\kappa(M - m)^2$. This energy density involves two curvature-elastic parameters, the bending rigidity κ and the spontaneous curvature m .

The Lagrange multiplier Σ' represents a tension. It can be shown⁹ that the Lagrange multiplier tension Σ' is equal to the mechanical tension Σ in Eq (1), that is,

$$\Sigma' = \Sigma = K_A\Delta a. \quad (4)$$

This equality applies to all (meta)stable states of the vesicle membrane as obtained by the constrained minimization of the elastic shape functional F_{el} in Eq (3). This minimization of F_{el} leads to the local shape equation, which has the general form²

$$\Delta P \approx \Sigma_{tot} M \quad \text{with } \Sigma_{tot} \equiv \Sigma + 2\kappa m^2 \quad (5)$$

to first order in the mean curvature M and defines the total membrane tension Σ_{tot} and the spontaneous or curvature-elastic tension $2\kappa m^2$. The curvature-elastic tension can be directly measured by micropipette aspiration of tubulated GUVs.¹⁰






Fig. 2 Simulation snapshot of a symmetric and tensionless bilayer. The lipid bilayer has a thickness of about 4 nm. The smallest curvature radius of its midsurface (red) is about 6 nm, the radius of the dashed circles.¹⁴

2.3 Shape fluctuations of fluid membranes

In the low tension regime, GUVs exhibit shape fluctuations in the form of bending undulations that are clearly visible in the optical microscope. Likewise, these undulations can be observed in molecular dynamics simulation, see Fig 2. The influence of membrane tension on the membrane's shape fluctuations led to a long and controversial debate. This debate started in the early 1980s with several studies^{11–13}, which considered both liquid-liquid interfaces with low interfacial tension^{11,13} and fluid membranes with low membrane tension^{12,13}. These two tensions must be distinguished, however, because they have fundamentally different properties.

Interfacial tension versus mechanical membrane tension.

First, the interfacial tension is always positive as follows from the thermodynamic stability of the corresponding liquid-liquid interface¹⁵, whereas the mechanical membrane tension Σ in Eq (1) can be positive or negative. Second, the mechanical membrane tension depends on the size and shape of the membrane, as demonstrated for multispherical vesicle shapes in Sect 5.1 below. This shape dependence of Σ also follows when we apply the identity $\Sigma' = \Sigma$ in Eq (4) to the curvature-elastic theory of vesicle adhesion¹⁶. In contrast, the interfacial tension is independent of the size and shape of the liquid-liquid interface. Third, the shape fluctuations of membranes and interfaces have very different scaling properties as described in Sect S2[†]. In particular, in contrast to interfaces, fluid membranes have a finite persistence length ξ_p for the correlations of their normal vectors, with different types of shape fluctuations on length scales below and above the persistence length.

Persistence length of fluid membranes. Below their persistence length, fluid membranes undergo fluctuations with relatively small amplitudes around their average shape as in Fig 2. In contrast, the membranes attain crumpled states with no average orientation of their normal vectors when their sizes are large compared to the persistence length ξ_p . Likewise, boundary conditions imposed on the membrane have no effect on those membrane segments that have a large spatial separation $\Delta L \gg \xi_p$ from this boundary.

The persistence length ξ_p of a fluid membrane grows exponentially with the ratio of the bending rigidity κ to the thermal energy

$k_B T$ according to^{12,13,17}

$$\xi_p = \ell \exp[c_p \kappa / k_B T] \quad \text{with a molecular length scale } \ell \quad (6)$$

and the dimensionless coefficient $c_p = 4\pi/3$ as determined by Monte Carlo simulations of discretized fluid membranes¹⁷.

From a conceptual point of view, the persistence length as given by Eq (6) provides an important cross-over length between two different regimes of shape fluctuations. In practise, this length scale is not relevant for lipid bilayers. Indeed, using the bending rigidity $\kappa \simeq 20 k_B T$ for lipid bilayers at room temperature and the molecular length scale $\ell \simeq 6$ nm, corresponding to the smallest curvature radius in Fig 2, we obtain the estimate that ξ_p is of the order of 10^{23} km which is astronomical compared to the size of the largest biomembranes. In fact, a fluid membrane is likely to form a pore across the bilayer when the lateral size of the membrane is still much smaller than the persistence length. Indeed, as shown in Sect 3.3 below, any small but finite mechanical tension $\Sigma > 0$ leads to poration for a sufficiently large membrane.

3 Membrane tensions of planar bilayers

We now model the membranes as molecular bilayers assembled from one or several lipid components. These bilayers can be studied by molecular dynamics simulations. Using such simulations, one can determine several types of tensions: the mechanical bilayer tension, the mechanical leaflet tensions, and the fluctuation tension. The mechanical tensions are computed via the stress profile across the bilayer. The fluctuation tension is deduced from the spectrum of bending undulations on length scales that are small compared to the persistence length.

In order to reduce boundary effects, all simulation results for planar bilayers as described here have been obtained for periodic boundary conditions. Thus, we will not consider any “frame tension” as sometimes discussed in the literature. In fact, if we fixed the edge of the membrane by a frame, we would also have to prescribe the normal vectors of the membrane along this frame.

3.1 Mechanical tensions of planar bilayers and leaflets

Each planar bilayer consists of two leaflets, the lower and the upper leaflet labeled by ll and ul , respectively. Both leaflets represent two-dimensional liquids, which are subject to the leaflet tensions Σ_{ll} and Σ_{ul} . In mechanical equilibrium, each of these leaflet tensions must be laterally uniform. As a consequence, the bilayer tension Σ can be decomposed according to

$$\Sigma = \Sigma_{ll} + \Sigma_{ul}. \quad (7)$$

The computation of these tensions from the stress profile across the bilayer is described in Sect S3[†].

3.2 Leaflet tension space for planar bilayers

Planar bilayers with a fixed total number of lipids, $N_{ll} + N_{ul}$, define a two-dimensional leaflet tension space, as depicted in Fig 3 for $N_{ll} + N_{ul} = 1682$. The origin of this leaflet tension space as given by $\Sigma_{ll} = \Sigma_{ul} = 0$ defines the relaxed reference state with two tensionless leaflets. This reference state of the planar bilayers can be






Fig. 3 Leaflet tension space for planar bilayers that contain a total number of $N_{ll} + N_{ul} = 1682$ lipids.¹⁸ The two coordinates are the leaflet tensions Σ_{ll} and Σ_{ul} in the lower and upper leaflets. Negative and positive leaflet tensions describe compressed and stretched leaflets. The reference state with tensionless leaflets, corresponding to $\Sigma_{ll} = \Sigma_{ul} = 0$, is obtained for the symmetric bilayer with $N_{ll} = N_{ul} = 841$ lipids. The red data points describe elastic deformations with equal leaflet tensions (ELTs), $\Sigma_{ll} = \Sigma_{ul}$. The black data represent bilayers with opposite leaflet tensions (OLTs), $\Sigma_{ll} = -\Sigma_{ul}$ and $\Sigma = 0$. All OLT states can be obtained from the reference state by reshuffling lipids from one leaflet to the other and adjusting the base area of the simulation box to obtain tensionless bilayers.

obtained by focusing on symmetric bilayers with $N_{ll} = N_{ul} = 841$ and identifying the unique bilayer with vanishing bilayer tension $\Sigma = 0$. To characterize the elastic response of the reference state, it is useful to distinguish two types of elastic deformations, corresponding to the red and black data in Fig 3.

The red data represent elastic deformations of the reference state with equal leaflet tensions (ELTs), that is, with

$$\Sigma_{ll} = \Sigma_{ul} \quad (\text{equal leaflet tensions}). \quad (8)$$

The ELT states are located along the main diagonal of the leaflet tension space in Fig 3. These ELT states are described in more detail in Sect S4[†] and Figs S2[†] – S4[†]. On the other hand, the black data in Fig 3 correspond to elastic deformations of the reference state with opposite leaflet tensions (OLTs), that is, with

$$\Sigma_{ll} = -\Sigma_{ul} \quad (\text{opposite leaflet tensions}), \quad (9)$$

The OLT states are located on the diagonal which is orthogonal to the main diagonal in Fig 3. All OLT states can be obtained from the reference state by reshuffling lipids from one leaflet to the other, keeping the total lipid number $N_{ll} + N_{ul}$ constant and imposing the constraint of vanishing bilayer tension $\Sigma = \Sigma_{ll} + \Sigma_{ul} = 0$. As a consequence, one leaflet becomes compressed by a negative leaflet tension whereas the other leaflet becomes stretched by a positive leaflet tension.

The different OLT states of the planar bilayer can be distin-

guished by their stress asymmetry^{6,19}

$$\Delta\Sigma \equiv \Sigma_{ul} - \Sigma_{ll}. \quad (10)$$

The stress asymmetry $\Delta\Sigma$ vanishes for all ELT states of the planar bilayer, for which both leaflets contain the same number of lipids, i.e., for which $N_{ul} = N_{ll}$. In contrast, all OLT states of the planar bilayer exhibit a nonzero stress asymmetry, which increases in magnitude as we move away from the reference state with $\Sigma_{ll} = \Sigma_{ul} = 0$. Furthermore, all bilayer states with the same nonzero value of the stress asymmetry, $\Delta\Sigma = \text{const}$, are located on a line parallel to the main diagonal. Stress asymmetry in planar bilayers has also been studied by Deserno and coworkers^{20,21}, who used the term “differential stress” instead of “stress asymmetry”. The elastic response of the planar bilayer leaflets to the leaflet tensions can be described both in terms of area per lipid and of volume per lipid (Sect S4[†], Figs S2[†], and S3[†]), where volume per lipid is obtained via Voronoi tessellation (Fig S4[†]).

3.3 Formation of stable bilayer pores

For a sufficiently large and positive bilayer tension Σ , the bilayer will rupture and form a pore. To illustrate this poration process, we now consider a planar and symmetric bilayer, which contains $N/2$ identical lipid molecules in each leaflet. We assume that the membrane area stored in the shape fluctuations of the bilayer can be ignored. As a consequence, the surface area A of the bilayer membrane is close to its projected area $A_{||}$, which can be directly controlled in molecular dynamics simulations by the geometry of the simulation box. Indeed, for a planar bilayer that spans a cubic simulation box, the projected area of the bilayer is equal to the base area of this box. Furthermore, the bilayer has an optimal projected area $A_{||} = A_0$ for which the mechanical tension $\Sigma = (A_{||} - A_0)/A_0$ as given by Eq (1) vanishes. When $A_{||}$ is larger than A_0 , the bilayer is stretched by a positive mechanical tension $\Sigma > 0$.

Now, let us compare, for a given projected area $A_{||} > A_0$, two different bilayer states: State (Un) corresponds to the unporated state with constant area per lipid, $a = 2A_{||}/N$, state (Po) to a porated bilayer with a single circular pore of radius R . The stretching free energy of state (Un) is equal to

$$F_{\text{un}} = \frac{1}{2} K_A (A_{||} - A_0)^2 / A_0 \quad (\text{State (Un)}) \quad (11)$$

as follows from Eq (2). The free energy F_{po} of the porated bilayer state (Po) has two contributions: (i) the stretching energy of the remaining bilayer with area $A_{||} - A_0 - \pi R^2$ and (ii) the edge energy of the pore with circumference $2\pi R$ and edge tension λ , which implies^{22,23}

$$F_{\text{po}} = \frac{1}{2} K_A (A_{||} - A_0 - \pi R^2)^2 / A_0 + 2\pi R \lambda \quad (\text{State (Po)}). \quad (12)$$

The free energy difference ΔF between (Po) and (Un) is

$$\Delta F(R) = F_{\text{po}} - F_{\text{un}} = 2\pi R \lambda - \pi \Sigma R^2 + \frac{1}{2} \pi^2 K_A R^4 / A_0 \quad (13)$$

with the mechanical tension $\Sigma = K_A (A_{||} - A_0) / A_0 = K_A \Delta a$ as in Eq (1). The free energy difference ΔF depends on the pore radius



R and involves two material parameters, the area compressibility modulus K_A and the edge tension λ . In addition, ΔF depends on the mechanical tension Σ , which is proportional to the relative area dilation $\Delta a = (A_{\parallel} - A_0)/A_0$ and, thus, represents a geometric control parameter that can be varied by changing the base area A_{\parallel} of the simulation box.

When the bilayer is compressed by a negative value of Σ , all terms on the right hand side of Eq (13) are positive and $\Delta F(R)$ is a monotonically increasing function of R , which implies that the unporated bilayer state (Un) is always the state of lower free energy. On the other hand, when the bilayer is stretched by a positive value of Σ , the free energy difference $\Delta F(R)$ in Eq (13) develops an inflection point for mechanical tension

$$\Sigma_1 = c_1 K_A^{1/3} \lambda^{2/3} / A_0^{1/3} \quad (14)$$

with $c_1 = 3\pi^{1/3}/2^{2/3} = 2.77$. For $\Sigma > \Sigma_1$, the inflection point splits up into a maximum and a minimum of $\Delta F(R)$. The minimum at $R = R_2$ describes a bilayer with a stable pore, which becomes more stable with increasing Σ until it attains the same free energy as the unporated state. This coexistence of the porated and unporated state is obtained for the mechanical tension

$$\Sigma_2 = c_2 K_A^{1/3} \lambda^{2/3} / A_0^{1/3} \quad (15)$$

with $c_2 = 2^{1/3}c_1 = 3.49$. For $\Sigma > \Sigma_2$, the porated bilayer provides the state of lower free energy.

Therefore, as we vary the mechanical bilayer tension Σ within a symmetric bilayer, the stretched bilayer has no pore for $\Sigma \leq \Sigma_1$, forms a metastable pore for $\Sigma_1 < \Sigma < \Sigma_2$, and a globally stable pore for $\Sigma > \Sigma_2$. The threshold values Σ_1 and Σ_2 as given by Eqs (14) and (15) are in good agreement with the results of molecular dynamics simulations.²³ It is important to note that both tension thresholds Σ_1 and Σ_2 are proportional to $1/A_0^{1/3}$ and thus go to zero for large bilayer area A_0 . As a consequence, sufficiently large bilayers can always lower their free energy by forming a stable pore,

3.4 Leaflet tensions of tensionless planar bilayers

In order to avoid pore formation and bilayer rupture, we will now focus on tensionless bilayers with $\Sigma = \Sigma_{ul} + \Sigma_{ll} = 0$ or $\Sigma_{ll} = -\Sigma_{ul}$, corresponding to OLT states of the bilayer. For bilayers with a total lipid number $N_{ll} + N_{ul} = 1682$ as in Fig 3, the tensionless bilayers or OLT states are given by the black data points in this figure. The corresponding leaflet tensions are displayed in Fig 4 as functions of the lipid number N_{ul} assembled in the upper leaflet. As shown in this figure, tensionless leaflets are obtained for the symmetric bilayer with $N_{ll} = N_{ul} = 841$.

The stability regime for planar tensionless bilayers as displayed in Fig 4 is bounded by two instability lines, the left instability line at $N_{ul} = 957$ and $N_{ll} = 725$ as well as the right instability line at $N_{ul} = 725$ and $N_{ll} = 957$. For $N_{ul} > 957$, the lipids in the compressed upper leaflet start to undergo flip-flops into the stretched lower leaflet. For $N_{ul} < 725$, the lipids in the compressed lower leaflet start to flip-flop into the stretched upper leaflet. In addition to flip-flops of lipid molecules, tensionless planar bilayers




Fig. 4 Leaflet tensions of tensionless planar bilayers: Upper leaflet tension Σ_{ul} (blue) and lower leaflet tension Σ_{ll} (red) versus lipid number N_{ul} for constant total lipid number $N_{ul} + N_{ll} = 1682$.¹⁹ The lipid number N_{ul} is reduced by moving lipids from the upper to the lower leaflet, thereby increasing the upper leaflet tension and decreasing the lower one. Both leaflet tensions vanish for $N_{ul} = 841$ (vertical dashed line) which defines the relaxed reference state of the planar bilayers. The green data display the bilayer tension $\Sigma = \Sigma_{ul} + \Sigma_{ll}$, which is close to zero. For $945 \geq N_{ul} \geq 737$, the bilayer remains stable but becomes unstable (i) for $N_{ul} \geq 957$ via flip-flops from the compressed upper to the stretched lower leaflet and (ii) for $N_{ul} \leq 725$ via flip-flops from the compressed lower to the stretched upper leaflet.

undergo structural instabilities outside of the stability regime in Fig 4. Data for lipid flip-flops and structural instabilities are displayed in Figs S5[†] and S6[†]. On the other hand, if a multicomponent and tensionless lipid bilayer contains a lipid species such as cholesterol that undergoes frequent flip-flops, these flip-flops act to decrease the stress asymmetry between the bilayer leaflets (Sect S5[†] and Fig S7[†]).²⁴

3.5 Fluctuation tension of planar bilayers

In this last subsection on planar bilayers, we briefly look at the effect of the membrane's shape fluctuations. These fluctuations can be decomposed into Fourier modes, which leads to the fluctuation spectrum

$$S(q) \approx \frac{1}{A_{\parallel}} \frac{k_B T}{\Sigma_{fl} q^2 + \kappa q^4} \quad (16)$$

for small wavenumber q . The prefactor of the q^2 -term defines the fluctuation tension Σ_{fl} . For large q -values, the spectrum is changed by molecular protrusions.

One interesting and open question is the relation between the fluctuation tension Σ_{fl} in Eq (16) and the mechanical bilayer tension Σ . In our original molecular dynamics study of the fluctuation spectrum¹⁴, we assumed that vanishing mechanical tension, $\Sigma = 0$, implies vanishing fluctuation tension, $\Sigma_{fl} = 0$. This assumption has been confirmed by the fluctuation spectra of planar bilayers with two lipid components, see simulation data in Fig S8[†].

On the other hand, several molecular dynamics simulations found small differences between the numerical values of the two tensions Σ and Σ_{fl} ²⁵⁻²⁷, in agreement with the theory described in Ref 28. However, the most recent study²⁹, which addressed this issue by Monte Carlo simulations, came to the conclusion that the two tensions are, in fact, identical, that is, that $\Sigma_{fl} = \Sigma$, in agreement with our original assumption¹⁴.



4 Membrane tensions of nanovesicles

For a closed vesicle, we distinguish the inner from the outer leaflet by the subscripts il and ol and denote the corresponding leaflet tensions by Σ_{il} and Σ_{ol} . The bilayer tension Σ is again equal to the sum of the two leaflet tensions, that is, $\Sigma = \Sigma_{il} + \Sigma_{ol}$.

4.1 Mechanical tensions of spherical bilayers and leaflets

For a spherical bilayer, the stress profile depends on the radial coordinate r , which measures the distance from the center of the sphere. The corresponding mechanical tension Σ can be obtained from the stress profile $s = s(r)$ via

$$\Sigma = \int_0^{+\infty} dr s(r), \quad (17)$$

The bilayer tension Σ of a spherical vesicle depends on the vesicle volume in close analogy to the experimental procedures of osmotic deflation and inflation. Osmotic deflation leads to a reduction of the bilayer tension whereas osmotic inflation increases this tension.

The inner leaflet occupies a certain range of r -values, which lie below the r -values for the outer leaflet. The molecular interface between these two leaflets defines the midsurface of the spherical bilayer at $r = r_{\text{mid}}$.¹ The leaflet tensions are then obtained from

$$\Sigma_{il} = \int_0^{r_{\text{mid}}} dr s(r) \quad \text{and} \quad \Sigma_{ol} = \int_{r_{\text{mid}}}^{+\infty} dr s(r). \quad (18)$$

4.2 Leaflet tension space for spherical bilayers

Now, consider an ensemble of spherical vesicles, which all have the same total lipid number $N_{il} + N_{ol}$. The corresponding leaflet tension space is displayed in Fig 5 for total lipid number $N_{il} + N_{ol} = 2525$. The origin of this space, corresponding to the relaxed reference state with tensionless leaflets $\Sigma_{il} = \Sigma_{ol} = 0$, is obtained for $N_{il} = 840$ lipids in the inner leaflet and $N_{ol} = 1685$ lipids in the outer leaflet. Therefore, the bilayer state with two tensionless leaflets is obtained when the outer lipid number N_{ol} is about twice as large as the inner lipid number N_{il} .

Because the reference state has a large lipid number asymmetry, it is difficult to identify the ELT states with $\Sigma_{ol} = \Sigma_{il}$ for spherical bilayers. In contrast, it is relatively simple to generate the OLT states with $\Sigma_{ol} = -\Sigma_{il}$ from the relaxed reference state with $\Sigma_{ol} = \Sigma_{il} = 0$ by (i) reshuffling lipids between the two leaflets and (ii) adjusting the vesicle volume to obtain a tensionless bilayer.^{18,19,30} The resulting OLT states correspond to the black data points in Fig 5. The different OLT states are distinguished by the stress asymmetry

$$\Delta\Sigma_{\text{ve}} \equiv \Sigma_{ol} - \Sigma_{il} \quad (19)$$

which increases in magnitude as we move away from the reference state with $\Sigma_{il} = \Sigma_{ol} = 0$ along the black data points in Fig 5.

To generate the green data points for vesicle inflation or deflation (VID) in Fig. 5, we start again from the reference state with $\Sigma_{il} = \Sigma_{ol} = 0$ but now change the vesicle volume in order to increase or decrease the bilayer tension, thereby mimicking the experimental procedure of osmotic inflation or deflation. Note




Fig. 5 Leaflet tension space for vesicle bilayers with a total number of $N_{il} + N_{ol} = 2525$ lipids in both leaflets. The two coordinates are the leaflet tensions Σ_{il} and Σ_{ol} in the inner and outer leaflets. The reference state with tensionless leaflets, corresponding to $\Sigma_{il} = \Sigma_{ol} = 0$, is obtained for a vesicle bilayer with $N_{il} = 840$ lipids in the inner leaflet and $N_{ol} = 1685$ lipids in the outer one.¹⁸ Negative and positive leaflet tensions describe compressed and stretched leaflets. The black data correspond to elastic OLT deformations obtained from the reference state by reshuffling lipids from one leaflet to the other and adjusting the vesicle volume to obtain tensionless bilayers with $\Sigma = \Sigma_{il} + \Sigma_{ol} = 0$. The green data represent the elastic deformations arising from changes in vesicle volume, corresponding to vesicle inflation or deflation (VID).

that the green data do not follow the main diagonal, in contrast to the equal leaflet tension (ELT) data in Fig 3, but become parallel to the main diagonal for large leaflet tensions, which implies a constant stress asymmetry $\Delta\Sigma_{\text{ve}}$.

4.3 Leaflet tensions of tensionless vesicle bilayers

The leaflet tensions of tensionless vesicle bilayers, which are assembled from a total number of $N_{il} + N_{ol} = 2875$ lipids, are displayed in Fig 6. For these vesicle bilayers, the relaxed reference state with tensionless leaflets is obtained for $N_{ol} = 1921$ lipids in the outer leaflet and $N_{il} = 954$ lipids in the inner leaflet, corresponding to the dashed vertical line in Fig 6. Thus, when both leaflet tensions vanish, the lipid number N_{ol} in the tensionless outer leaflet is more than twice as large as the lipid number N_{il} in the tensionless inner leaflet.

The white stability regime in Fig 6 is bounded by two instability lines. When we reach the left instability line at $N_{ol} = 2105$, lipids start to flip-flop from the compressed outer leaflet to the stretched inner leaflet. Likewise, at the right stability line with $N_{ol} = 1755$, the compressed inner leaflet becomes unstable with respect to flip-flops from the inner to the outer leaflet. The cumulative distribution function for the first flip-flop is displayed in Fig S9[†]. In addition to flip-flops, tensionless vesicle bilayers undergo structural instabilities outside of their stability regime as illustrated in Fig S10[†].






Fig. 6 Leaflet tensions for tensionless vesicle bilayers with a total number of $N_{il} + N_{ol} = 2875$ lipids. The vesicles have a diameter of $23.8d$ or 19 nm . The blue and red data represent the outer and inner leaflet tensions.¹⁹ The green data correspond to the bilayer tension $\Sigma = \Sigma_{ol} + \Sigma_{il}$, which is close to zero. During the run time of the simulations, we observed no flip-flops within the stability regime (white), corresponding to $1775 \leq N_{ol} \leq 2095$. The left vertical line at $N_{ol} = 2105$ represents the instability line at which the lipids start to undergo flip-flops from the compressed outer to the stretched inner leaflet. The right vertical line at $N_{ol} = 1775$ represents the instability line at which the lipids start to undergo flip-flops from the compressed inner to the stretched outer leaflet.

4.4 Volume per lipid and lateral volume compressibility

The elastic response of a planar bilayer to mechanical tension can be described in terms of area per lipid or in terms of volume per lipid. In order to determine area per lipid for nanovesicles, we need to introduce some projection onto the curved surfaces of the vesicle bilayers, which represents an ambiguous procedure. Such an ambiguity can be avoided by using the volume per lipid, v , to characterize the lipid bilayer.¹⁸

The volume per lipid can be obtained from the following procedure based on Voronoi tessellation. In the coarse-grained molecular model, the molecules are built up from different types of spherical beads. Voronoi tessellation assigns a polyhedral cell to each bead, such that all points in this cell are closer to the center of the chosen bead than to the center of any other bead. In Fig S4[†], we illustrate the tessellation method for a planar bilayer and for an individual lipid molecule. Next, the volumes V_{il} and V_{ol} of the inner and outer leaflets are computed by summing up the volumes of all polyhedral cells that belong to the respective leaflet. Finally, the volumes per lipid, v_{il} and v_{ol} , of the inner and outer leaflets are then obtained by dividing the leaflet volumes by the lipid numbers, N_{il} and N_{ol} , which leads to

$$v_{il} = \frac{V_{il}}{N_{il}} \quad \text{and} \quad v_{ol} = \frac{V_{ol}}{N_{ol}}. \quad (20)$$

These lipid volumes vary with the leaflet tensions Σ_{il} and Σ_{ol} in the inner and outer leaflets.

One example for this dependence is shown in Fig. 7 for a vesicle bilayer assembled from a total number of $N_{il} + N_{ol} = 1500$ lipids. Inspection of this figure shows that the optimal lipid volumes for the outer and inner leaflet have almost identical but slightly different values as given by¹⁸ $v_{il,0} = (3.5790 \pm 0.0004)d^3$ for the inner leaflet and by $v_{ol,0} = (3.5612 \pm 0.0003)d^3$ for the outer leaflet of the vesicle, corresponding to the vertical dashed lines in Fig. 7. Therefore, the optimal lipid volumes for the inner




Fig. 7 Leaflet tensions versus volume per lipid for nanovesicles with a total number of $N_{il} + N_{ol} = 1500$ lipids.¹⁸ Leaflet tensions $\Sigma_{le} = \Sigma_{ol}$ and $\Sigma_{le} = \Sigma_{il}$ of the outer (black triangles) and inner leaflets (red triangles) versus volumes per lipid, $v_{le} = v_{il}$ and $v_{le} = v_{ol}$, in the two leaflets of the tensionless bilayers. The two dotted vertical lines are located at the optimal lipid volumes $v = v_{ol}^0 \approx 3.56d^3$ and $v = v_{il}^0 \approx 3.58d^3$ of the outer and inner leaflet.

and outer leaflets differ by less than one percent.

The deviations of the lipid volumes from their optimal values define the relative volume dilations

$$\Delta v_{il} \equiv \frac{v_{il} - v_{il}^0}{v_{il}^0} \quad \text{and} \quad \Delta v_{ol} \equiv \frac{v_{ol} - v_{ol}^0}{v_{ol}^0} \quad (21)$$

for the inner and outer leaflets. The elastic response of the two leaflets to the leaflet tensions Σ_{il} and Σ_{ol} is then described by the tension–volume relations

$$\Sigma_{il} \approx B_{il} \Delta v_{il} = B_{il} \frac{v_{il} - v_{il}^0}{v_{il}^0} \quad (22)$$

and

$$\Sigma_{ol} \approx B_{ol} \Delta v_{ol} = B_{ol} \frac{v_{ol} - v_{ol}^0}{v_{ol}^0} \quad (23)$$

to first order in the relative volume dilations, which define the lateral volume compressibilities B_{il} and B_{ol} . These two elastic moduli are significantly different for the two leaflets as follows from the data in Fig 7.¹⁸

4.5 Tension-induced fusion of nanovesicles

Membrane fusion of two nanovesicles can be induced by their bilayer tensions or by their leaflet tensions. A particularly interesting fusion process is observed for tensionless bilayers with $\Sigma = 0$ and $\Sigma_{ol} = -\Sigma_{il}$, which avoids bilayer rupture as an alternative pathway to fusion.

Membrane fusion induced by bilayer tension. Experimentally, tension-induced fusion was originally studied by osmotic swelling of vesicles that adhere to a planar bilayer.³¹ The molecular mechanisms underlying tension-induced fusion have been visualized and elucidated by molecular dynamics simulations.^{32–34} More recently, several experimental studies provided additional evidence for tension-induced fusion^{35–38}, which can be strongly enhanced by electrostatic interactions between oppositely charged membranes³⁶. In the simulations, the close proximity of two lipid bilayers can lead to a variety of alternative out-






Fig. 8 Fusion of nanovesicle with planar bilayer: The vesicle bilayer consists of 6869 lipids (orange heads, yellow chains) and has a diameter of about 30 nm, while the planar bilayer contains 6911 lipids (red heads, green chains). The water beads originally inside the vesicle are blue, the water beads in the exterior compartment are not shown for clarity. The figure displays four time lapse snapshots which illustrate the development of the fusion process from 78.5 ns, when the first contact between the two membranes occurs and lipid tails start to undergo interbilayer flip-flops, until the opening of the fusion pore after 1334 ns.³³

comes such as stable hemifused bilayers and bilayer rupture but fusion events are always observed within a certain range of positive bilayer tensions.

One example for membrane fusion as observed in molecular dynamics simulations is depicted in Fig 8 for a nanovesicle in contact with a planar bilayer.^{33,39} In order to visualize the fusion pathway, the lipids in the planar bilayer have red head groups and green tails whereas the lipids in the vesicle have orange head groups and yellow tails.

The observed fusion pathway proceeds via three distinct sub-steps. After the vesicle and the planar bilayer have come into contact at time $t = 78.5$ ns, the vesicle starts to spread onto the bilayer, thereby forming a highly curved membrane segment along the contact line with the planar bilayer. The lipids along the contact line then undergo interbilayer flip-flops from the vesicle to the planar bilayer, see the yellow lipid tails that appear in the lower leaflet of the planar bilayer. These flip-flops lead to intermixing and partial fusion of the two bilayers, followed by the nucleation of a bean-shaped hemifusion diaphragm at about $t = 1177$ ns. The bean-shape of the diaphragm can be clearly seen when one looks at sections across the planar bilayer, which are perpendicular to the side views in Fig. 8. Finally, this diaphragm ruptured close to the contact line at $t = 1334$ ns, thereby opening up a strongly asymmetric fusion pore that provides a water channel between the interior vesicle solution (blue) and the water compartment (white) above the planar bilayer.

The fusion process displayed in Fig 8 was induced by increasing the area per lipid, a , of the planar bilayer, thereby increasing the mechanical tension Σ experienced by this bilayer. The resulting fusion process competes with several alternative pathways such as membrane adhesion, hemifusion, and rupture. Because of these alternative pathways, the fraction of successful fusion




Fig. 9 Two identical vesicles with tensionless bilayers undergo fusion, when their outer leaflets are stretched by a sufficiently large leaflet tension Σ_{ol} and their bilayers experience a sufficiently large stress asymmetry. Each vesicle contains $N_{il} = 4500$ lipids in its inner and $N_{ol} = 5600$ lipids in its outer leaflet, corresponding to the outer leaflet tension $\Sigma_{ol} = +1.02k_B T/d^2$ and the inner leaflet tension $\Sigma_{il} = -1.02k_B T/d^2$. The vesicles come into contact at time $t = 0$ and undergo fusion within $0.3 \mu s$.¹

events, which defines the fusion probability, depends strongly on the area per lipid, a , and on the mechanical tension Σ within the planar bilayer. Indeed, as shown in Fig S11[†], the fusion probability increases linearly with increasing $a \sim \Sigma$, reaches a maximum, and then decreases again for larger values of $a \sim \Sigma$. The decay of the fusion probability for large a is caused by bilayer rupture.

Vesicle fusion driven by leaflet tensions. Bilayer rupture is strongly suppressed when we consider tensionless bilayers with vanishing mechanical tension, $\Sigma = 0$, corresponding to one stretched and one compressed leaflet. Somewhat surprisingly, two nanovesicles with $\Sigma = 0$ also undergo fusion, provided the outer leaflet is stretched by a sufficiently large leaflet tension Σ_{ol} . One example for such a fusion process is displayed in Fig 9.

The fusion of nanovesicles with tensionless bilayers but stretched outer leaflets can be understood as follows. First, within the emerging contact area, the shielding of the hydrophobic cores of the two bilayers is reduced by stretching the two outer leaflets and increasing their areas and volumes per lipid. Second, both leaflet tensions drive the subsequent opening of the fusion pore. Indeed, the stretched outer leaflet tries to reduce its area, thereby pulling on the emerging fusion pore, whereas the compressed inner leaflet tries to increase its area, thereby curving the rim of the pore. Furthermore, recent molecular dynamics simulation show that this fusion process proceeds without bilayer rupture and without leakage of vesicle contents (Bartosz Rozycki, Rikhia Ghosh, and Reinhard Lipowsky, in preparation).

5 Membrane tensions of giant vesicles

In this section, we will address membrane tensions of giant vesicles and will focus on multispherical and two-sphere shapes as well as on different shapes of tubular membranes. All of these shapes involve closed membrane necks, which also represent the necessary initial step for vesicle division. Additional aspects of multispherical shapes are discussed in Sect S6[†], more details on positive two-sphere shapes are provided in Sect S7[†], which also describes the experimentally observed division of positive two-sphere shapes⁴⁰. Negative two-sphere shapes are further examined in Sect S8[†].






Fig. 10 (a-d) $(1+N_s)$ -spheres for positive spontaneous curvatures, consisting of one large sphere and N_s small spheres, all with positive mean curvature. The scale bars in (a) and (d) are $10\mu\text{m}$, the scale bar in (a) applies to (a-c); and (e-h) $(1+N_s)$ -spheres for negative spontaneous curvatures, consisting of one large sphere with positive mean curvature and N_s small spheres with negative mean curvature. The membrane necks of the positive and negative multispheres have a positive and negative effective mean curvature, respectively, as follows from Eq (24). The scale bar in (e) is $5\mu\text{m}$ and applies to (e-h).⁴¹

5.1 Multispherical shapes of giant vesicles

Multispherical shapes of giant vesicles can be obtained by exposing the two leaflets of the vesicle membranes to two different sugar solutions which generate a spontaneous curvature within the membranes.^{41,42} If the inner leaflet is in contact with sucrose and the outer leaflet exposed to glucose, the spontaneous curvature m is positive. On the other hand, the GUV membrane acquires a negative spontaneous curvature, $m < 0$, when the interior leaflet is exposed to glucose whereas the outer leaflet is in contact with sucrose. Some examples for the resulting multispheres are displayed in Fig 10.

Multispheres consist of large and small spheres. Inspection of Fig 10 reveals that each multisphere involves only two types of spheres, a large sphere with radius R_l and positive mean curvature $M_l = +1/R_l$ as well as small spheres with identical radius R_s and identical mean curvature $M_s = \pm 1/R_s$. This general property of multispheres is a direct consequence of the local shape equation for spherical membrane segments as given by Eq (S24)[†], which is quadratic in the constant mean curvature of the spherical segment. For positive spontaneous curvature, multispheres can also consist of several equally sized spheres, corresponding to a degenerate (double) root of the quadratic shape equation, see the example in Fig 12c below.

Limit shapes versus persistent shapes. Each multispherical shape can be characterized by its stability regime, that is, by a certain subregion of the morphology diagram, which depends on two dimensionless shape parameters, the volume-to-area ratio \bar{v} and the rescaled spontaneous curvature \bar{m} as defined by Eqs (S9)[†] and (S10)[†]. The stability regime of each multisphere is bounded by two lines of *limit shapes*. When we leave the stability regime by crossing a line of limit shapes, at least one closed membrane neck opens up. On the other hand, away from these limit shapes, the multispherical shapes represent *persistent shapes* that depend only on \bar{v} but are independent of \bar{m} .⁴³

Positive and negative multispheres. Each closed neck between two connected spheres i and j with mean curvature M_i and M_j can be characterized by its effective mean curvature as given by⁴³

$$M_{ij}^{\text{eff}} \equiv \frac{1}{2} (M_i + M_j) = \frac{1}{2} \left(\pm \frac{1}{R_i} \pm \frac{1}{R_j} \right). \quad (24)$$

This effective curvature represents a purely geometric quantity that can be directly obtained from microscopy images as in Fig 10. Therefore, we can distinguish positive necks with $M_{ij}^{\text{eff}} > 0$ from negative necks with $M_{ij}^{\text{eff}} < 0$.

In fact, for each multisphere, all necks have the same sign. Indeed, all membrane necks are positive for each multisphere in Fig 10a-d and negative for each multisphere in Fig 10e-h. As a consequence, we can distinguish positive multispheres with positive membrane necks from negative multispheres with negative membrane necks.

Mechanical membrane tension of multispheres. The constant mean curvature M_{sp} of a spherical membrane segment is obtained by solving the local shape equation as given by Eq (S24)[†]. This equation is quadratic in M_{sp} and depends on ΔP and Σ . The two solutions of this equation are equal to the mean curvatures of the large and the small spheres, M_l and M_s , which build up the multispheres as in Fig 10. It is possible to invert these solutions to obtain ΔP and Σ , which leads to the mechanical tension⁴³

$$\Sigma = 2\kappa m (M_l + M_s - m) = 2\kappa m \left(\frac{1}{R_l} \pm \frac{1}{R_s} - m \right). \quad (25)$$

The \pm sign applies to positive and negative multispheres, which are characterized by small spheres with positive and negative mean curvature, respectively. The relationship in Eq (25) remains valid for the degenerate case of a single solution with $M_l = M_s = M_*$ and $R_l = R_s = R_*$.

The mechanical tension as given by Eq (25) involves both the curvature-elastic parameters κ and m as well as on mean curvatures M_l and M_s (or M_*) of the large and small spheres. Therefore, Eq (25) directly demonstrates that the mechanical tension Σ depends on the size and shape of the multispherical vesicle.

5.2 Two-sphere shapes of giant vesicles

The simplest multispheres are provided by two spheres connected by a closed membrane neck as shown in panels a and e of Fig 10. The positive two-sphere shape in Fig 10a consists of one large sphere and one smaller spherical out-bud, the negative two-sphere shape in Fig 10e of one large sphere and one smaller spherical in-bud. The stability regime of the positive two-spheres is described in the next paragraph and displayed in Fig 11, the stability regime of the negative two-spheres in Sect S8[†] and Fig S13[†].

Stability regime for positive two-sphere shapes. For positive spontaneous curvature, the two-sphere shape in Fig 10a involves a positive *ls*-neck between the large and the small sphere. The condition for this *ls*-neck to be stably closed is given by the in-



equality⁴³

$$m \geq M_{ls}^{\text{eff}} = \frac{1}{2} (M_l + M_s) = \frac{1}{2} \left(\frac{1}{R_l} + \frac{1}{R_s} \right) > 0. \quad (26)$$

In the limiting case, in which this inequality becomes an equality, we obtain the neck closure condition that determines the line of limit shapes L_{1+1}^{pos} , see the upper boundary line of the stability regime in Fig 11.

The lower boundary line of the stability regime in Fig 11 is provided by another line of limit shapes, denoted by L_{2*} and consisting of two equally sized spheres with radius $R_l = R_s = R_* = R_{ve}/\sqrt{2}$. In the morphology diagram, the L_{2*} -line is located at $\bar{v} = \bar{v}_{2*} = 1/\sqrt{2}$ and $\bar{m} \geq \bar{m}_{2*} = \sqrt{2}$.

The two lines of limit shapes, L_{1+1}^{pos} and L_{2*}^{pos} , meet at the corner point with $\bar{v} = \bar{v}_{2*} = 1/\sqrt{2}$ and $\bar{m} = \bar{m}_{2*} = \sqrt{2}$. At this corner point, the limit shape consists of two equally sized spheres with mean curvature $M_* = 1/R_* = m_{2*}$, which implies zero bending energy, $E_{be} = 0$, as follows from Eq (S3)[†].

Mechanical membrane tension of positive two-spheres.

Along the line of limit shapes L_{1+1}^{pos} in Fig 11, the neck closure condition $M_l + M_s = 2M_{ls}^{\text{eff}} = 2m$ implies the mechanical tension

$$\Sigma = 2\kappa m^2 = 2\kappa \left(M_{ls}^{\text{eff}} \right)^2 \quad \text{for all limit shapes } L_{1+1}^{\text{pos}} \quad (27)$$

as follows from Eq (25). The geometry of the limit shape L_{1+1}^{pos} changes with the spontaneous curvature m because the effective mean curvature $M_{ls}^{\text{eff}} = m$ of the ls -neck increases with increasing m . The relation in Eq (25) also implies that

$$\Sigma = 0 \quad \text{for } m = M_l + M_s, \quad (28)$$

which is valid for the persistent shapes along the red dashed line in Fig 11.




Fig. 11 Stability regime (light blue) for positive (1+1)-spheres with positive spontaneous curvature \bar{m} . The stability regime is bounded by two lines (dark blue) of limit shapes, L_{1+1}^{pos} and L_{2*}^{pos} , which meet at the corner point (purple circle) with $\bar{m} = \sqrt{2}$ and $\bar{v} = 1/\sqrt{2}$. The persistent (1+1)-spheres along the red dashed line are characterized by tensionless membranes with mechanical tension $\Sigma = 0$; this tension is positive and negative to the left and to the right of the red dashed line. The inset displays a persistent (1+1)-shape that remains unchanged when we vary the spontaneous curvature along the horizontal dashed line (orange). This shape invariance is a direct consequence of the area and volume relations in Eqs (S27)[†] and (S28)[†].

Along the line of limit shapes L_{2*}^{pos} with $\bar{v} = \bar{v}_{2*} = 1/\sqrt{2}$ and $\bar{m} \geq \bar{m}_{2*} = \sqrt{2}$, the mean curvature M_* of the equally sized spheres is equal to $M_* = m_{2*} = \sqrt{2}/R_{ve}$. It then follows from Eq (25) with $M_l = M_s = M_*$ that the associated membrane mechanical tension is given by

$$\Sigma = 2\kappa m \left(\frac{2\sqrt{2}}{R_{ve}} - m \right) \quad \text{for the limit shapes } L_{2*}^{\text{pos}}. \quad (29)$$

This tension vanishes at $m = 2m_{2*} = 2\sqrt{2}/R_{ve}$, which represents the lower end point of the dashed red line in Fig 11. Furthermore, the mechanical tension Σ as given by Eq (29) behaves as $\Sigma \approx -2\kappa m^2$ for large m , that is, the mechanical tension Σ becomes large and negative for large and positive spontaneous curvature m .

As shown in Fig 11, the L_{1+1} -line meets the L_{2*} -line of two equally sized spheres at the corner point with $\bar{m} = \bar{m}_{2*} = \sqrt{2}$ and $\bar{v} = \bar{v}_{2*} = 1/\sqrt{2}$ where the mechanical tension Σ has the value

$$\Sigma = 2\kappa m_{2*}^2 = 4\kappa/R_{ve}^2 \quad (\text{corner point}). \quad (30)$$

Division of positive two-sphere vesicles. Closed membrane necks of multispherical vesicles also play a pivotal role for the division of these vesicles. As an example, consider the positive two-sphere vesicle in the inset of Fig 11. This shape remains unchanged when we increase the spontaneous curvature m of the vesicle membrane. However, as explained in Sect S7.2[†] such an increase of m also increases the constriction force f^{pos} acting against the positive closed neck of the two-sphere vesicle, as described by Eq (S30)[†]. A sufficiently large constriction force leads to the cleavage of the closed neck and to the division of the two-sphere vesicle into two daughter vesicles as observed experimentally for GUVs⁴⁰. Because the constriction force is a local neck property, such a force is present at each closed neck of all multispheres.

5.3 Different shapes of tubular membranes

Tubular membranes and vesicles can exhibit different shapes, corresponding to multispherical, unduloidal, and cylindrical shapes.

Multispherical shapes of membrane tubes. We first consider multispherical shapes of membrane tubes as shown in Fig 12. In Fig 12b, the tube is part of a (1+14)-sphere and consists of 14 small spheres emanating from a somewhat larger sphere. In Fig 12c, the multispherical tube is built up from 15 equally sized spheres, corresponding to a (15*)-sphere. The stability regime for the (1+14)-spheres is displayed in Fig 12a as the light blue subregion of the morphology diagram. As for (1+1)-spheres, the stability regime for the (1+14)-spheres is bounded by two lines of limit shapes, denoted by L_{1+14}^{pos} and L_{15*}^{pos} .

The line of limit shapes L_{1+14}^{pos} in Fig 12a is given by

$$\bar{v} = \bar{v}_{1+14}(\bar{m}) = \left(1 - \frac{14}{\bar{m}^2} \right)^{3/2} + \frac{14}{\bar{m}^3}, \quad (31)$$

the line of limit shapes L_{15*}^{pos} by

$$\bar{m} \geq \bar{m}_{15*} \equiv \sqrt{15} \quad \text{and} \quad \bar{v} = \bar{v}_{15*} \equiv 1/\sqrt{15} \quad (32)$$






Fig. 12 (a) Stability regime (light blue) for positive (1+14)-spheres. The stability regime is bounded by two lines of limit shapes (dark blue), L_{1+14}^{pos} and $L_{15^*}^{\text{pos}}$. These two lines meet at the corner point with $\bar{m} = \bar{m}_{15^*} = \sqrt{15} = 3.87$ and $\bar{v} = \bar{v}_{15^*} = 1/\sqrt{15} = 0.258$. The persistent (1+14)-shapes along the red dashed line are enclosed by membranes with vanishing mechanical tension, $\Sigma = 0$; (b) Example for a GUV that forms a (1+14)-sphere consisting of one somewhat larger sphere at the left end and a multispherical tube of 14 small spheres; and (c) Example for a GUV forming a (15 *)-sphere with 15 equally sized spheres. The images in (b) and (c) are from Ref 41, both scale bars are 10 μm .

as follows from the area and volume relations in Eqs (S27) and (S28) with small-sphere radius $r_s = 1/\bar{m}$. The red dashed line in Fig 12a provides the location for vesicle membranes with vanishing mechanical tension, $\Sigma = 0$. Along this line, the spontaneous curvature m is equal to $2M_{15}^{\text{eff}}$, which can be combined with Eqs (S27) and (S28) to obtain a parametric plot⁴².

The small spheres of the (1+14)-spheres in Fig 12b transform into small prolates along an instability line as given by the relationship

$$r_s - \frac{r_s^2}{\sqrt{1 - 14r_s^2}} = \frac{3}{\bar{m}} \quad (33)$$

between the small-sphere radius r_s and the spontaneous curvature \bar{m} . This equation has no solution for $\bar{m} < \bar{m}_{ss} \equiv 21.47$, one solution for $\bar{m} = \bar{m}_{ss}$, and two solutions for $\bar{m} > \bar{m}_{ss}$. Therefore, the small spheres of the persistent (1+14)-spheres are stable for $\bar{m} < \bar{m}_{ss} = 21.47$ but transform into small prolates for $\bar{m} > \bar{m}_{ss}$ and a certain \bar{m} -dependent range of \bar{v} -values. At $\bar{m} = \bar{m}_{ss}$, the instability line emerges from the single point $(\bar{m}_{ss}, \bar{v}_{ss}) = (21.47, 0.4035)$, which opens up into a parabola-like line for $\bar{m} > \bar{m}_{ss}$. For large \bar{m} , the upper and the lower branch of this parabola-like line approach the two lines of limit shapes, L_{1+14}^{pos} and $L_{15^*}^{\text{pos}}$, respectively. This bifurcation is located outside of the morphology diagram displayed in Fig 12a.

Multispherical tubes via fusion of proteoliposomes. Membrane nanotubes have been prepared by the fusion of proteoliposomes that contain certain membrane proteins, which hydrolyse GTP.^{44,45} Some insight into this process of nanotube formation from proteoliposomes can be obtained from the following thought experiment. Thus, let us consider N_* proteoliposomes that undergo fusion to form an (N_*)-multisphere. In the absence of GTP, the individual proteoliposome are taken to be relaxed in the sense that the mean curvature of the vesicle membranes are close to the membrane's spontaneous curvature m , which implies

that the radius R of the proteoliposomes is close to $1/m$ and that their bending energy as given by Eq (S3)[†] is close to zero. The dimensionless shape parameters \bar{v} and \bar{m} in Eqs. (S9)[†] and (S10)[†] are then equal to $\bar{v} = \bar{v}_1 = 1$ and $\bar{m} = \bar{m}_1 = 1$ for each proteoliposome.

When we add GTP, the proteoliposomes start to fuse and to form multispherical tubes. More precisely, the fusion of N_* proteoliposomes leads to an (N_*)-multisphere with the shape parameters $\bar{v} = \bar{v}_{N_*} \equiv 1/\sqrt{N_*}$ and $\bar{m} = \bar{m}_{N_*} \equiv \sqrt{N_*}$, provided both the total membrane area and the total vesicle volume are conserved during the fusion process. For $N_* = 2$, the shape parameters $\bar{v} = \bar{v}_{2^*} = 1/\sqrt{2}$ and $\bar{m} = \bar{m}_{2^*} = \sqrt{2}$ represent the corner point of the stability regime in Fig 11, at which the two lines of limit shapes, L_{1+1}^{pos} and $L_{2^*}^{\text{pos}}$, meet. For $N_* = 15$, the shape parameters $\bar{v} = \bar{v}_{15^*} \equiv 1/\sqrt{15}$ and $\bar{m} = \bar{m}_{15^*} \equiv \sqrt{15}$ define the location of the corner point in Fig 12a.

Unduloidal shapes of membrane tubes. So far, super-resolution microscopy has not been used to resolve the shapes of nanotubes reconstituted from proteoliposomes *in vitro*. However, structured illumination microscopy has been applied to the nanotubular networks of the endoplasmic reticulum *in vivo*,^{46,47} which revealed that these shapes resemble unduloids⁴⁸ as depicted in Fig 13. Within the morphology diagram in the (\bar{m}, \bar{v}) -plane, long unduloids with mean curvature $M = m$ and zero bending energy are located between two hyperbolae with $\bar{v} = \bar{v}_1(\bar{m}) \equiv 1/\bar{m}$ and $\bar{v} = \bar{v}_2(\bar{m}) \equiv 3/(4\bar{m})$. Such unduloid shapes will form *via* the fusion of proteoliposomes when some membrane proteins get lost during the fusion process, thereby reducing the spontaneous curvature \bar{m} , or when the fusion process is leaky and the interior volume of the proteoliposomes is reduced during the process.

If one of the membrane necks as indicated by the blue arrow heads in Fig 13 became closed for an extended period of time, the neck can undergo fission, thereby breaking the nanotubes up into several fragments. For reticular networks, such a fragmentation of the nanotubes has indeed been induced experimentally, both *in vivo* and *in vitro*, by downregulating the biosynthesis of GTP-hydrolyzing membrane proteins⁴⁹ and by interrupting the supply of GTP⁴⁴. As a consequence, any mechanism that keeps




Fig. 13 Time-lapse snapshots of an irregular pentagon formed by membrane nanotubes and five three-way junctions as observed by structured illumination microscopy.⁴⁶ Time points in seconds (upper left corners); scale bar in (c) also applies to (a) and (b). The pentagon consists of five three-way junctions connected by five tubular segments. One tube segment is marked by the white dotted rectangle in (b). The four blue arrow heads point to transient membrane necks (or constrictions) formed along three tubular segments, which resemble peristaltic shape deformations of these tubules.⁴⁸



the membrane necks open also prevents nanotube fragmentation.

Peristaltic modes of cylindrical nanotubes. The unduloidal tubes displayed in Fig 13 can be viewed as peristaltic modes of cylindrical nanotubes. Thus, consider a cylindrical tube of length L and radius R_{cy} . The shape fluctuations of such a cylindrical tube can be decomposed into Fourier modes. We will focus on peristaltic modes which preserve the axisymmetry of the cylinder and include the most unstable mode. It is convenient to introduce a fluctuation tension Σ_{fl} that governs the excess area stored in these shape fluctuations. An analogous tension term has been considered for planar membrane segments⁵⁰ and for quasi-spherical vesicles^{51,52}.

For a cylindrical tube with radius R_{cy} and length L , the most unstable mode, denoted by l_{n*} , is characterized by wavenumber $p = p_* = 1/R_{cy}$, period $2\pi R_{cy}$, and mean-squared amplitude⁴⁸

$$\langle |l_{n*}|^2 \rangle = \frac{k_B T}{2\pi R_{cy} L \Sigma_{fl} p_*^2} = \frac{k_B T}{2\pi (L/R_{cy}) \Sigma_{fl}}, \quad (34)$$

which grows as $1/\Sigma_{fl}$ for small Σ_{fl} . Thus, the tube should become fragmented for sufficiently small tensions, for which the amplitude of the most unstable mode exceeds the tube radius, that is, for

$$\langle |l_{n*}|^2 \rangle \geq R_{cy}^2 \quad \text{or} \quad \Sigma_{fl} \leq \Sigma_{th,1} \equiv \frac{k_B T}{2\pi R_{cy} L}. \quad (35)$$

The tension threshold $\Sigma_{th,1}$ is very low. As an example, consider the tube segment within the dotted white rectangle in Fig 13b, which has a length of $L = 2.5 \mu\text{m}$ and a radius of $R_{cy} = 100 \text{nm}$. For this tube, the tension $\Sigma_{th,1}$ has the value $4 \times 10^{-6} \text{mN/m}$ at room temperature.

In Fig 13, the peristaltic modes do not lead to tube fragmentation. Inspection of this figure indicates that the most unstable modes have amplitudes of about half the cylinder radius which defines another tension threshold $\Sigma_{th,2}$ via

$$\langle |l_{n*}|^2 \rangle = (R_{cy}/2)^2 \quad \text{or} \quad \Sigma_{th,2} \equiv \frac{4k_B T}{2\pi R_{cy} L} = 4\Sigma_{th,1}. \quad (36)$$

Thus, the fragmentation of the membrane nanotubes can be avoided by increasing the fluctuation tension Σ_{fl} from $\Sigma_{th,1}$ to $\Sigma_{th,2} = 4\Sigma_{th,1}$.

6 Membranes in contact with condensate droplets

The term ‘condensate droplet’ implies that the droplet is enclosed by a liquid–liquid rather than by a liquid–gas interface. Aqueous two-phase (or biphasic) systems based on biopolymers such as PEG and dextran have been applied for several decades in biochemical analysis and biotechnology⁵³ and are intimately related to water-in-water emulsions⁵⁴. The aqueous phase separation leads to the formation of two coexisting liquid phases, denoted here by α and β as in Fig 14. The two liquid phases α and β are separated by a liquid–liquid interface with interfacial tension $\Sigma_{\alpha\beta}$, which represents a new tension scale for the vesicle-droplet systems.

The aqueous phase separation of PEG-dextran solutions provides an example for *segregative* phase separation, in which one phase is enriched in one macromolecular component such as PEG




Fig. 14 Partial wetting geometry of vesicle-droplet systems, which involve three liquid phases α (white), β (green), and γ (light red). The phases α and β represent two coexisting phases that arise via segregative or associative liquid–liquid phase separation; the γ phase is an inert spectator phase: (a) Phase separation of the exterior solution and adhesion of an exterior β droplet to the outer leaflet of the vesicle membrane; and (b) Phase separation of the interior solution creating one interior α and one interior β droplet, both of which are in contact with the inner leaflet of the membrane.⁶⁷

whereas the other phase is enriched in the other macromolecular component such as dextran. The segregative behavior implies that the different species of macromolecules effectively repel each other. The phase diagram and the interfacial tension of aqueous PEG-dextran solutions is described in Sect S9[†] and Fig S14[†]. Another type of aqueous two-phase system is created by *associative* phase separation, for which one phase is enriched in the macromolecular components whereas the other phase represents a dilute solution of the macromolecules^{55–58}. The associative behavior implies that the different macromolecular species effectively attract each other. Associative phase separation is observed, for instance, in solutions of two, oppositely charged polyelectrolytes^{57,58},

Condensate droplets have also been observed in living cells where they provide separate liquid compartments or organelles which are not enclosed by intracellular membranes. These biomolecular condensates are believed to form via liquid–liquid phase separation in the cytoplasm^{59,60} and can be reconstituted in vitro^{61–64}. They are enriched in certain types of proteins that have intrinsically disordered domains and interact via multivalent macromolecular interactions^{60,63–66}.

6.1 Geometry of vesicle-droplet systems

When a condensate droplet comes into contact with a vesicle membrane, attractive interactions between droplet and membrane lead to the adhesion of the droplet to the membrane. The geometry of these vesicle-droplet systems involves three liquid phases α , β , and γ as shown in Fig 14. The two phases α and β are formed by segregative or associative liquid–liquid phase separation and are separated by the $\alpha\beta$ interface. When the droplet partially wets the membrane, the $\alpha\beta$ interface forms a contact line with the membrane, which divides the membrane up into two segments, the $\alpha\gamma$ segment exposed to the α and γ phases






Fig. 15 Complete engulfment of condensate droplets by GUV membranes (red): (a,b) Two-sphere shape of the membrane, which completely engulfs the dextran-rich β droplet (green) via an in-bud;⁷⁰ (c,d) Two-sphere shape of the membrane, which forms two spherical segments around both the dextran-rich β (green) and the PEG-rich α (black) droplet, connected by a closed contact line neck.⁷¹ In both cases, the systems's free energy is strongly reduced by eliminating the $\alpha\beta$ interface.




Fig. 16 Spontaneous tubulation of the $\alpha\gamma$ membrane segment in contact with the PEG-rich phase α leads to different patterns of nanotubes:⁷² (a) Disordered pattern corresponding to a vesicle membrane that is completely wetted by the PEG-rich phase; and (b) Layer of densely packed tubes corresponding to a membrane that is partially (de)wettered by both aqueous phases. All tubes are connected to the outer vesicle membranes (big circles). In both images, the diameter of the fluorescently labeled tubes is below the diffraction limit of the light microscope.

as well as the $\beta\gamma$ segment in contact with the β and γ phases. In Fig 14a and 14b, the coexisting phases α and β are located outside and inside the vesicle, respectively. Therefore, Fig 14a displays one exterior β droplet at the outer leaflet of the bilayer membrane whereas Fig 14b shows one interior α and one interior β droplet in contact with the inner leaflet of the membrane.

6.2 Remodeling of GUV membranes by condensate droplets

The adhesion of condensate droplets to vesicles generates strong morphological responses of the vesicle membranes. One example for this response is provided by the apparent kinks, which are observed in the optical microscope along the contact line between the adhering droplet and the vesicle membrane, see Fig 14. On nanoscopic scales, these kinks represent membrane segments with a very high curvature, caused by the capillary forces that the $\alpha\beta$ interface exerts onto the membrane at the contact line.^{68,69} For PEG-dextran systems, particularly interesting processes are the complete engulfment of the droplets by the membranes^{70,71} and the formation of membrane nanotubes⁷² as shown in Figs 15 and 16, respectively. The different patterns of nanotubes displayed in Fig 16a,b reveal complete and partial wetting of the GUV membrane by the aqueous two-phase system. More details on this wetting behavior are provided in Sect S9[†] and Fig S15[†].

6.3 Interfacial tension versus curvature-elastic tensions

The two morphological pathways displayed in Figs 15 and 16 arise from the competition between the interfacial tension of the liquid-liquid interface with the curvature-elastic membrane tensions of the two membrane segments $\alpha\gamma$ and $\beta\gamma$.

When the area $A_{\alpha\beta}$ of the $\alpha\beta$ interface is reduced by $\Delta A_{\alpha\beta}$, the interfacial free energy of the vesicle-droplet system decreases by

$$\Delta E_{\text{int}} = -\Sigma_{\alpha\beta} \Delta A_{\alpha\beta} \quad (37)$$

which is proportional to the interfacial tension $\Sigma_{\alpha\beta}$. The latter tension is a material parameter, which is determined by the molecular composition of the α and β phases but is independent of the size and shape of the interface. When the β droplet is completely engulfed by the membrane as in Figs 15, the $\alpha\beta$ interface is completely eliminated from the vesicle-droplet system, thereby reducing its free energy by $\Sigma_{\alpha\beta} A_{\alpha\beta}$ with the total interfacial area $A_{\alpha\beta}$. To better visualize the shape of the vesicle-droplet systems for completely engulfed droplets, the optical images in panels a and c of Fig 15 are compared to schematic drawings in panels b and d of the same figure.

For GUVs filled with PEG-dextran solutions, inward-pointing nanotubes are formed by the membrane segment $\alpha\gamma$ exposed to the PEG-rich phase α , as displayed in Fig 16. The associated spontaneous curvature $m_{\alpha\gamma}$ is generated by the adsorption of PEG molecules onto the membranes.⁷² The free energy reduction by the nanotube formation competes with the free energy reduction by the complete engulfment of the droplets. Which process is energetically more favorable depends on three parameters, the interfacial tension $\Sigma_{\alpha\beta}$ and the curvature-elastic tensions

$$\sigma_{\alpha\gamma} \equiv 2\kappa_{\alpha\gamma} m_{\alpha\gamma}^2 \quad \text{and} \quad \sigma_{\beta\gamma} \equiv 2\kappa_{\beta\gamma} m_{\beta\gamma}^2 \quad (38)$$

of the two membrane segments.⁶⁹

When the $\alpha\gamma$ membrane segment has a large spontaneous curvature $m_{\alpha\gamma} \gg 1/R_{\text{ve}}$ but a small mean curvature, this segment has a large bending energy of the order of $\sigma_{\alpha\gamma} A_{\alpha\gamma}$ which is proportional to the curvature-elastic tension $\sigma_{\alpha\gamma}$ of the $\alpha\gamma$ segment and to the surface area $A_{\alpha\gamma}$ of this segment. Now, when the membrane area ΔA_{me} of the weakly curved mother vesicle is transferred to form a nanotube of mean curvature $M \simeq m_{\alpha\gamma}$, we gain the bending energy

$$\Delta E_{\text{be}} = -\sigma_{\alpha\gamma} \Delta A_{\text{me}} = -2\kappa_{\alpha\gamma} m_{\alpha\gamma}^2 \Delta A_{\text{me}}. \quad (39)$$

We now repeat this argument for the membrane segment $\beta\gamma$ in contact with the dextran-rich phase β , which generates the spontaneous curvature $m_{\beta\gamma}$ of the $\beta\gamma$ segment. When the membrane area ΔA_{me} of this segment forms a nanotube with mean curvature $M \simeq m_{\beta\gamma}$, we gain the bending energy

$$\Delta E_{\text{be}} = -\sigma_{\beta\gamma} \Delta A_{\text{me}} = -2\kappa_{\beta\gamma} m_{\beta\gamma}^2 \Delta A_{\text{me}}. \quad (40)$$

Because the membrane segment $\beta\gamma$ in contact with the dextran-rich phase is not observed to form nanotubes, we can directly conclude from Eqs (39) and (40) that the curvature-elastic tension $\sigma_{\beta\gamma}$ of the $\beta\gamma$ segment is smaller than the curvature-elastic tension $\sigma_{\alpha\gamma}$ of the $\alpha\gamma$ segment.



The reduction of the membrane's bending energy by nanotube formation as given by Eq (39) remains to be compared with the reduction of the interfacial free energy by the engulfment of the droplet as described by Eq (37). We then conclude that complete droplet engulfment should be observed for⁶⁹

$$\Sigma_{\alpha\beta} \gg \sigma_{\alpha\gamma} = 2\kappa_{\alpha\gamma}m_{\alpha\gamma}^2 \quad (\text{complete engulfment}). \quad (41)$$

that is, when the interfacial tension $\Sigma_{\alpha\beta}$ is large compared to the curvature-elastic tension $\sigma_{\alpha\gamma}$ of the membrane segment $\alpha\gamma$ exposed to the PEG-rich α phase. On the other hand, the $\alpha\gamma$ membrane segment will form nanotubes if

$$\sigma_{\alpha\gamma} \gg \Sigma_{\alpha\beta} \quad (\text{nanotube formation}). \quad (42)$$

that is, if the curvature-elastic tension $\sigma_{\alpha\gamma}$ of the $\alpha\gamma$ segment is large compared to the interfacial tension $\Sigma_{\alpha\beta}$. For the PEG-dextran two-phase system, the interfacial tension $\Sigma_{\alpha\beta}$ varies by several orders of magnitude (Fig S14b[†]). As a consequence, this two-phase system can lead to both $\Sigma_{\alpha\beta} \gg \sigma_{\alpha\gamma}$ and $\sigma_{\alpha\gamma} \gg \Sigma_{\alpha\beta}$, that is, to complete engulfment of the condensate droplet and to the formation of membrane nanotubes as in Figs 15 and 16.

7 Conclusions

In this paper, we elucidate the notion of membrane tension for three biomimetic membrane systems as provided by planar lipid bilayers, unilamellar nanovesicles, and GUVs. For planar bilayers and nanovesicles, we distinguish the bilayer tension Σ from the two leaflet tensions, Σ_1 and Σ_2 , with $\Sigma = \Sigma_1 + \Sigma_2$. A global view of the leaflet tensions is provided by the two-dimensional parameter spaces in Figs 3 and 5 for planar and vesicle bilayers, respectively. Particularly interesting bilayer states are tensionless but have opposite leaflet tensions (OLTs) as in Fig 4 for planar bilayers and in Fig 6 for nanovesicles. Outside their stability regimes, planar and vesicle bilayers exhibit lipid flip-flops (Figs S5[†] and S6[†]) and structural instabilities (Figs S9[†] and S10[†]).

For GUV membranes, the total membrane tension Σ_{tot} is equal to the mechanical membrane tension Σ plus the curvature-elastic tension $2\kappa m^2$. The mechanical membrane tension depends on the size and shape of the membrane as demonstrated for multispheres (Fig 10) in Sect 5.1 and for positive two-sphere shapes (Fig 11) in Sect 5.2. We then addressed the different shapes of tubular membranes, which can be multispherical (Fig 12), unduloidal (Fig 13), or cylindrical. Unduloidal shapes represent peristaltic deformations of cylindrical shapes. The fluctuation tension Σ_{fl} for the excess membrane stored in the peristaltic modes exhibits two tension thresholds as given by Eqs (35) and (36).

The adhesion of condensate droplets to GUVs is considered in Sect 6. The geometry of the vesicle-droplet systems (Fig 14) involves three surfaces, the liquid-liquid interface with interfacial tension $\Sigma_{\alpha\beta}$ as well as two membrane segments $\alpha\gamma$ and $\beta\gamma$ with curvature-elastic tensions $\sigma_{\alpha\gamma}$ and $\sigma_{\beta\gamma}$ as defined in Eq (38). The relative magnitude of these tensions determines the remodeling of the GUV membranes: For large interfacial tension, the condensate droplets are completely engulfed by the vesicle membranes as in Fig 15 whereas one of the membrane segments forms membrane nanotubes as in Fig 16 for small interfacial tension.

Finally, it follows from the finite persistence length of tensionless fluid membranes as given by Eq (6) and from the poration of sufficiently large membranes in Sect 3.3 that the thermodynamic route to membrane tension is ill-defined, in contrast to the thermodynamic definition of interfacial tension by J. Willard Gibbs.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

I thank Bartosz Różycki, Udo Seifert, Gerhard Gompper, and Alberto Imperato for useful correspondence, and all my collaborators for enjoyable and fruitful interactions. This research was conducted within the Max Planck School "Matter to Life" supported by the German Federal Ministry of Education and Research (BMBF) in collaboration with the Max Planck Society and the Max Planck Institute of Colloids and Interfaces.

References

- 1 R. Lipowsky, R. Ghosh, V. Satarifard, A. Sreekumari, M. Zamaletdinov, B. Różycki, M. Miettinen and A. Grafmüller, *Biomolecules*, 2023, **13**, 926.
- 2 R. Lipowsky, *Faraday Discuss.*, 2013, **161**, 305–331.
- 3 R. Lipowsky, *Adv. Biology*, 2022, **6**, 2101020.
- 4 M. D. Collins and S. L. Keller, *PNAS*, 2008, **105**, 124 – 128.
- 5 A. Orth, L. Johannes, W. Römer and C. Steinem, *Chem. Phys. Chem.*, 2012, **13**, 108 – 114.
- 6 B. Różycki and R. Lipowsky, *J. Chem. Phys.*, 2015, **142**, 054101.
- 7 L. D. Landau and E. M. Lifshitz, *Theory of Elasticity*, Pergamon Press, New York, 1986.
- 8 E. Evans and R. Hochmuth, *Current Topics in Membranes and Transport*, Elsevier, 1978, vol. 10, pp. 1–64.
- 9 R. Lipowsky, *Adv. Colloid Interface Sci.*, 2014, **208**, 14–24.
- 10 T. Bhatia, J. Agudo-Canalejo, R. Dimova and R. Lipowsky, *ACS Nano*, 2018, **12**, 4478–4485.
- 11 P.-G. De Gennes and C. Taupin, *J. Phys. Chem.*, 1982, **86**, 2294–2304.
- 12 W. Helfrich and W. Harbich, *Chemica Scripta*, 1985, **25**, 32–36.
- 13 L. Peliti and S. Leibler, *Phys. Rev. Lett.*, 1985, **54**, 1690–1693.
- 14 R. Goetz, G. Gompper and R. Lipowsky, *Phys. Rev. Lett.*, 1999, **82**, 221–224.
- 15 J. Rowlinson and B. Widom, *Molecular Theory of Capillarity*, Oxford University Press, Oxford, 1989.
- 16 U. Seifert and R. Lipowsky, *Phys. Rev. A*, 1990, **42**, 4768–4771.
- 17 G. Gompper and D. Kroll, *Phys. Rev. E*, 1995, **51**, 514–525.
- 18 M. Zamaletdinov, M. Miettinen and R. Lipowsky, *Soft Matter*, 2023, **19**, 6929–6944.
- 19 A. Sreekumari and R. Lipowsky, *Soft Matter*, 2022, **18**, 6066–6078.
- 20 A. Hossein and M. Deserno, *Biophys. J.*, 2020, **118**, 624–642.
- 21 M. Varma and M. Deserno, *Biophys. J.*, 2022, **121**, 4001 –



- 4018.
- 22 O. Farago, *J. Chem. Phys.*, 2003, **119**, 596–605.
- 23 T. V. Tolpekina, W. K. den Otter and W. J. Briels, *J. Chem. Phys.*, 2004, **121**, 8014 – 8020.
- 24 M. Miettinen and R. Lipowsky, *Nano Lett.*, 2019, **19**, 5011–5016.
- 25 A. Imparato, J. Shillcock and R. Lipowsky, *Europhys. Lett.*, 2005, **69**, 650–656.
- 26 A. Imparato, *J. Chem. Phys.*, 2006, **124**, 154714.
- 27 J. Stecki, *J. Chem. Phys.*, 2006, **125**, 154902.
- 28 J.-B. Fournier and C. Barbetta, *Phys. Rev. Lett.*, 2008, **100**, 078103.
- 29 H. Shiba, H. Noguchi and J.-B. Fournier, *Soft Matter*, 2016, **12**, 2373–2380.
- 30 R. Ghosh, V. Satarifard, A. Grafmüller and R. Lipowsky, *Nano Lett.*, 2019, **19**, 7703–7711.
- 31 J. Z. A. Finkelstein and F. Cohen, *Anu. Rev. Physiol.*, 1986, **48**, 163–174.
- 32 J. Shillcock and R. Lipowsky, *Nature Materials*, 2005, **4**, 225–228.
- 33 A. Grafmüller, J. Shillcock and R. Lipowsky, *Phys. Rev. Lett.*, 2007, **98**, 218101.
- 34 L. Gao, R. Lipowsky and J. C. Shillcock, *Soft Matter*, 2008, **4**, 1208–1214.
- 35 T.-T. Kliesch, J. Dietz, L. Turco, P. Halder, E. Polo, M. Tarrantola, R. Jahn and A. Janshoff, *Sci. Rep.*, 2017, **7**, 12070.
- 36 R. B. Lira, T. Robinson, R. Dimova and K. A. Riske, *Biophys. J.*, 2019, **116**, 79 – 91.
- 37 I. Ivanov, R. B. Lira, T.-Y. D. Tang, T. Franzmann, A. Klosin, L. C. da Silva, A. A. Hyman, K. Landfester, R. Lipowsky, K. Sundmacher and R. Dimova, *Advanced Biosystems*, 2019, **3**, 1800314.
- 38 S. Deshpande, S. Wunna, D. Hueting and C. Dekker, *Small*, 2019, **15**, 1902898.
- 39 A. Grafmüller, J. C. Shillcock and R. Lipowsky, *Biophys. J.*, 2009, **96**, 2658–2675.
- 40 J. Steinkühler, R. L. Knorr, Z. Zhao, T. Bhatia, S. Bartelt, S. Wegner, R. Dimova and R. Lipowsky, *Nature Commun.*, 2020, **11**, 905.
- 41 T. Bhatia, S. Christ, J. Steinkühler, R. Dimova and R. Lipowsky, *Soft Matter*, 2020, **16**, 1246–1258.
- 42 R. Lipowsky, *Adv. Colloid Interface Sci.*, 2022, **301**, 102613.
- 43 R. Lipowsky, *The Giant Vesicle Book*, Taylor & Francis, 2019, ch. 5, pp. 73–168.
- 44 R. E. Powers, S. Wang, T. Y. Liu and T. A. Rapoport, *Nature*, 2017, **543**, 257–272.
- 45 M. A. Betancourt-Solis, T. Desai and J. A. McNew, *J. Biol. Chem.*, 2018, **293**, 18514–18524.
- 46 D. Holcman, P. Parutto, J. E. Chambers, M. Fantham, L. J. Young, S. J. Marciniak, C. F. Kaminski, D. Ron and E. Avezov, *Nat. Cell Biol.*, 2018, **20**, 1118–1125.
- 47 C. J. Obara, A. S. Moore and J. Lippincott-Schwartz, *Cold Spring Harb. Perspect. Biol.*, 2023, **15**, a041259.
- 48 R. Lipowsky, S. Pramanik, A. S. Benk, M. Tarnawski, J. P. Spatz and R. Dimova, *ACS Nano*, 2023, **17**, 11957–11968.
- 49 G. Orso, D. Pendin, S. Liu, J. Toso, T. J. Moss, J. E. Faust, M. Micaroni, A. Egorova, A. Martinuzzi, J. A. McNew and A. Daga, *Nature*, 2009, **460**, 978–983.
- 50 W. Helfrich and R.-M. Servuss, *Il Nuovo Cimento*, 1984, **3D**, 137–151.
- 51 S. Milner and S. Safran, *Phys. Rev. A*, 1987, **36**, 4371–4379.
- 52 U. Seifert, *Z. Physik B*, 1995, **97**, 299–309.
- 53 P. A. Albertsson, *Partition of Cell Particles and Macromolecules: Separation and Purification of Biomolecules, Cell Organelles Membranes, and Cells in Aqueous Polymer Two-Phase Systems and Their Use in Biochemical Analysis and Biotechnology*, Wiley, 3rd edn, 1986.
- 54 J. Esquena, *Curr. Opin. Colloid Interface Sci.*, 2016, **25**, 109–119.
- 55 C. D. Crowe and C. D. Keating, *Interface Focus*, 2018, **8**, 20180032.
- 56 H. G. B. de Jong and H. R. Kruyt, *Proc. K. Ned. Akad. Wet.*, 1929, **32**, 849 – 856.
- 57 C. E. Sing, *Adv. Colloid Interface Sci.*, 2017, **239**, 2 – 16.
- 58 L. Li, S. Srivastava, M. Andreev, A. B. Marciel, J. J. de Pablo and M. V. Tirrell, *Macromolecules*, 2018, **51**, 2988 – 2995.
- 59 C. P. Brangwynne, C. R. Eckmann, D. S. Courson, A. Rybarska, C. Hoeghe, J. Gharakhani, F. Jülicher and A. A. Hyman, *Science*, 2009, **324**, 1729 – 1732.
- 60 S. F. Banani, H. O. Lee, A. A. Hyman and M. K. Rosen, *Nature Rev.: Mol. Cell Biol.*, 2017, **18**, 285–295.
- 61 P. Li, S. Banjade, H.-C. Cheng, S. Kim, B. Chen, L. Guo, M. Llaguno, J. V. Hollingsworth, D. S. King, S. F. Banani, P. S. Russo, Q.-X. Jiang, B. T. Nixon and M. K. Rosen, *Nature*, 2012, **483**, 336 – 341.
- 62 A. Patel, H. O. Lee, L. Jawerth, S. Maharana, M. Jahnel, M. Y. Hein, S. Stoynov, J. Mahamid, S. Saha, T. M. Franzmann, S. Grill, D. Drechsel, A. A. Hyman and S. Alberti, *Cell*, 2015, **162**, 1066 – 1077.
- 63 Y. Lin, D. S. W. Protter, M. K. Rosen and R. Parker, *Mol. Cell*, 2015, **60**, 208 – 219.
- 64 A. Molliex, J. Temirov, J. Lee, H. J. Kim, T. Mittag and J. P. Taylor, *Cell*, 2015, **163**, 123 – 133.
- 65 T. W. Han, M. Kato, S. Xie, L. C. Wu, H. Mirzaei, J. Pei, M. Chen, Y. Xie, J. Allen, G. Xiao and S. L. McKnight, *Cell*, 2012, **149**, 768 – 779.
- 66 L. Guo and J. Shorter, *Mol. Cell*, 2015, **60**, 189 – 192.
- 67 R. Lipowsky, *Membranes*, 2023, **13**, 223.
- 68 H. Kusumaatmaja, Y. Li, R. Dimova and R. Lipowsky, *Phys. Rev. Lett.*, 2009, **103**, 238103.
- 69 R. Lipowsky, *J. Phys. Chem. B*, 2018, **122**, 3572–3586.
- 70 Y. Li, H. Kusumaatmaja, R. Lipowsky and R. Dimova, *J. Phys. Chem. B*, 2012, **116**, 1819–1823.
- 71 R. Dimova and R. Lipowsky, *Adv. Mater. Interf.*, 2017, **4**, 1600451.
- 72 Y. Liu, J. Agudo-Canalejo, A. Grafmüller, R. Dimova and R. Lipowsky, *ACS Nano*, 2016, **10**, 463–474.



Title: The many faces of membrane tension for biomembranes and vesicles

Author: Reinhard Lipowsky

[View Article Online](#)

DOI: 10.1039/D4FD00184B

Data Availability Statement

This paper provides a theoretical analysis of simulation data and experimental data that are available in the published and cited literature.

