

CORRECTION

[View Article Online](#)
[View Journal](#) | [View Issue](#)


Cite this: *Green Chem.*, 2024, **26**, 8878

Correction: Metal-free visible-light-induced phosphorylation of unactivated alkyl iodides with white phosphorus as the P-atom source

Fushan Chen, Jialiang Peng, Yue Ying, Yinwei Cao, Pengxiang Xu,* Guo Tang* and Yufen Zhao

DOI: 10.1039/d4gc90072c


rsc.li/greenchem

Correction for 'Metal-free visible-light-induced phosphorylation of unactivated alkyl iodides with white phosphorus as the P-atom source' by Fushan Chen *et al.*, *Green Chem.*, 2023, **25**, 6629–6634, <https://doi.org/10.1039/D3GC01579C>.

The authors regret that the structures for compounds **4n** and **4o** in Scheme 2 of their original publication were incorrect. The correct version of Scheme 2 is shown below and the original ESI is replaced by a revised version. Additional NMR spectra have also been included in the revised ESI.

Department of Chemistry, College of Chemistry and Chemical Engineering, and the Key Laboratory for Chemical Biology of Fujian Province, Xiamen University, Xiamen, Fujian 361005, China. E-mail: t12g21@xmu.edu.cn, xpengxiang@xmu.edu.cn





Scheme 2 Scope of secondary phosphine oxides. Reaction conditions from Table 1, entry 1; after irradiation by blue LEDs for 12 h, the reaction mixtures were stirred at room temperature for 1 h in air (oxidation of R_2PH by air afforded DAPOs).

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

