

## PAPER

View Article Online  
View Journal | View Issue



Cite this: *Org. Biomol. Chem.*, 2024, **22**, 1915

Received 29th November 2023,  
Accepted 2nd February 2024

DOI: 10.1039/d3ob01943h

rsc.li/obc

# TiF<sub>4</sub>-catalyzed direct amidation of carboxylic acids and amino acids with amines†


Abdulkhaliq A. Alawaed and P. Veeraraghavan Ramachandran \*

Unlike other metal fluorides, catalytic titanium tetrafluoride enhances the direct amidation of aromatic and aliphatic carboxylic acids and *N*-protected amino acids in refluxing toluene. While aromatic acids were converted to amides with 10 mol% of the catalyst within 24 h, aliphatic acids underwent a faster reaction (12 h), with lower catalyst loading (5 mol%). This protocol is equally efficient with alkyl and aryl amines providing a variety of carboxamides and peptides in 60–99% yields.

## Introduction

The prevalence of amides in organic molecules and biomolecules has made their preparation, structure, and reactivity a much-discussed subject. The direct amidation of carboxylic acids is the most straightforward protocol to access them, although it is a challenging reaction<sup>1</sup> due to the formation of ammonium salt. Dehydration of the salt to an amide can be achieved at high temperatures. For milder reactions, the conversion of the acid to a derivative capable of undergoing facile nucleophilic acyl substitution, such as an acid chloride or anhydride, is opted for. Also, a variety of carbodiimide reagents, such as dicyclohexylcarbodiimide (DCC),<sup>2</sup> 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC),<sup>3</sup> and 1,1'-carbonyldiimidazole (CDI)<sup>4</sup> or phosphate agents, such as benzotriazol-1-yl-oxytripyrrolidinophosphonium hexafluorophosphate (PyBOP),<sup>5</sup> Castro's reagent (BOP),<sup>6</sup> T3P,<sup>7</sup> HBTU,<sup>8</sup> HATU,<sup>9</sup> *etc.* have been developed for the dehydrative coupling of an acid with an amine.<sup>10,11</sup> However, in addition to the reagent cost, the need for additional equivalents of amines and excess reagents adds to the difficulties in the separation of the amides from stoichiometric quantities of the side products. Leading the intense activity in developing alternate catalyzed direct amidation protocols<sup>12,13</sup> are boron-<sup>14</sup> and silicon-based<sup>15</sup> catalysts as well as transition metal salts, particularly those of Ti,<sup>16–18</sup> Mn,<sup>19</sup> Fe,<sup>20</sup> Zn,<sup>21,22</sup> Zr,<sup>17,23,24</sup> In,<sup>25</sup> Hf,<sup>26</sup> Ta,<sup>27,28</sup> *etc.* Indeed, some of these reactions are carried out in sealed tubes and are effective only for aliphatic acids and several of these protocols are inefficient for amidation with anilines.

We have recently reported the effect of catalytic titanium tetrachloride (TiCl<sub>4</sub>) for the borane-amine mediated reduction of ketones,<sup>29</sup> acids,<sup>30</sup> amides,<sup>31</sup> and nitriles,<sup>32</sup> deoxygenation of esters to ethers<sup>33</sup> and deoxyhalogenation of carbonyl compounds.<sup>34</sup> During these projects, we undertook a systematic evaluation of the catalytic ability of several metal halides and established that the reaction depended on the Lewis acidity of these halides and TiCl<sub>4</sub> is a critical and superior catalyst. Interested in TiCl<sub>4</sub>'s applicability for amidation, a literature search was undertaken, which led to a recent, impractical process with 300 mol% of TiCl<sub>4</sub> in pyridine as the solvent (Scheme 1(i)).<sup>18</sup> A prior report comparing several Lewis acids



**Scheme 1** Comparison of titanium halide mediated amidation catalyzed amidation.

Department of Chemistry, Purdue University, West Lafayette, Indiana 47907, USA.

E-mail: chandran@purdue.edu

†Electronic supplementary information (ESI) available: NMR spectra of products. See DOI: <https://doi.org/10.1039/d3ob01943h>



for amidation reported that  $\text{ZrCl}_4$  is superior to  $\text{TiCl}_4$  (Scheme 1(ii)).<sup>17</sup> The first report from half a century ago using large excess of amines is inconclusive (Scheme 1(iii)).<sup>16</sup>

## Results and discussion

After attempting an amidation of benzoic acid (**1a**) with benzylamine (**2a**) without any catalyst in refluxing toluene (24 h, 15% yield; Table 1, entry 1), a similar reaction in the presence of 10 mol%  $\text{TiCl}_4$  provided 89% yield of the corresponding amide **3aa**<sup>35</sup> (Table 1, entry 2).  $\text{TiBr}_4$  was equally effective (91% yield, entry 3). Other group 4 metal chlorides, such as  $\text{ZrCl}_4$  and  $\text{HfCl}_4$ , were poor amidation catalysts (entries 4 and 5) under similar conditions. Mader and Helquist have reported that titanium isopropoxide (TIPO) is a poor catalyst for intermolecular amidation.<sup>36</sup> However, Adolfsson and coworkers reported successful amidation in THF in a sealed tube with 1.5 equiv. of the acid and catalytic TIPO or  $\text{Ti}(n\text{-BuO})_4$ .<sup>37</sup> Attempts to scale-up the reaction with  $\text{TiCl}_4$  and  $\text{TiBr}_4$  were unrealistic due to the corrosive nature of the reagents, so we turned our attention to  $\text{TiF}_4$ .

We had observed during our earlier-mentioned deoxyhalogenation of carbonyls<sup>34</sup> using  $\text{TiCl}_4$  and  $\text{TiBr}_4$  that the corresponding fluoride did not participate in the nucleophilic transfer of the fluoride. This led us to believe that  $\text{TiF}_4$  might be a stable catalyst for Lewis acid-mediated reactions. Surprisingly, there are only scarce reports on direct amidation using metal fluorides as catalysts. Metal fluoride salts, such as  $\text{ZnF}_2$  and


$\text{AlF}_3$ , are ineffective compared to chlorides for the formylation of amines.<sup>21</sup> Also, a recent report on the screening of different fluoride salts such as  $\text{CsF}$ ,  $\text{NaF}$ ,  $\text{KF}$ ,  $\text{AgSbF}_6$ , and  $\text{KPF}_6$  as promoters for a similar esterification produced poor results with all of them, except  $\text{KPF}_6$ .<sup>38</sup> The hexafluorophosphate was effective for amidation as well.<sup>38</sup> Our long-time interest in the chemistry of fluorinated molecules stimulated us to examine  $\text{TiF}_4$  for the direct amidation of carboxylic acids. The literature suggested that unlike the chloride analog,  $\text{TiF}_4$  is easier to handle, stable in air and water,<sup>39</sup> and has excellent solubility in dimethoxyethane and THF.<sup>40,41</sup> Our concerns were the possible complexation of  $\text{TiF}_4$  with amines,<sup>42</sup> which could be detrimental to their nucleophilicity. The results documented herein unequivocally confirm that 5–10% of  $\text{TiF}_4$  is a catalytic promoter for the direct amidation of both aliphatic and aromatic carboxylic acids with both classes of amines (Scheme 1(iv)).

A reaction of an equivalent each of **1a** with **2a** in refluxing toluene in the presence of 10 mol%  $\text{TiF}_4$  for 24 h provided 77% of **3aa**<sup>35</sup> (entry 6), which was increased to 96% with an additional 10% of the acid (entry 7). A large scale 50 mmol reaction was carried out for the synthesis of *N*-benzylbenzamide (**3aa**), which resulted in a slightly reduced yield of 77%. We then explored the limitations of the process by varying the catalyst loading, solvent, reaction temperature, etc. (entries 8–14) and concluded that 10% catalyst with 1.1 equiv. of the acid provided the optimal yield.  $\text{BF}_3\cdot\text{Et}_2\text{O}$ ,<sup>43</sup>  $\text{ZnF}_2$  and,  $\text{ZrF}_4$ , were also compared (entries 15–17), which gave unsatisfactory results.

Having determined the optimal conditions for the amidation of **1a** with **2a**, the generality of the protocol was examined by varying the acids and amines (Fig. 1 and 2). In all cases, pure amides were isolated after a simple acid–base workup. Initially, we maintained the amine (benzylamine) and focused on the effect of the aromatic acid (Fig. 1). Substitutions on the benzene ring of **1a** were examined with benzoic acid substituents bearing electron-withdrawing groups (EWGs) and electron-donating groups (EDGs) (**1b–1j**). All of the amides were obtained in good to excellent yields. However, moderate yield was realized with *ortho*-substituted benzoic acids **1b** and **1g**, probably due to the increased steric effects. Among those tested, 4-nitrobenzoic acid failed to undergo amidation. Cinnamic acid (**1k**) and nicotinic acid (**1l**) were also treated with **2a** to yield the corresponding amides **3ka** and **3la** in 99% and 93% yields, respectively. Replacing benzylamine with cyclohexylamine (**2c**) resulted in 24% yield of the amide **3ac**. However, increasing the catalyst to 50 mol% afforded 90% yield of the product.

Preparation of tertiary amides from aromatic acids was examined using morpholine (**2d**) as the representative secondary amine. Thus, benzoic acids with EWGs and EDGs were amidated with **2d** in 90–98% yield, except 2-methoxybenzoic acid **1g**, which, as with **2a**, provided **3gd** in 61% yield. Significantly, amidations of benzoic acids and anilines each bearing EDGs and EWGs were complete within 24 h, and the products were isolated in 77–96% yields. It is noteworthy that


Table 1 Optimization of amidation conditions




Entry	LA	Solvent	Yield, <sup>a</sup> %
1	None	Toluene	15
2	$\text{TiCl}_4$	Toluene	89
3	$\text{TiBr}_4$	Toluene	91
4	$\text{ZrCl}_4$	Toluene	45
5	$\text{HfCl}_4$	Toluene	47
6	$\text{TiF}_4$	Toluene	77 <sup>b</sup>
7	$\text{TiF}_4$	Toluene	96
8	$\text{TiF}_4$	Toluene	28 <sup>c</sup>
9	$\text{TiF}_4$	Toluene	NR <sup>d</sup>
10	$\text{TiF}_4$	Xylene	69
11	$\text{TiF}_4$	Benzene	NR <sup>c</sup>
12	$\text{TiF}_4$	Cyclohexane	NR <sup>c</sup>
13	$\text{TiF}_4$	THF	NR <sup>c</sup>
14	$\text{TiF}_4$	$\text{CH}_3\text{CN}$	NR <sup>c</sup>
15	$\text{BF}_3\cdot\text{Et}_2\text{O}$	Toluene	79 (83) <sup>e</sup>
16	$\text{ZnF}_2$	Toluene	43
17	$\text{ZrF}_4$	Toluene	41

<sup>a</sup> Isolated yields for reactions using 1.1 equiv. of **1a** and 1.0 equiv. of **2a**. <sup>b</sup> 1.0 equiv. of carboxylic acid. <sup>c</sup> 5 mol% of  $\text{TiF}_4$  used. <sup>d</sup> Reaction at 80 °C. <sup>e</sup> From ref. 43 in refluxing benzene (100 h) in the presence of  $\text{Et}_3\text{N}$ . NR = No reaction.





**Fig. 1** Amidation of aryl carboxylic acids using 10 mol%  $\text{TiF}_4$ , 1.1 equiv. of carboxylic acid and 1 equiv. of amine. Isolated yields shown. <sup>a</sup>Yield for the 50 mmol scale reaction. <sup>b</sup>Yield with 50 mol% catalyst.



**Fig. 2** Amidation of alkyl carboxylic acids using 5 mol%  $\text{TiF}_4$ , 1.1 equiv. of carboxylic acid and 1 equiv. of amine. Isolated yields shown. <sup>a</sup>Indicates product yield with 20% catalyst loading. <sup>b</sup>Yield with 50 mol% catalyst.


amidations with anilines are seldom reported for several of the procedures described in the literature. Under similar conditions, metoclopramide (**3mh**), an antiemetic and gut motility stimulator,<sup>44</sup> was synthesized from 4-amino-5-chloro-2-methoxybenzoic acid (**1m**) and *N,N*-diethylethane-1,2-diamine (**2h**).

The direct amidation of aliphatic acids (**4a–4r**) required only 5 mol%  $\text{TiF}_4$  and were faster (12 h) (Fig. 2). Formic, acetic, cyclic, linear, and arylacetic acids reacted smoothly and the amides (**5**)<sup>35</sup> were obtained in good to excellent yields (Fig. 2). The amide **5ia** from adamantylcarboxylic acid (**4i**) and benzylamine (**2a**) was obtained in only 47% yield under the standard conditions. However, it improved to 83% with 50 mol% of the catalyst.

The reaction of the anti-inflammatory drug ibuprofen (**4l**) and **2a** produced the corresponding amide **5la** in 98% yield. A decreased yield of 47% was observed for amide **5ma** from mandelic acid (**4m**) under the standard conditions, which was improved to 65% with 50 mol% of the catalyst. The need for excess catalyst could be due to the reaction of the hydroxyl group with the catalyst. No side reaction was observed with substrates having distal double (**4p**) and triple (**4q**) bonds in the carboxylic acids and amides **5pa** and **5qa** were obtained in 86% and 94% yields, respectively. Nevertheless, the yield of the amide **5ji** was low (29%) when phenylacetic acid (**4j**) was

reacted with a secondary amine (**2i**). An optically pure amine, (*R*)-(+)- $\alpha$ -methylbenzylamine (**2j**), was treated with **4j** to examine the stereochemical retention during the amidation. The chiral amide **5jj**, which was isolated in 57% yield, revealed an optical rotation comparable to the pure chiral amide reported in the literature,<sup>45</sup> demonstrating the applicability of this process for chiral substrates.

The new amidation protocol was next examined to prepare peptides from amino acids. The amidation of the sterically hindered *N*-Fmoc and *N*-Boc protected glycine (**6a** and **6b**) with benzylamine **2a** (Fig. 3) yielded the target amides **7aa**<sup>35</sup> and **7ba** in 62% and 88% yields, respectively.



**Fig. 3** Direct amidation of amino acids and primary amines using catalytic  $\text{TiF}_4$  (5%).






Fig. 4 Proposed mechanism of titanium tetrafluoride-catalyzed amidation of carboxylic acids.

More challenging di- and tripeptide couplings were also successfully achieved *via* the combination of *N*-Fmoc-Gly-OH (**6a**) and *N*-Boc-Gly-OH (**6b**), with ethyl  $\alpha$ -amino esters of phenylalanine **2k** under the described conditions when the amides **7ak** and **7bk** were obtained in excellent (99%) yields. Subsequently, dipeptide **7bk** was deprotected to give the intermediate Gly-Phe-OEt, which was further reacted with *N*-Boc-Gly-OH (**6b**) to furnish the tripeptide *N*-Boc-Gly-Gly-Phe-OEt **7ck** in 63% yield, demonstrating the capability of the reaction for sequential amidation reactions.

The  $\text{TiF}_4$ -mediated amidation is believed to proceed *via* the coordination of the catalyst with the acid, which prepared the carbonyl for the nucleophilic attack by the amine, which was followed by a proton transfer, elimination of water and regeneration of the catalyst (Fig. 4). A similar mechanism proposed for the amidation of carboxylic acids with catalytic  $\text{ZrCl}_4$ <sup>46</sup> forms the basis of our hypothesis.  $\text{TiF}_4$  is known to complex with amines.<sup>42</sup> It is known to complex with ketones and carboxylic acids as well.<sup>41</sup> However, the formation of an ammonium salt upon the mixing of the carboxylic acid and amine may preclude the formation of the amine-titanium complex. In refluxing toluene, the ammonium salt is probably cleaved, and the amine attacks the Ti-complexed carbonyl carbon.

## Conclusion

In conclusion, we have described a direct amidation protocol using catalytic  $\text{TiF}_4$  (10 mol% for aromatic or 5 mol% for aliphatic acids) in refluxing toluene. The former is amidated within 24 h, while the latter requires only 12 h. This procedure can provide secondary and tertiary amides from both aliphatic and aromatic acids and amines in very high yields. This is applicable for the synthesis of peptides from amino esters and *N*-protected amino acids and has several advantages, such as: (i) ready availability and (ii) very low molecular weight of the catalyst, (iii) non-chromatographic purification, (iv) no require-

ment of removal of water using azeotropic/molecular sieves for small scale reactions, and (v) consistently high yields. We continue to explore ways to improve and extend this amidation procedure.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

We thank the Hebert C. Brown Center for Borane Research for financial support and Henry J. Hamann of our group for helpful discussions.

## References

- 1 D. G. Brown and J. Bostrom, *J. Med. Chem.*, 2016, **59**, 4443–4458.
- 2 J. C. Sheehan and G. P. Hess, *J. Am. Chem. Soc.*, 1955, **77**, 1067–1068.
- 3 N. Goyal, *Synlett*, 2010, 335–336.
- 4 R. Paul and G. W. Anderson, *J. Am. Chem. Soc.*, 1960, **82**, 4596–4600.
- 5 J. Coste, D. Lenguyen and B. Castro, *Tetrahedron Lett.*, 1990, **31**, 205–208.
- 6 B. Castro, J. R. Dormoy, G. Evin and C. Selve, *Tetrahedron Lett.*, 1975, 1219–1222.
- 7 H. Wissmann and H. J. Kleiner, *Angew. Chem., Int. Ed. Engl.*, 1980, **19**, 133–134.
- 8 V. Dourtoglou, B. Gross, V. Lambropoulou and C. Zioudrou, *Synthesis*, 1984, 572–574.
- 9 L. A. Carpino, *J. Am. Chem. Soc.*, 1993, **115**, 4397–4398.
- 10 E. Valeur and M. Bradley, *Chem. Soc. Rev.*, 2009, **38**, 606–631.
- 11 J. R. Dunetz, J. Magano and G. A. Weisenburger, *Org. Process Res. Dev.*, 2016, **20**, 140–177.
- 12 J. Magano, *Org. Process Res. Dev.*, 2022, **26**, 1562–1689.
- 13 A. S. Santos, A. M. S. Silva and M. M. B. Marques, *Eur. J. Org. Chem.*, 2020, 2501–2516.
- 14 M. T. Sabatini, L. T. Boulton and T. D. Sheppard, *Sci. Adv.*, 2017, **3**, e1701028.
- 15 J. J. Davies, D. C. Braddock and P. D. Lickiss, *Org. Biomol. Chem.*, 2021, **19**, 6746–6760.
- 16 J. D. Wilson and H. Weingart, *Can. J. Chem.*, 1970, **48**, 983.
- 17 C. L. Allen, A. R. Chhatwal and J. M. J. Williams, *Chem. Commun.*, 2012, **48**, 666–668.
- 18 A. Leggio, J. Bagalà, E. L. Belsito, A. Comande, M. Greco and A. Liguori, *Chem. Cent. J.*, 2017, **11**, 87.
- 19 Z. Q. Fu, X. H. Wang, S. Tao, Q. Q. Bu, D. H. Wei and N. Liu, *J. Org. Chem.*, 2021, **86**, 2339–2358.
- 20 Basavaprabhu, K. Muniyappa, N. R. Panguluri, P. Veladi and V. V. Sureshbabu, *New J. Chem.*, 2015, **39**, 7746–7749.





- 21 A. C. Shekhar, A. R. Kurnar, G. Sathaiah, V. L. Paul, M. Sridhar and P. S. Rao, *Tetrahedron Lett.*, 2009, **50**, 7099–7101.
- 22 J.-G. Kim and D. O. Jang, *Bull. Korean Chem. Soc.*, 2010, **31**, 2989–2991.
- 23 H. Lundberg, F. Tinnis, N. Selander and H. Adolfsson, *Chem. Soc. Rev.*, 2014, **43**, 2714–2742.
- 24 H. Lundberg, F. Tinnis and H. Adolfsson, *Chem. – Eur. J.*, 2012, **18**, 3822–3826.
- 25 J.-G. Kim and D. O. Jang, *Synlett*, 2010, 1231–1234.
- 26 M. Hong and G. Xiao, *J. Fluorine Chem.*, 2013, **146**, 11–14.
- 27 A. M. Gabdullin, R. N. Kadikova, O. S. Mozgovoij and I. R. Ramazanov, *ChemistrySelect*, 2023, **8**, 7.
- 28 W. Muramatsu and H. Yamamoto, *J. Am. Chem. Soc.*, 2019, **141**, 18926–18931.
- 29 P. V. Ramachandran, A. A. Alawaed and H. J. Hamann, *J. Org. Chem.*, 2022, **87**, 13259–13269.
- 30 P. V. Ramachandran, A. A. Alawaed and H. J. Hamann, *Org. Lett.*, 2022, **24**, 8481–8486.
- 31 P. V. Ramachandran, A. A. Alawaed and A. Singh, *Molecules*, 2023, **28**, 4575.
- 32 P. V. Ramachandran and A. A. Alawaed, *Molecules*, 2023, **28**, 60.
- 33 P. V. Ramachandran, A. A. Alawaed and H. J. Hamann, *Org. Lett.*, 2023, **25**, 6902–6906.
- 34 P. V. Ramachandran, A. A. Alawaed and H. J. Hamann, *Org. Lett.*, 2023, **25**, 4650–4655.
- 35 The first letter of the amide number refers to the acid and the second letter refers to the amine used.
- 36 M. Mader and P. Helquist, *Tetrahedron Lett.*, 1988, **29**, 3049–3052.
- 37 H. Lundberg, F. Tinnis and H. Adolfsson, *Synlett*, 2012, 2201–2204.
- 38 Sonam, V. N. Shinde and A. Kumar, *J. Org. Chem.*, 2022, **87**, 2651–2661.
- 39 E. Jorda, A. Tuel, R. Teissier and J. Kervennal, *Zeolites*, 1997, **19**, 238–245.
- 40 G. B. Nikiforov, C. Knapp, J. Passmore and A. Decken, *J. Fluorine Chem.*, 2006, **127**, 1398–1404.
- 41 E. L. Muetterties, *J. Am. Chem. Soc.*, 1960, **82**, 1082–1087.
- 42 J. A. Chandler and R. S. Drago, *Inorg. Chem.*, 1962, **1**, 356.
- 43 J. Tani, T. Oine and I. Inoue, *Synthesis*, 1975, 714–715.
- 44 T. J. Shen, V. T. Hanh, T. Q. Nguyen, M. K. Jhan, M. R. Ho and C. F. Lin, *Front. Cell. Infect. Microbiol.*, 2020, **10**, 606743.
- 45 J. H. Dam, G. Osztrovsky, L. U. Nordstrøm, and R. Madsen, *Chem. – Eur. J.*, 2010, **16**, 6820–6827.
- 46 H. Lundberg, F. Tinnis, J. J. Zhang, A. G. Algarra, F. Himo and H. Adolfsson, *J. Am. Chem. Soc.*, 2017, **139**, 2286–2295.

