

CORRECTION

View Article Online
View Journal | View Issue



Cite this: *Org. Biomol. Chem.*, 2024, **22**, 395

Correction: π -Facial selectivity in the Diels–Alder reaction of glucosamine-based chiral furans and maleimides


Cornelis H. M. van der Loo,^a Rutger Schim van der Loeff,^a Avelino Martín,^c Pilar Gomez-Sal,^c Mark L. G. Borst,^b Kees Pouwer^b and Adriaan J. Minnaard^{*a}

DOI: 10.1039/d3ob90168h

rsc.li/obc

Correction for ' π -Facial selectivity in the Diels–Alder reaction of glucosamine-based chiral furans and maleimides' by Cornelis H. M. van der Loo *et al.*, *Org. Biomol. Chem.*, 2023, **21**, 1888–1894, <https://doi.org/10.1039/D2OB02221D>.

The authors regret that there was an error in the depiction of a specific stereocenter found in the key structures **14–16** and **18** throughout the article (Scheme 1, Scheme 4, Fig. 1, Scheme 7 and Scheme 9). The stereocenter is inverted in each instance. The correct figures and schemes are shown below.




Scheme 1 Recent advances in the application of carbohydrate derived furans in direct Diels–Alder cycloaddition reactions with maleimides.

^aDepartment of Chemical Biology, Stratingh Institute for Chemistry, University of Groningen, Groningen, The Netherlands. E-mail: A.J.Minnaard@RUG.nl

^bSymeres B.V., Kadijk 3, 9747 AT Groningen, The Netherlands

^cDepartamento de Química Orgánica y Química Inorgánica, Instituto de Investigación Química "Andrés M. Del Río" (IQAR), Universidad de Alcalá, Alcalá de Henares, 28805 Madrid, Spain





Scheme 4 Diels-Alder reactions of Di-HCF (10) with substituted maleimides afford a mixture of four different diastereomers. Periodate oxidation simplifies the structure to facilitate NMR analysis. Top ^1H -NMR spectrum: before periodate treatment; bottom ^1H -NMR spectrum: after periodate treatment.






Fig. 1 The ^1H -NMR signal used to determine the product distribution. The proton of the primary alcohol (highlighted in purple) has a different chemical shift for each of the four diastereomeric products. NMR solvent: $\text{DMSO}-d_6$.



Scheme 7 Synthesis of crystalline derivative **16** to enable X-ray diffraction.



Scheme 9 Diels-Alder cycloaddition of Di-HAF (1) and *N*-phenylmaleimide in chloroform–DMSO at $-10\text{ }^\circ\text{C}$. *endo/exo*-ratio: 2 : 8 and *exo*-facial selectivity: 4 : 1, isolated yield of major isomer: 59%.



The same error was also found in the electronic supplementary information (ESI) (section S12). The structure was corrected in the updated replacement ESI file on 6th December 2023.

Additionally, in the Experimental section in the ESI two chemical names (in sections S9 and S10) contain the wrong stereo-annotation. The corrected chemical names are shown below.

Section S9:

benzyl ((4*R*,7*R*)-7-((*R*)-1,2-dihydroxyethyl)-1,3-dioxo-2-phenyl-2,3,3a,4,7,7a-hexahydro-1*H*-4,7-epoxyisoindol-5-yl)carbamate (**DAP 14**) should read benzyl ((3a*S*,4*S*,7*S*,7a*R*)-7-((*R*)-1,2-dihydroxyethyl)-1,3-dioxo-2-phenyl-2,3,3a,4,7,7a-hexahydro-1*H*-4,7-epoxyisoindol-5-yl)carbamate (**DAP 14**)

Section S10:

benzyl ((4*R*,5*R*,7*R*)-7-((*R*)-1,2-dihydroxyethyl)-1,3-dioxo-2-phenyloctahydro-1*H*-4,7-epoxyisoindol-5-yl)carbamate (**DAPRED 15**) should read benzyl ((3a*S*,4*S*,5*S*,7*S*,7a*R*)-7-((*R*)-1,2-dihydroxyethyl)-1,3-dioxo-2-phenyloctahydro-1*H*-4,7-epoxyisoindol-5-yl)carbamate (**DAPRED 15**)

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

