



Cite this: *Org. Biomol. Chem.*, 2024, **22**, 2395

## Synthesis of galactomannan fragments to help NMR assignment of polysaccharides extracted from lichens†

Louis-Philippe David,<sup>a</sup> Solenn Ferron,<sup>b</sup> Bénédicte Favreau,<sup>a</sup> Ozgur Yeni,<sup>c</sup> Simon Ollivier,<sup>d,e</sup> David Ropartz,<sup>d,e</sup> Isabelle Compagnon,<sup>id</sup> Vincent Ferrières,<sup>id</sup>\*<sup>a</sup> Françoise Le Dévéhat\*<sup>b</sup> and Laurent Legentil<sup>id</sup>\*<sup>a</sup>

The synthesis of six model trisaccharides representative of galactomannans produced by lichens was performed through stereoselective glycosylation. These standards include linear and branched galactomannans bearing either galactofuranosyl or galactopyranosyl entities. The complete assignment of <sup>1</sup>H and <sup>13</sup>C signals for both forms of synthetically reduced oligosaccharides was performed. The resulting NMR data were used to quickly demonstrate the structural characteristics of minor polysaccharides within different extracts of three representative lichens.

Received 10th January 2024,  
Accepted 20th February 2024


DOI: 10.1039/d4ob00047a

rsc.li/obc

## Introduction

Lichens constitute a “self-sustaining ecosystem” where the main partners are photosynthetic organisms (microalgae and/or cyanobacteria) and fungi.<sup>1</sup> Such symbiosis relies on the fruitful exchange of metabolites and nutrients between both partners and allows the lichens to grow in all habitats even under harsh conditions like sun-exposed rocks. To date, lichen chemistry has mainly focused on secondary metabolites (more than 1000 compounds have been isolated and identified in the last compendium).<sup>2</sup> Such low molecular weight metabolites represent 3–20% of the lichen dry weight while free sugars and polysaccharides represent about 60% of the extractable biomass.<sup>3</sup> These polysaccharides contribute to the extreme tolerance of the lichens towards harsh conditions, their fascinating revival ability, their very unique management of water and the maintenance of the complex partnership, including a specific microflora within this long-living microecosystem.<sup>4–6</sup> Three main families of polysaccharides could be found among

the few lichens that have been studied for their carbohydrate constituents.<sup>7,8</sup>  $\beta$ - and  $\alpha$ -glucans are common homopolymers found throughout the species while galactomannans are much more heterogeneous. The latter biomolecules are built on a mannan backbone with different decorations depending on the symbiotic partners. For example, polysaccharides from *Cetraria islandica*<sup>9</sup> and *Lasallia pustulata*<sup>10</sup> have a common (1  $\rightarrow$  6) linked  $\alpha$ -D-mannopyranosyl main chain branched for the first one with  $\beta$ -D-galactopyranose (Galp) at positions 2 and 4, and for the second one with  $\beta$ -D-galactofuranose (Galf) at the same positions (Fig. 1A). The quite rare galactofuranose moiety was also identified within the galactomannans of *Umbilicaria*,<sup>11</sup> *Usnea*,<sup>12</sup> *Gyrophora*<sup>13</sup> or *Roccella*.<sup>14</sup> For the last lichen, the mannan backbone is made up of  $\alpha$ -(1  $\rightarrow$  4) linked mannopyranose with galactofuranose at the non-reducing end



**Fig. 1** General structures of (A) polysaccharides present in *Lasallia*, *Cetraria* and (B) *Roccella* lichens.

<sup>a</sup>Univ Rennes, Ecole Nationale Supérieure de Chimie de Rennes, CNRS, ISCR – UMR 6226, F-35000 Rennes, France. E-mail: laurent.legentil@ensc-rennes.fr

<sup>b</sup>Univ Rennes, CNRS, ISCR – UMR 6226, F-35000 Rennes, France

<sup>c</sup>Univ Lyon, Université Claude Bernard Lyon 1, CNRS, Institut Lumière Matière, F-69622 Villeurbanne, France

<sup>d</sup>INRAE, UR BIA, F-44316 Nantes, France

<sup>e</sup>INRAE, BIBS Facility, F-44316 Nantes, France

† Electronic supplementary information (ESI) available: Experimental procedures and related spectroscopic data for all intermediaries 4–23, <sup>1</sup>H and <sup>13</sup>C indexation table of trisaccharides 1–3, 2D HSQC maps of lichen extracts, and <sup>1</sup>H and <sup>13</sup>C NMR spectra of all intermediates and final products. See DOI: <https://doi.org/10.1039/d4ob00047a>



(Fig. 1B). Recently, such diversity of structures was outlined as a chemotaxonomic character to classify lichens from different species.<sup>8</sup> Some of them were also evaluated for their antitumor, antiviral and immunostimulating properties.<sup>15</sup> For example, both polysaccharides KI-M-7 and thamnolan, isolated from *C. islandica* and *Thamnolia subuliformis*, respectively, showed an increase in the phagocytic activity of granulocytes in the presence of polymers and anticomplementary effects.<sup>16,17</sup>

Nevertheless, the studies on such structures remain scarce, due to the difficulty associated with their harvest, extraction and characterisation. The sensitivity of galactofuranosides towards mild hydrolysis conditions also hampers reliable characterisation studies.

Previously, their primary structures were determined after different fractionation steps followed by NMR analysis. The characterisation of the regioisomery relied mainly on <sup>13</sup>C data tables that dated from the early 70s.<sup>18</sup> Recently, Nifantiev *et al.* highlighted the difficulty of correlating such <sup>13</sup>C chemical shifts with the sequence of the polysaccharide by comparing synthetic galactofuranomannans with the extracellular galactomannan of *Aspergillus fumigatus*.<sup>19</sup> No such systematic comparison by NMR of oligogalactomannans with different extracts of lichens has been reported yet.

To efficiently perform such screening, we describe herein the synthesis of a library of six trisaccharides that could serve as analytical standards (Fig. 2). Linear  $\beta$ -D-Galf-(1  $\rightarrow$  6)- $\alpha$ -D-Manp-(1  $\rightarrow$  4)- $\alpha$ -D-Manp **1a**, linear and branched trisaccharides  $\beta$ -D-Galf-(1  $\rightarrow$  4)- $\alpha$ -D-Manp-(1  $\rightarrow$  6)- $\alpha$ -D-Manp **2a** and  $\alpha$ -D-Manp-(1  $\rightarrow$  6)-[ $\beta$ -D-Galf-(1  $\rightarrow$  4)]- $\alpha$ -D-Manp **3a** were first synthesized as representative galactomannans from *Rocella decipiens*<sup>14</sup> and *Lasallia pustulata*<sup>10</sup> species. Moreover, in order to pursue our efforts in the development of analytical approaches allowing differentiation between galactosyl tautomers,<sup>20</sup> we also prepared the trisaccharides bearing Galp residues **1b–3b**. This is crucial as Galp-Man motifs were also described in some lichens like *Cetraria islandica*.<sup>9</sup> All standards were obtained as

hemiacetals, the main product of hydrolysis of natural galactomannans.

To complete this study, the NMR spectra of all trisaccharides were compared with different fractions of extracts of our representative lichen thalli. We demonstrate here that the presence of biologically relevant galactomannans could be detected in unsuspected fractions, not reported before.

## Results and discussion


### Synthesis of trisaccharides

The strategy implemented in this study capitalized on recent works on the synthesis of the four possible regioisomers of Galf-Manp and Galp-Manp disaccharides.<sup>20–22</sup> In each case, the synthesis of **a**-family derivatives (Galf family) and **b**-family compounds (Galp one) followed a similar sequence of reactions.

The synthetic cascade to afford linear trisaccharides **1** started from known thioglycosides **4**<sup>20</sup> with the established link Gal-(1  $\rightarrow$  6)-Manp. They were first converted into the corresponding trichloroacetimidates **6** (Scheme 1). Hydrolysis of **4** assisted by *N*-iodosaccharin afforded acetals **5**, which were reacted with trichloroacetonitrile to give **6**. Then, glycosylation of acceptor **7** with the free *O*-4 position was performed in the presence of a catalytic amount of trimethylsilyl trifluoromethanesulfonate (TMSOTf). Subsequent cleavage of the acetonide group in **8** in the presence of an excess of trifluoroacetic acid in dichloromethane gave **9**. This reaction turned out to be sluggish for obtaining **9a** and maximum yield was limited to 30%. Addition of dithiothreitol (DTT) allowed the increase of the yield to 41%. As for **9b**, a good yield of 66% was obtained without DTT. All ester groups were then removed by Zemplen transesterification.

Finally, in order to get as close as possible to structures that would result from controlled hydrolysis of native polysaccharides, the targeted triglycosides **1a** and **1b** were obtained with excellent yields thanks to the action of the bromonium ion and water on thioglycosides **10**.


A very similar approach was performed to synthesize trisaccharides, characterized by a Manp-(1  $\rightarrow$  6)-Manp skeleton grafted with a Gal residue at position 4 (Scheme 2). While building block **11a** was already described,<sup>20</sup> the pyranosidic counterpart **11b** had to be synthesized first through glycosylation of thiophenyl 4-hydroxyl mannopyranoside **7** by tetra-*O*-benzoyl- $\beta$ -D-galactopyranosyl trichloroacetimidate according to a standard procedure. Then, to gain access to linear compounds **2**, the thioglycosides **11** in hand were first transformed into the corresponding trichloroacetimidate donors **13** in two steps. Despite the absence of a participating group at the *C*-2 position of both donors, glycosylation of **14** yielded the desired Gal-(1  $\rightarrow$  4)-Manp-(1  $\rightarrow$  6)-Manp trisaccharides **15a** and **15b** with  $\alpha$ -(1  $\rightarrow$  6)-linkages only. Removal of the acetonide group was performed this time in hot aqueous acetic acid media to avoid by-product formation. This was followed by deacylation



**Fig. 2** Targeted trisaccharides **1–3** as standards for NMR comparison with extracted fractions from the lichen thalli (series **a** correspond to Galf-containing trisaccharides and **b** to Galp-containing ones).








**Scheme 2** Synthesis of the trisaccharides **2** and **3** based on the Manp-(1,6)-Manp skeleton. Reagents and conditions: (i) NBS, H<sub>2</sub>O/Me<sub>2</sub>CO; (ii) Cl<sub>3</sub>CCN, CsCO<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>; (iii) **14**, TMSOTf, CH<sub>2</sub>Cl<sub>2</sub>; (iv) AcOH, H<sub>2</sub>O; (v) NaOMe, MeOH; (vi) NIS, CH<sub>3</sub>CN/H<sub>2</sub>O; (vii) BzCl, Pyr; (viii) AcCl, MeOH; (ix) **21**, TMSOTf, CH<sub>2</sub>Cl<sub>2</sub>.

presence of around 15% of acetylated glucose in pustulan. As for the minor constituent of both fractions, <sup>1</sup>H and <sup>13</sup>C chemical shifts characteristic of galactomannans could be found (Fig. 3A). Interestingly, such a pattern was never reported before on the hot-water extract A2. Therefore, the <sup>1</sup>H signal at 5.06 ppm ( $\delta_C$  107.53 ppm) showed a good match with the anomeric proton of trisaccharides **2a** and **3a**. Interestingly the anomeric proton at 4.93 ppm ( $\delta_C$  99.25 ppm) fits well with signals corresponding to the branched trisaccharides **3a** rather than the linear one **2a**. This confirmed the prevalence of the branched galactomannan oligomer over Galf<sub>2</sub>-ending mannans.

No Galp was identified in the extract which is in accordance with the literature.

The same trend was observed using the extracts of the *Roccella thalli*. Again unexpectedly, hot water extract A2 contained galactose in its furanose form ( $\delta_H$  5.00 ppm and  $\delta_C$  107.69 ppm in Fig. 3B). A comparison with standard **1a** confirmed the previously reported Galf-(1 → 6)-Manp-(1 → 4) sequence ( $\delta_H$  5.17 ppm and  $\delta_C$  100.68 ppm for mannose anomeric proton). No correlation with the pyranose analogues was identified. Finally, we compared **1–3** with the water-soluble fractions from *Cetraria*. While a good match was found





**Fig. 3** Identification of galactomannans in the natural extract from lichens. HSQC spectrum of the anomeric region of the fractions A2 of *L. pustulata* (A), *R. fuciformis* (B) and *C. islandica* (C) and comparison with trisaccharides 1–3 as standards (D). HSQC correlations are shown in black for lichen extracts; blue for 1a; red for 2a; pink for 2b; green for 3a. Signals in the anomeric region corresponding to putative galactomannans are labelled.

with standards 2b and 3b, thus confirming the presence of the  $\beta\text{-Galp}(1 \rightarrow 4)\text{-}\alpha\text{-Manp}$  motif ( $\delta_{\text{H}} 4.44$  ppm and  $\delta_{\text{C}} 103.06$  ppm in Fig. 3C), some correlation spots were also visible around 107 ppm ( $\delta_{\text{H}} 5.20$  ppm and  $\delta_{\text{C}} 106.72$  ppm). While this chemical shift differed from the one found for  $\beta\text{-Galp}(1 \rightarrow 4)\text{-}$  and  $\beta\text{-Galp}(1 \rightarrow 6)\text{-}\alpha\text{-Manp}$ , it is nevertheless representative of the presence of the furanosyl moiety, probably linked in  $\beta\text{-}(1 \rightarrow 2)$  on the mannopyranose core. It is worth noting that this signal could also be found in both *Lassalia* and *Rocella* fractions.

## Conclusion

Six different oligogalactomannans were synthesized and their HSQC maps were compared with the natural polysaccharides extracted from three different lichens. Thanks to this non-destructive method, the presence of galactomannans could be identified in most fractions, even in the hot water extract which is supposed to contain mostly pustulan. Unexpectedly, the rare hexofuranose form was identified in lichens that are

supposed to contain only the pyranose isomers. It confirms that such galactofuranomannans constitute a unique feature of the symbiosis.

## Experimental

### General procedure for hydrolysis of thiomannosides using *N*-iodosaccharin (A1)

To a 9/1 (v/v) mixture of  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ , thiosaccharide and *N*-iodosaccharin were successively added. The reaction mixture was stirred at RT and monitored by TLC. After the disappearance of the starting compound, the reaction was quenched by adding triethylamine and concentrated under reduced pressure. The product was partitioned between  $\text{CH}_2\text{Cl}_2$  and water, and the organic layer was washed with water. The combined organic layers were dried over  $\text{MgSO}_4$ , filtered and concentrated under reduced pressure. The product was finally chromatographically purified.



### General procedure for hydrolysis of thiomannosides using *N*-bromosuccinimide (A2)

To a 9/1 (v/v) mixture of acetone/H<sub>2</sub>O, thiosaccharide and *N*-bromosuccinimide were successively added. The reaction mixture was stirred at RT and monitored by TLC. After the disappearance of the starting compound, the reaction medium was diluted with CH<sub>2</sub>Cl<sub>2</sub> and the resulting solution was washed with a saturated aqueous solution of NaHCO<sub>3</sub> (3×). The resulting organic phase was dried over MgSO<sub>4</sub>, filtered and concentrated under reduced pressure. The product was finally chromatographically purified.

### General procedure for Zemplén transesterification (B)

Into anhydrous MeOH, acylated trisaccharide and a volume of a 0.54 M solution of MeONa in MeOH were successively added. The mixture was stirred overnight at RT and then neutralized with IR 120 (H<sup>+</sup>-form). After filtration, the solvent was removed under reduced pressure. The residue was suspended in water and the biphasic solution was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3×). The resulting aqueous layer was finally freeze-dried.

### General procedure for glycosylation (C)

Into anhydrous CH<sub>2</sub>Cl<sub>2</sub>, both the acceptor and donor and then activated 4 Å molecular sieves were added. The mixture was cooled to 0 °C and stirred for 15 min before dropwise addition of TMSOTf. The temperature was maintained at 0 °C and the reaction was monitored by TLC (cyclohexane/AcOEt 7/3). When no starting material remained, triethylamine was added and the solution was filtered on a pad of Celite. The resulting filtrate was evaporated under vacuum before flash-chromatography.

### General procedure for synthesis of trichloroacetimidate (D)

Into anhydrous CH<sub>2</sub>Cl<sub>2</sub>, the reducing saccharide, Cs<sub>2</sub>CO<sub>3</sub>, and trichloroacetonitrile were successively added. The resulting mixture was stirred at RT until no starting material remained. After filtration, the resulting filtrate was concentrated under reduced pressure and the product was purified by flash-chromatography.

### β-D-Galactofuranosyl-(1 → 6)-α-D-mannopyranosyl-(1 → 4)-α,β-D-mannopyranose (1a)

To a solution of **10a** (47 mg, 0.079 mmol) in a CH<sub>3</sub>CN/H<sub>2</sub>O 1 : 1 mixture (5 mL), *N*-iodosaccharin (87 mg, 0.28 mmol) was added. The reaction mixture was stirred at RT for 3 h. After the disappearance of the starting compound, the reaction was quenched by adding triethylamine and concentrated under reduced pressure. The resulting crude was partitioned between CH<sub>2</sub>Cl<sub>2</sub> and water, and the aqueous layer was washed with CH<sub>2</sub>Cl<sub>2</sub> (3×). The resulting aqueous phase was freeze-dried and the resulting residue was purified by column chromatography (AcOEt/MeOH/H<sub>2</sub>O gradient from 8 : 1 : 1 up to 6 : 2 : 2) to afford **1a** (36 mg, 90%) as a white solid. TLC: (BuOH/AcOH/H<sub>2</sub>O, 6 : 2 : 2), *R*<sub>f</sub> = 0.23.

**1aα**: <sup>1</sup>H NMR (D<sub>2</sub>O, 400 MHz): δ 5.20 (1H, d, *J*<sub>1b,2b</sub> 1.5, H-1b), 5.17 (1H, d, *J*<sub>1a,2a</sub> 1.8, H-1a), 5.05 (1H, d, *J*<sub>1c,2c</sub> 1.8, H-1c), 4.14 (1H, dd, *J*<sub>2c,3c</sub> 3.7, *J*<sub>2c,1c</sub> 1.8, H-2c), 4.09–4.03 (2H, m, H-2b, H-6b), 4.07 (1H, dd, *J*<sub>3c,4c</sub> 6.5, *J*<sub>3c,2c</sub> 3.7, H-3c), 4.00 (1H, dd, *J*<sub>4c,3c</sub> 6.5, *J*<sub>4c,5c</sub> 4.2, H-4c), 3.97 (1H, dd, *J*<sub>3a,4a</sub> 9.0, *J*<sub>3a,2a</sub> 3.4, H-3a), 3.92–3.76 (8H, m, H-2a, H-4a, H-5a, H-6a, H-3b, H-5b, H-5c), 3.76–3.63 (4H, m, H-4b, H-6'b, H-6c). <sup>13</sup>C NMR (D<sub>2</sub>O, 126 MHz): δ 107.75 (C-1c), 101.74 (C-1b), 93.83 (C-1a), 82.83 (C-4c), 81.05 (C-2c), 76.85 (C-3c), 75.10 (C-4a), 72.56 (C-5b), 71.17 (C-2a), 71.03, 70.87 (C-5a, C-5c), 70.60 (C-3a), 70.27 (C-2b, C-3b), 66.80 (C-6b), 66.62 (C-4b), 62.76 (C-6c), 61.19 (C-6a).

**1aβ**: <sup>1</sup>H NMR (D<sub>2</sub>O, 500 MHz): δ 5.25 (1H, d, *J*<sub>1b,2b</sub> 1.7, H-1b), 5.05 (1H, d, *J*<sub>1c,2c</sub> 1.8, H-1c), 4.89 (1H, s, *J*<sub>1a,2a</sub> 1.8, H-1a), 4.14 (1H, dd, *J*<sub>2c,3c</sub> 3.7, *J*<sub>2c,1c</sub> 1.8, H-2c), 4.09–4.03 (2H, m, H-2b, H-6b), 4.07 (1H, dd, *J*<sub>3c,4c</sub> 6.5, *J*<sub>3c,2c</sub> 3.7, H-3c), 4.00 (1H, dd, *J*<sub>4c,3c</sub> 6.5, *J*<sub>4c,5c</sub> 4.2, H-4c), 3.92–3.76 (6H, m, H-2a, H-3a, H-6a, H-3b, H-5b, H-5c), 3.76–3.63 (6H, m, H-4a, H-6'a, H-4b, H-6'b, H-6c), 3.53–3.44 (1H, m, H-5a). <sup>13</sup>C NMR (D<sub>2</sub>O, 100 MHz): δ 107.75 (C-1c), 101.74 (C-1b), 93.61 (C-1a), 82.83 (C-4c), 81.05 (C-2c), 76.85 (C-3c), 74.77, 74.69 (C-4a, C-5a), 73.48 (C-3a), 72.56 (C-5b), 71.17 (C-2a), 70.87 (C-5c), 70.27 (C-2b, C-3b), 66.80 (C-6b), 66.62 (C-4b), 62.76 (C-6c), 60.93 (C-6a).

HRMS (ESI): calcd for C<sub>18</sub>H<sub>32</sub>O<sub>16</sub>Na [M + Na]<sup>+</sup> 527.1588, found 527.1585.

### β-D-Galactopyranosyl-(1 → 6)-α-D-mannopyranosyl-(1 → 4)-α,β-D-mannopyranose (1b)

To a solution of **10b** (60 mg, 0.1 mmol) in a 9 : 1 acetone/H<sub>2</sub>O mixture (6 mL), *N*-bromosuccinimide (21 mg, 0.12 mmol) was added. The reaction mixture was stirred at RT for 3 h. After the disappearance of the starting compound, the reaction was quenched by adding triethylamine and concentrated under reduced pressure. The resulting crude was partitioned between CH<sub>2</sub>Cl<sub>2</sub> and water, and the aqueous layer was washed with CH<sub>2</sub>Cl<sub>2</sub> (3×). The resulting aqueous phase was freeze-dried and the resulting residue was purified by column chromatography (AcOEt/MeOH/H<sub>2</sub>O gradient from 8 : 1 : 1 up to 6 : 2 : 2) to afford **1b** (50 mg, 99%) as a white solid. TLC: (AcOEt/MeOH/H<sub>2</sub>O, 6 : 2 : 2), *R*<sub>f</sub> = 0.12.

**1bα**: <sup>1</sup>H NMR (D<sub>2</sub>O, 400 MHz): δ 5.22 (1H, d, *J*<sub>1b,2b</sub> 1.8, H-1b), 5.17 (1H, d, *J*<sub>1a,2a</sub> 1.9, H-1a), 4.45 (1H, d, *J*<sub>1c,2c</sub> 7.8, H-1c), 4.22 (1H, d, *J*<sub>6b,6'b</sub> 9.5, H-6b), 4.07 (1H, dd, *J*<sub>2b,3b</sub> 3.3, *J*<sub>2b,1b</sub> 1.8, H-2b), 3.97 (1H, dd, *J*<sub>3a,4a</sub> 9.0, *J*<sub>3a,2a</sub> 3.5, H-3a), 3.94–3.85 (6H, m, H-2a, H-5a, H-6a, H-5b, H-6'b, H-4c), 3.84–3.69 (7H, m, H-4a, H-6'a, H-3b, H-4b, H-5c, H-6c), 3.67 (1H, dd, *J*<sub>3c,2c</sub> 10.0, *J*<sub>3c,4c</sub> 3.4, H-3c), 3.57 (1H, dd, *J*<sub>2c,3c</sub> 10.0, *J*<sub>2c,1c</sub> 7.8, H-2c). <sup>13</sup>C NMR (D<sub>2</sub>O, 100 MHz): δ 103.36 (C-1c), 101.71 (C-1b), 93.78 (C-1a), 75.14 (C-5c), 74.94 (C-4a), 72.62, 72.59 (C-5b, C-3c), 71.13 (C-2a), 70.90 (C-5a), 70.81 (C-2c), 70.61 (C-3a), 70.21 (C-2b, C-3b), 68.66 (C-6b, C-4c), 66.33 (C-4b), 61.16, 61.01 (C-6a, C-6c).

**1bβ**: <sup>1</sup>H NMR (D<sub>2</sub>O, 400 MHz): δ 5.22 (1H, d, *J*<sub>1b,2b</sub> 1.8, H-1b), 4.89 (1H, d, *J*<sub>1a,2a</sub> 1.0, H-1a), 4.45 (1H, d, *J*<sub>1c,2c</sub> 7.8, H-1c), 4.22 (1H, d, *J*<sub>6b,6'b</sub> 9.5, H-6b), 4.07 (1H, dd, *J*<sub>2b,3b</sub> 3.3, *J*<sub>2b,1b</sub> 1.8, H-2b), 3.97 (1H, dd, *J*<sub>3a,4a</sub> 9.0, *J*<sub>3a,2a</sub> 3.5, H-3a), 3.94–3.85 (5H,



m, H-2a, H-6a, H-5b, H-6'b, H-4c), 3.84–3.69 (7H, m, H-4a, H-6'a, H-3b, H-4b, H-5c, H-6c), 3.67 (1H, dd,  $J_{3c,2c}$  10.0,  $J_{3c,4c}$  3.4, H-3c), 3.57 (1H, dd,  $J_{2c,3c}$  10.0,  $J_{2c,1c}$  7.8, H-2c), 3.47 (1H, ddd,  $J_{5a,4a}$  9.6,  $J_{5a,6'a}$  6.1,  $J_{5a,6a}$  2.2, H-5a).  $^{13}\text{C}$  NMR ( $\text{D}_2\text{O}$ , 100 MHz):  $\delta$  103.33 (C-1c), 101.65 (C-1b), 93.57 (C-1a), 75.14 (C-5c), 74.58 (C-5a), 71.48 (C-4a), 72.62, 72.59 (C-5b, C-3c), 71.70 (C-2a), 70.81 (C-2c), 70.61 (C-3a), 70.21 (C-2b, C-3b), 68.66 (C-6b, C-4c), 66.33 (C-4b), 61.16, 61.01 (C-6a, C-6c).

HRMS (ESI): calcd for  $\text{C}_{18}\text{H}_{32}\text{O}_{16}\text{Na}$  [ $\text{M} + \text{Na}$ ] $^+$  527.1588, found 527.1582.

#### $\beta$ -D-Galactofuranosyl-(1 $\rightarrow$ 4)- $\alpha$ -D-mannopyranosyl-(1 $\rightarrow$ 6)- $\alpha,\beta$ -D-mannopyranose (2a)

To a solution of **17a** (40 mg, 0.07 mmol) in an acetone/ $\text{H}_2\text{O}$  9 : 1 mixture (6 mL), *N*-bromosuccinimide (36 mg, 0.2 mmol) was added. The reaction mixture was stirred at RT for 3 h. After the disappearance of the starting compound, the reaction was quenched by adding triethylamine and concentrated under reduced pressure. The resulting crude was partitioned between  $\text{CH}_2\text{Cl}_2$  and water, and the aqueous layer was washed with  $\text{CH}_2\text{Cl}_2$  (3 $\times$ ). The resulting aqueous phase was freeze-dried and the resulting residue was purified by column chromatography (AcOEt/MeOH/ $\text{H}_2\text{O}$  gradient from 8 : 1 : 1 up to 6 : 2 : 2) to afford **2a** (20 mg, 59%) as a white solid.

**2a $\alpha$** :  $^1\text{H}$  NMR ( $\text{D}_2\text{O}$ , 500 MHz):  $\delta$  5.17 (1H, d,  $J_{1a,2a}$  1.9, H-1a), 5.08 (1H, d,  $J_{1c,2c}$  2.3, H-1c), 4.72 (1H, d,  $J_{1b,2b}$  1.1, H-1b), 4.17 (1H, dd,  $J_{6a,6'a}$  11.1,  $J_{6a,5a}$  2.0, H-6a), 4.12–4.07 (4H, m, H-2b, H-2c, H-3c, H-4c), 3.99–3.91 (3H, m, H-2a, H-5a, H-6b), 3.89–3.77 (4H, m, H-3a, H-6'a, H-6'b, H-5c), 3.77–3.59 (5H, m, H-4a, H-3b, H-4b, H-6c), 3.52–3.46 (1H, m, H-5b).  $^{13}\text{C}$  NMR ( $\text{D}_2\text{O}$ , 126 MHz):  $\delta$  107.91 (C-1c), 100.44 (C-1b), 94.09 (C-1a), 82.62 (C-2c), 80.91 (C-4c), 75.91 (C-3c), 75.14 (C-4b), 75.04 (C-5b), 71.44 (C-3b), 71.24 (C-5a), 70.58 (C-2a), 70.48 (C-3a), 70.23 (C-2b), 70.12 (C-5c), 68.60 (C-6a), 66.67 (C-4a), 62.67 (C-6c), 60.43 (C-6b).

**2a $\beta$** :  $^1\text{H}$  NMR ( $\text{D}_2\text{O}$ , 500 MHz):  $\delta$  5.08 (1H, d,  $J_{1c,2c}$  2.3, H-1c), 4.91 (1H, d,  $J_{1a,2a}$  1.0, H-1a), 4.73 (1H, d,  $J_{1b,2b}$  1.1, H-1b), 4.20 (1H, dd,  $J_{6a,6'a}$  11.3,  $J_{6a,5a}$  2.1, H-6a), 4.12–4.07 (4H, m, H-2b, H-2c, H-3c, H-4c), 3.99–3.91 (2H, m, H-2a, H-6b), 3.89–3.77 (5H, m, H-3a, H-5a, H-6'a, H-6'b, H-5c), 3.77–3.59 (5H, m, H-4a, H-3b, H-4b, H-6c), 3.56–3.50 (1H, m, H-5b).  $^{13}\text{C}$  NMR ( $\text{D}_2\text{O}$ , 126 MHz):  $\delta$  107.91 (C-1c), 100.47 (C-1b), 93.72 (C-1a), 82.62 (C-2c), 80.91 (C-4c), 75.91 (C-3c), 74.90 (C-5b), 72.94 (C-4b), 71.44 (C-3b), 71.16 (C-5a), 70.58 (C-2a), 70.48 (C-3a), 70.23 (C-2b), 70.12 (C-5c), 68.56 (C-6a), 66.43 (C-4a), 62.67 (C-6c), 61.04 (C-6b).

HRMS (ESI): calcd for  $\text{C}_{18}\text{H}_{32}\text{O}_{16}\text{Na}$  [ $\text{M} + \text{Na}$ ] $^+$  527.1588, found 527.15891.

#### $\beta$ -D-Galactofuranosyl-(1 $\rightarrow$ 4)- $\alpha$ -D-mannopyranosyl-(1 $\rightarrow$ 6)- $\alpha,\beta$ -D-mannopyranose (2b)

To a solution of **17b** (12 mg, 0.02 mmol) in a  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$  1 : 1 mixture (5 mL), *N*-iodosaccharine (14 mg, 0.045 mmol) was added. The reaction mixture was stirred at RT for 3 h. After the disappearance of the starting compound, the reaction was quenched by adding triethylamine and concentrated

under reduced pressure. The resulting crude was partitioned between  $\text{CH}_2\text{Cl}_2$  and water, and the aqueous layer was washed with  $\text{CH}_2\text{Cl}_2$  (3 $\times$ ). The resulting aqueous phase was freeze-dried and the resulting residue was purified by column chromatography (AcOEt/MeOH/ $\text{H}_2\text{O}$  gradient from 8 : 1 : 1 up to 6 : 2 : 2) to afford **2b** (6 mg, 60%) as a white solid.

**2b $\alpha$** :  $^1\text{H}$  NMR ( $\text{D}_2\text{O}$ , 500 MHz):  $\delta$  5.17 (1H, d,  $J_{1a,2a}$  1.7, H-1a), 4.91 (1H, d,  $J_{1b,2b}$  1.7, H-1b), 4.45 (1H, d,  $J_{1c,2c}$  7.8, H-1c), 4.06 (1H, dd,  $J_{2b,3b}$  3.4,  $J_{2b,1b}$  1.7, H-2b), 4.00–3.91 (6H, m, H-2a, H-5a, H-6a, H-3b, H-4c, H-6c), 3.91–3.79 (4H, m, H-3a, H-4b, H-5b, H-6'c), 3.78–3.72 (5H, m, H-4a, H-6'a, H-6b, H-5c), 3.68 (1H, dd,  $J_{3c,2c}$  9.9,  $J_{3c,4c}$  3.3, H-3c), 3.55 (1H, dd,  $J_{2c,3c}$  9.9,  $J_{2c,1c}$  7.8, H-2c).  $^{13}\text{C}$  NMR ( $\text{D}_2\text{O}$ , 126 MHz): 103.02 (C-1c), 99.33 (C-1b), 94.17 (C-1a), 76.40 (C-4b), 75.37 (C-5c), 72.50 (C-3c), 71.28 (C-5b), 70.95 (C-2c), 70.63 (C-5a), 70.58 (C-2a), 70.41 (C-3a), 69.41 (C-2b), 69.32 (C-3b), 68.59 (C-4c), 66.66 (C-4a), 65.91 (C-6a), 61.10 (C-6b), 60.23 (C-6c).

**2b $\beta$** :  $^1\text{H}$  NMR ( $\text{D}_2\text{O}$ , 500 MHz):  $\delta$  4.92 (1H, d,  $J_{1b,2b}$  1.8, H-1b), 4.90 (1H, d,  $J_{1a,2a}$  1.0, H-1a), 4.45 (1H, d,  $J_{1c,2c}$  7.8, H-1c), 4.06 (1H, dd,  $J_{2b,3b}$  3.4,  $J_{2b,1b}$  1.8, H-2b), 4.00–3.91 (6H, m, H-2a, H-3a, H-6a, H-3b, H-4c, H-6c), 3.91–3.79 (3H, m, H-4b, H-5b, H-6'c), 3.78–3.72 (4H, m, H-6'a, H-6b, H-5c), 3.68 (1H, dd,  $J_{3c,2c}$  9.9,  $J_{3c,4c}$  3.3, H-3c), 3.67–3.64 (1H, m, H-4a), 3.58–3.49 (1H, m, H-5a), 3.55 (1H, dd,  $J_{2c,3c}$  9.9,  $J_{2c,1c}$  7.8, H-2c).  $^{13}\text{C}$  NMR ( $\text{D}_2\text{O}$ , 125 MHz): 103.02 (C-1c), 99.39 (C-1b), 93.81 (C-1a), 76.40 (C-4b), 75.37 (C-5c), 74.25 (C-5a), 73.14 (C-2a), 72.50 (C-3c), 71.28 (C-5b), 71.14 (C-3a), 70.95 (C-2c), 69.38 (C-2b), 69.30 (C-3b), 68.59 (C-4c), 66.45 (C-4a), 65.99 (C-6a), 61.10 (C-6b), 60.23 (C-6c).

HRMS (ESI): calcd for  $\text{C}_{18}\text{H}_{32}\text{O}_{16}\text{Na}$  [ $\text{M} + \text{Na}$ ] $^+$  527.1588, found 527.1583.

#### $[\beta$ -D-Galactofuranosyl-(1 $\rightarrow$ 4)]- $\alpha$ -D-mannopyranosyl-(1 $\rightarrow$ 6)- $\alpha,\beta$ -D-mannopyranose (3a)

To a solution of **23a** (20 mg, 0.03 mmol) in a  $\text{CH}_3\text{CN}/\text{H}_2\text{O}$  9 : 1 mixture (5 mL), *N*-iodosaccharine (16 mg, 0.05 mmol) was added. The reaction mixture was stirred at RT for 2 h. After the disappearance of the starting compound, the reaction was quenched by adding triethylamine and concentrated under reduced pressure. The resulting crude was partitioned between  $\text{CH}_2\text{Cl}_2$  and water, and the aqueous layer was washed with  $\text{CH}_2\text{Cl}_2$  (3 $\times$ ). After freeze-drying of the aqueous phase, the target trisaccharide **3a** was chromatographically purified on silica gel (AcOEt/MeOH/ $\text{H}_2\text{O}$  gradient from 8 : 1 : 1 up to 6 : 2 : 2) and isolated (8 mg, 50%) as a white solid.

**3a $\alpha$** :  $^1\text{H}$  NMR ( $\text{D}_2\text{O}$ , 500 MHz):  $\delta$  5.17 (1H, d,  $J_{1a,2a}$  1.8, H-1a), 5.11 (1H, d,  $J_{1c,2c}$  2.2, H-1c), 4.93 (1H, d,  $J_{1b,2b}$  1.8, H-1b), 4.13–4.08 (3H, m, H-2c, H-3c, H-4c), 4.05–3.93 (5H, m, H-2a, H-3a, H-5a, H-6a, H-2b), 3.92–3.81 (4H, m, H-4a, H-3b, H-6b, H-5c), 3.80–3.71 (2H, m, H-6'a, H-6'b), 3.71–3.63 (4H, m, H-4b, H-5b, H-6c).  $^{13}\text{C}$  NMR ( $\text{D}_2\text{O}$ , 126 MHz):  $\delta$  107.77 (C-1c), 100.34 (C-1b), 93.88 (C-1a), 82.66 (C-2c), 81.04 (C-4c), 76.06 (C-3c), 75.25 (C-4a), 72.92 (C-5b), 70.50, 70.49 (C-5c, C-2a), 70.44 (C-3b), 69.86 (C-5a), 69.65 (C-2b), 68.76 (C-3a), 66.62 (C-4b), 66.15 (C-6a), 62.68 (C-6c), 60.84 (C-6b).



**3aβ**: <sup>1</sup>H NMR (D<sub>2</sub>O, 500 MHz): δ 5.11 (1H, d, *J*<sub>1c,2c</sub> 2.2, H-1c), 4.94 (1H, d, *J*<sub>1b,2b</sub> 1.8, H-1b), 4.90 (1H, s, H-1a), 4.13–4.08 (3H, m, H-2c, H-3c, H-4c), 4.05–3.93 (4H, m, H-2a, H-3a, H-6a, H-2b), 3.92–3.81 (3H, m, H-3b, H-6b, H-5c), 3.80–3.71 (3H, m, H-4a, H-6'a, H-6'b), 3.71–3.63 (4H, m, H-4b, H-5b, H-6c), 3.64–3.59 (1H, m, H-5a). <sup>13</sup>C NMR (D<sub>2</sub>O, 126 MHz): δ 107.73 (C-1c), 100.30 (C-1b), 93.77 (C-1a), 82.75 (C-2c), 81.00 (C-4c), 76.06 (C-3c), 74.95 (C-4a), 73.38 (C-5a), 72.92 (C-5b), 70.50, 70.49 (C-5c, C-2a), 70.44 (C-3b), 69.65 (C-2b), 68.76 (C-3a), 66.62 (C-4b), 66.06 (C-6a), 62.68 (C-6c), 60.87 (C-6b).

HRMS (ESI): calcd for C<sub>18</sub>H<sub>32</sub>O<sub>16</sub>Na [M + Na]<sup>+</sup> 527.1588, found 527.1590.

### [β-D-Galactopyranosyl-(1 → 4)]-α-D-mannopyranosyl-(1 → 6)-α,β-D-mannopyranose (3b)

To a solution of **23b** (33 mg, 0.06 mmol) in a CH<sub>3</sub>CN/H<sub>2</sub>O 9 : 1 mixture (5 mL), *N*-iodosaccharin (26 mg, 0.08 mmol) was added. The reaction mixture was stirred at RT for 2 h. After the disappearance of the starting compound, the reaction was quenched by adding triethylamine and concentrated under reduced pressure. The resulting crude was partitioned between CH<sub>2</sub>Cl<sub>2</sub> and water, and the aqueous layer was washed with CH<sub>2</sub>Cl<sub>2</sub> (3×). After freeze-drying of the aqueous phase, the target trisaccharide **3b** was chromatographically purified on silica gel (AcOEt/MeOH/H<sub>2</sub>O gradient from 8 : 1 : 1 up to 6 : 2 : 2) and isolated (25 mg, 83%) as a white solid.

**3bα**: <sup>1</sup>H NMR (D<sub>2</sub>O, 500 MHz): δ 5.17 (1H, s, H-1a), 4.93 (1H, s, H-1b), 4.44 (1H, d, *J*<sub>1c,2c</sub> 8.1, H-1c), 4.10–3.97 (4H, m, H-2a, H-5a, H-6a, H-2b), 3.96–3.87 (4H, m, H-3a, H-4a, H-6b, H-4c), 3.87–3.70 (7H, m, H-6'a, H-3b, H-5b, H-6'b, H-5c, H-6c), 3.70–3.64 (2H, m, H-4b, H-3c), 3.58–3.51 (1H, m, H-2c).

<sup>13</sup>C NMR (D<sub>2</sub>O, 126 MHz): δ 102.99 (C-1c), 100.22 (C-1b), 93.80 (C-1a), 76.32 (C-4a), 75.49 (C-5c), 72.89 (C-5b), 72.45 (C-3c), 70.89 (C-2c), 70.52 (C-3b), 70.03 (C-2b), 69.84 (C-2a), 69.62 (C-5a), 68.92 (C-3a), 68.61 (C-4c), 66.61 (C-4b), 65.99 (C-6a), 61.13 (C-6c), 60.95 (C-6b).

**3bβ**: <sup>1</sup>H NMR (D<sub>2</sub>O, 500 MHz): δ 4.94 (1H, s, H-1b), 4.91 (1H, s, H-1a), 4.44 (1H, d, *J*<sub>1c,2c</sub> 8.1, H-1c), 4.10–3.97 (3H, m, H-2a, H-6a, H-2b), 3.96–3.87 (3H, m, H-3a, H-6b, H-4c), 3.87–3.70 (8H, m, H-4a, H-6'a, H-3b, H-5b, H-6'b, H-5c, H-6c), 3.70–3.64 (2H, m, H-4b, H-3c), 3.58–3.50 (2H, m, H-5a, H-2c). <sup>13</sup>C NMR (D<sub>2</sub>O, 126 MHz): δ 102.99 (C-1c), 100.17 (C-1b), 93.67 (C-1a), 75.92 (C-4a), 75.49 (C-5c), 73.45 (C-5a), 72.84 (C-5b), 72.45 (C-3c), 70.90 (C-2c), 70.50 (C-3b), 70.03 (C-2b), 69.81 (C-2a), 68.92 (C-3a), 68.61 (C-4c), 66.65 (C-4b), 65.91 (C-6a), 61.13 (C-6c), 60.95 (C-6b).

HRMS (ESI): calcd for C<sub>18</sub>H<sub>32</sub>O<sub>16</sub>Na [M + Na]<sup>+</sup> 527.1588, found 527.1586.

### General procedure for lichen extraction

All lichen samples were collected in France. *Cetraria islandica*, *Lasallia pustulata* and *Roccella fuciformis* were collected respectively in the Alpes (Haute-Savoie (74), August 2009), in Sainte Urcize (Cantal (15), June 2005) and near the plage des Chevrets (St Malo, Ille et Vilaine (35), June 2005). Vouchers were stored

in the Herbarium of the lab under the codes (JB/09/106), (JB/19/227) and (JB/05/58), respectively.

According to Pereyra,<sup>10</sup> the thalli of lichens were separated from the support (soil and rock) debris, washed with water (3×) and dried overnight at 40 °C. 15 g of thalli were boiled for 30 min at 100 °C (3×) and filtered through a cheesecloth. After cooling, the soluble fraction was centrifuged (2000g, 15 min). The supernatant was precipitated with EtOH (v/v) and filtered through a Büchner filter. Fractions A2 (>10 kDa) were obtained after ultrafiltration using an Amicon device with a regenerated cellulose filter (10 kDa, Merck Milipore), a concentration step under vacuum and a lyophilization step. They were stored in the dark at room temperature. The insoluble fraction after hot water extraction was treated with NaOH 1 M (3 × 200 mL) at 25 °C followed by centrifugation (2000g, 15 min). The supernatant was precipitated with EtOH (v/v) and filtered through a Büchner filter. Fractions F1S (>10 kDa) were obtained after ultrafiltration using the Amicon device with a regenerated cellulose filter (10 kDa, Merck Milipore), a concentration step under vacuum and a lyophilization step.

## Author contributions

V. F., L. L., I. C. and D. R. contributed to the conceptualisation and acquisition of the financial support. B. F., L.-P. D. and L. L. synthesized all oligosaccharides and intermediates. L. L., S. F. and F. L. D. participated in the interpretation of the 2D HSQC maps of the lichen extracts. L. L. and V. F. shared the preparation of the initial draft. All contributors participated in the approval of the final version of the manuscript.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

Financial support from Agence Nationale de la Recherche (project ALGAIMS ANR-18-CE29-0006-02, <https://algaims-35.websself.net/accueil>) is gratefully acknowledged. Part of this work was performed using the PRISM core facility (Biogenouest, Univ Rennes, Univ Angers, INRAE, CNRS, France). Special thanks to Joel Esnault for the pictures of the lichens.

## References

- 1 D. L. Hawksworth and M. Grube, *New Phytol.*, 2020, **227**, 1281–1283.
- 2 K. Müller, *Appl. Microbiol. Biotechnol.*, 2001, **56**, 9–16.
- 3 P. Le Pogam, G. Herbette and J. Boustie, in *Recent Advances in Lichenology*, ed. D. K. Upreti, P. K. Divakar, V. Shukla and R. Bajpai, Springer India, 2015.



- 4 F. Gasulla, E. M. del Campo, L. M. Casano and A. Guéra, *Plants*, 2021, **10**, 807.
- 5 T. Spribille, G. Tagirdzhanova, S. Goyette, V. Tuovinen, R. Case and W. F. Zandberg, *FEMS Microbiol. Lett.*, 2020, **367**, fnaa023.
- 6 M. González-Hourcade, E. M. del Campo, M. R. Braga, A. Salgado and L. M. Casano, *Environ. Microbiol.*, 2020, **22**, 3096–3111.
- 7 P. A. J. Gorin and E. Barreto-Bergter, in *The Polysaccharides*, ed. G. O. Aspinall, Academic Press, 1983, pp. 365–409, DOI: [10.1016/B978-0-12-065602-8.50011-X](https://doi.org/10.1016/B978-0-12-065602-8.50011-X).
- 8 E. S. Olafsdottir and K. Ingólfssdottir, *Planta Med.*, 2001, **67**, 199–208.
- 9 P. A. J. Gorin and M. Iacomini, *Carbohydr. Res.*, 1984, **128**, 119–132.
- 10 M. T. Pereyra, A. Prieto, M. Bernabé and J. A. Leal, *Lichenologist*, 2003, **35**, 177–185.
- 11 E. R. Carbonero, F. R. Smiderle, A. H. P. Gracher, C. G. Mellinger, G. Torri, T. Ahti, P. A. J. Gorin and M. Iacomini, *Carbohydr. Polym.*, 2006, **63**, 13–18.
- 12 P. A. J. Gorin and M. Iacomini, *Carbohydr. Res.*, 1985, **142**, 253–267.
- 13 Y. Sone, M. Isoda-Johmura and A. Misaki, *Biosci., Biotechnol., Biochem.*, 1996, **60**, 213–215.
- 14 E. R. Carbonero, L. M. C. Cordeiro, C. G. Mellinger, G. L. Sasaki, E. Stocker-Wörgötter, P. A. J. Gorin and M. Iacomini, *Carbohydr. Res.*, 2005, **340**, 1699–1705.
- 15 G. Shrestha, L. L. St. Clair and K. L. O'Neill, *Phytother. Res.*, 2015, **29**, 317–322.
- 16 E. S. Olafsdottir, S. Omarsdottir, B. S. Paulsen, K. Jurcic and H. Wagner, *Phytomedicine*, 1999, **6**, 273–279.
- 17 K. Ingólfssdottir, K. Jurcic, B. Fischer and H. Wagner, *Planta Med.*, 1994, **60**, 527–531.
- 18 L. D. Hall and L. F. Johnson, *J. Chem. Soc. D*, 1969, 509–510, DOI: [10.1039/C29690000509](https://doi.org/10.1039/C29690000509).
- 19 V. B. Krylov, D. A. Argunov, A. S. Solovev, M. I. Petruk, A. G. Gerbst, A. S. Dmitrenok, A. S. Shashkov, J. P. Latgé and N. E. Nifantiev, *Org. Biomol. Chem.*, 2018, **16**, 1188–1199.
- 20 B. Favreau, O. Yeni, S. Ollivier, J. Boustie, F. Le Dévéhat, J.-P. Guégan, M. Fanuel, H. Rogniaux, R. Brédy, I. Compagnon, D. Ropartz, L. Legentil and V. Ferrières, *J. Org. Chem.*, 2021, **86**, 6390–6405.
- 21 O. Yeni, S. Ollivier, B. Moge, D. Ropartz, H. Rogniaux, L. Legentil, V. Ferrières and I. Compagnon, *J. Am. Chem. Soc.*, 2023, **145**, 15180–15187.
- 22 S. Ollivier, L. Legentil, O. Yeni, L.-P. David, V. Ferrières, I. Compagnon, H. Rogniaux and D. Ropartz, *J. Am. Soc. Mass Spectrom.*, 2023, **34**, 627–639.

