Volume 22 Number 35 21 September 2024 Pages 7063-7246

Organic & **Biomolecular Chemistry**

rsc.li/obc

ISSN 1477-0520

PAPER Patrick J. Guiry *et al.* $Zn(u)$ -catalyzed asymmetric $[3 + 2]$ cycloaddition of acyclic enones with azomethine ylides

Biomolect Biomolecular Chemistry

PAPER

Cite this: Org. Biomol. Chem., 2024 22, 7148

Received 22nd May 2024, Accepted 13th June 2024 DOI: 10.1039/d4ob00854e

rsc.li/obc

Introduction

The pyrrolidine ring is a pivotal core structure found in numerous natural products and bioactive compounds.¹ It is considered a privileged scaffold in pharmaceutical and medicinal chemistry, serving as a foundation for the development of novel classes of bioactive and medicinally important drugs.² Consequently various synthetic routes to assembling such types of frameworks have been developed. 3 Of all of the routes developed, the transition metal-catalyzed $\begin{bmatrix} 3 & 2 \end{bmatrix}$ asymmetric cycloaddition reaction stands out due to its convergent and atom-economical approach for the construction of pyrrolidines with more than one chiral center.⁴ Various classes of dipolarophiles, such as α , β -unsaturated esters, maleimides, α,β-unsaturated nitriles, acyclic/cyclic enones, nitroalkenes, vinyl sulfones, and fullerenes, have been extensively studied. The reactivity and selectivities in these reactions are significantly influenced by the steric and electronic properties of both the dipolarophiles and dipoles, as well as the Lewis acidity of the metal-ligand complex utilized. 4 More recently, dipolarophiles such as cyclobutanone,⁵ α -halonitroalkenes,⁶ and electron-rich benzofulvenes⁷ have also been explored. Among these, acyclic/cyclic enones have garnered significant attention due to their synthetic potential. $8-14$ However, achieving high endo/exo ratios and enantioselectivities with these substrates remains challenging. PAPER
 (A) Check for updates
 (A) Check for updates
 EVALUATE CONDUCT CO

However, Carretero reported that the $Cu(1)$ -Fesulphos complex catalyzed the $[3 + 2]$ azomethine ylide cycloaddition of acyclic/cyclic enones, with the endo/exo selectivity being highly dependent on the *cis* or *trans* geometry of the enones.⁸ In contrast, use of Cu(I)-imFerroS or Ag-imFerroS produced

$Zn(i)$ -catalyzed asymmetric $[3 + 2]$ cycloaddition of acyclic enones with azomethine ylides†

Sundaravel Vivek Kumar, Jeremiah Olusegun and Patrick J. Guiry \mathbb{D}^*

The Zn(II)/UCD-Imphanol-catalyzed highly endo-selective [3 + 2] asymmetric cycloaddition of acyclic enones and azomethine ylides has been developed. Moderate to high yields (up to 94%) with excellent endo/exo selectivities (99 : 1) and enantioselectivities up to 96.5 : 3.5 er were obtained.

> endo-cycloadducts from azomethine ylide cycloaddition of cyclic/acyclic enones, respectively.⁹ Similarly, the Ag(1)-ThioClickFerrophos catalytic system afforded high levels of endo-selectivity in cycloadduct formation, irrespective of the cis or *trans* geometry of the enones.¹⁰ The application of the Ag(I)-Xingphos catalytic system demonstrated solvent-dependent selectivities.¹¹

> Nájera observed metal-dependent diastereoselectivity in the application of the Cu(I)-DM-SegPhos system in the formation of an exo-cycloadduct with 70% ee, whereas the Ag(I)-SegPhos system yielded an endo-cycloadduct with 88% ee.¹² Oh reported a metal-dependent enantiodivergent $\begin{bmatrix} 3 & + & 2 \end{bmatrix}$ cycloaddition, where the Cu (i) -brucine diol system provided endoselective products with good yields and enantioselectivities, compared to the Ag(i)-brucine diol system.¹³ Additionally, the $Au(I)$ - (S_2) -Binap catalytic system was reported to deliver good yields of endo-cycloadducts with 60–80% ee, compared to the Ag(I)-(S_a)-Binap catalytic system.¹⁴

> Both $Cu(i)$ and Ag (i) -based catalysts have been extensively explored, often offering complementary results in terms of endo/exo selectivity.⁴ However, this trend becomes variable when acyclic enones are used as dipolarophiles. $8-14$ These procedures frequently encounter one or more limitations, such as modest diastereo- or enantioselectivities and/or restricted substrate scope.

> On the other hand, despite its unique advantages such as $cost$ -effectiveness, abundance $(24th$ most abundant element), non-toxicity, and environmental friendliness, the use of zinc catalysis in azomethine ylide cycloadditions is rather limited.¹⁵ Recently, our group developed the UCD-Imphanol family of ligands, which proved highly efficient for Zn(II)-catalyzed [3 + 2] asymmetric cycloaddition of azomethine ylides with maleimides. These $Zn(n)$ -UCD-Imphanol catalytic system produced endo-selective cycloadducts in good yields, achieving enantioselectivities of up to 99.9% ee.^{15c} In continuation of our work, we present a highly *endo-selective* asymmetric $[3 + 2]$ cycloaddition of azomethine ylides with chalcones catalyzed by $Zn(n)/UCD$ -Imphanol (Scheme 1).

Centre for Synthesis and Chemical Biology, School of Chemistry, University College Dublin, Belfield, Dublin 4, Ireland. E-mail: p.guiry@ucd.ie

[†]Electronic supplementary information (ESI) available. CCDC 2299375. For ESI and crystallographic data in CIF or other electronic format see DOI: [https://doi.](https://doi.org/10.1039/d4ob00854e) [org/10.1039/d4ob00854e](https://doi.org/10.1039/d4ob00854e)

Scheme 1 Asymmetric $[3 + 2]$ cycloaddition of azomethine ylides with acyclic enones.

Results and discussion

We began our investigation by using (E) -chalcone 1a, ethyl (E) -2-((4-chlorobenzylidene)amino)acetate 2a as model substrates employing DABCO as base, $\text{Zn}(\text{OTf})_2$ as the metal catalyst to screen the series of UCD-Imphanol family of ligands including the non-planar chiral N,O-ligand 6. The use of ligand (S, S, S_P) -

4a gave the cycloadduct 3a in 55% yield with a 93:7 endo/exo ratio and $95:5$ er (Table 1, entry 1), whereas the use of 4-methylbenzyl- or pentafluorobenzyl-substituted ligands (S,S, $S_{\rm P}$)-4b and $(S, S, S_{\rm P})$ -4c led to the cycloadduct 3a in slightly reduced yield and selectivity (Table 1, entries 2 and 3). Use of ligand 4d afforded the cycloadduct 3a in 79% yield with a 94 : 6 diastereoselectivity and 94.5 : 5.5 er (Table 1, entry 4). In contrast, the diastereomeric ligands (S, S, R_P) -5a–c showed poor levels of reactivity and selectivity (Table 1, entries 5–7). The non-planar chiral N,O-ligand 6 gave a 38% yield with moderate diastereoselectivity (78 : 12) but poor enantioselectivity (54 : 46 er) (Table 1, entry 8), which indicated that the planar chiral paracyclophane unit was necessary to achieve high levels of reactivity and selectivity. The metal salts AgOAc and $Cu(OTf)_{2}$ were examined; the former produced the cycloadduct 3a in 58% yield with high diastereoselectivity and a 55 : 45 er, while the latter yielded only a trace amount of product (for full optimization studies see the ESI†). The screening of base and solvent revealed that dichloromethane and DABCO were optimal choices of solvent and base, respectively (see ESI†). Organic & Biomolecular Chemistry

Properties

Published on 14 Six articles. Published on 14 Six articles. The published on 14 Six articles. The published under the use of the state of the state of the state of the state o

Gratifyingly, the use of an excess (2 equiv.) of 2a led to an increase in yield to 90% without affecting selectivity (Table 1, entry 9). Further, lowering the reaction temperature to 0 \degree C produced the cycloadduct 3a in 95% yield with an excellent

 a All reactions were carried out with Zn(OTf)2 (10 mol%), L (11.5 mol%), 1a (0.15 mmol), 2a (0.225 mmol), DABCO (10 mol%) in CH2Cl2 (1.5 mL) for 24 h. ^b Isolated yield after purification. ^c The *endo/exo* ratios were determined by analyzing the ¹H NMR spectra of the crude product.
^d Determined by chiral HPLC analysis. ^e 0.3 mmol of 2a was used. ^f 4 ^d Determined by chiral HPLC analysis. e 0.3 mmol of 2a was used. f 48 h. g Zn(OTf)₂ (5 mol%) and 4d (5.75 mol%) for 72 h.

endo/exo ratio and 96.5 : 3.5 er, our best result in this optimization study (Table 1, entry. 10). Further decreasing the catalyst loading (5 mol% of $Zn(OTf)₂$ and 5.75 mol% of 4d) led to long reaction time with slight reduction of yield and 99 : 1 endo/exo ratio and 96 : 4 er.

Having optimized conditions in hand, we then examined the scope of this reaction for a focused library of acyclic enones which afforded the products 3a–k with excellent levels of diastereocontrol (99 : 1) (Scheme 2). A series of chalcones with 4-OMe, 4-Cl, and 3-Br-substituted β-aryl groups gave the corresponding products 3b–c, 3e in moderate to good yields with enantioselectivities of 85 : 15–90 : 10 er. In contrast, the reaction of $4-NO₂$ -substituted chalcone led to a complicated mixture; the expected product 3d was not formed, which may presumably be due to the competitive coordination of the nitro group with the Zn metal. The sterically hindered 1-naphthyl and 3-thienyl derived substrates were also tolerated under these reaction conditions to yield the corresponding cycloadducts 3f and 3g in 94% and 50% yield with 96 : 4 and 91 : 9 er, respectively. However, the reaction with β-methyl-substituted chalcone afforded the cycloadduct 3h in a lower yield (35%) with a lower level of enantioselectivity (87 : 13). Next, various α-benzoyl-substituted chalcones also reacted to deliver products 3i–j in moderate to good yields with 88 : 12 and 87 : 13 er's. The use of (E) -benzylidene acetone gave rise to the cycloadduct 3k in a disappointing yield of 21% with 82 : 18 er. Paper

Paper Constrained on 2022. The strained of the comparison of the compa

Next, our attention turned to exploring the scope of the α-iminoesters (Scheme 3). We were pleased to find that a wide range of α-iminoesters with varying electronics and sterics were tolerated to afford the desired products 3l–w in 20–81% yields with 72 : 28–94 : 6 enantioselectivities and 99 : 1 endo/exo selectivities. In particular, α -iminoesters with electrondeficient and neutral substituents on the aryl ring led to the products 3m–p in 84–71% yields with enantioselectivities of 91 : 9–94 : 6 er. The α-iminoesters with sterically hindered substituents such as o-tolyl, 2-naphthyl, and ferrocenyl underwent the reaction to give the products 3q, 3s–t in 20–54% yields with 84 : 16–90 : 10 er's. Notably, the 1-naphthyl-substituted α-iminoester provided the cycloadduct 3r with the opposite stereochemistry (12 : 88 er) in 72% yield. Furthermore, 2-thienyl- and cyclohexyl-substituted iminoesters also showed poor reactivity (28–37% yield) for the formation of products 3u–v with enantioselectivities in the range 72 : 28–82 : 18 er's. Moreover, the α -methyl-substituted iminoester also displayed moderate reactivity, forming the product 3w in 48% yield with 75 : 25 er.

To demonstrate the synthetic potential of the present reaction, we performed the reaction on a 1 mmol scale, furnishing 3a in 86% yield and excellent diastereo- and enantioselectivity (99 : 1 endo/exo, 96.5 : 3.5 er) (Scheme 4a). Further treatment of the product with chloroacetyl chloride led to the formation of 7, which then reacted with 40% aq. MeNH_2 ,¹⁷ affording the diketopiperazine derivative 8 in 80% yield with 96 : 4 er (Scheme 4b).

A plausible transition state was tentatively proposed based on the Zn-complex 9, which was prepared from the reaction of ligand 4a with Et₂Zn, affording the homoleptic zinc (n) complex 9 (Scheme 5). The homoleptic complex was confirmed by single crystal X-ray diffraction studies (Fig. 1), and this complex would generate the heteroleptic $Zn(n)$ complex by interaction with iminoester 2 under basic conditions that can readily react with acyclic enones.

The transition state contains the azomethine ylide coordinating with the $Zn(\mu)$ -imidazolinyl-paracyclophanol in a tetrahedral complex (Fig. 2). The sterically crowded paracyclophane and C-4 phenyl group effectively shield the top-face approach to the Si-face of the ylide, while the Re-face is less crowded thus allowing the acyclic enone approach in an endo fashion (TSI) to deliver the products with good to moderate levels of enantioselectivity. Although it cannot be completely ruled out, the exceptional endo-selectivity that was observed may perhaps be due to the additional coordination of the α ,β-unsaturated ketone carbonyl group with the metal center. Additionally,

Fig. 1 Single crystal XRD analysis for 9. All H-atoms are omitted for clarity.

Fig. 2 Proposed stereochemical models.

TSII may explain the noted opposite enantiomer in the case of 1-naphthyl iminoester since TSI presumably leads to a disfavored steric clash between the 1-naphthyl iminoester and the paracyclophane unit of ligand.

Conclusions

In summary, we have developed the $Zn(\pi)$ -catalyzed highly endo-selective asymmetric $\begin{bmatrix} 3 & + & 2 \end{bmatrix}$ cycloaddition of acyclic enones with azomethine ylides by the use of the planar chiral UCD-Imphanol family of ligands, leading to chiral pyrrolidines in moderate to good yields with $99:1$ endo/exo selectivities and enantioselectivities up to 96.5 : 3.5 er. A variety of acyclic enones and iminoester-derived azomethine ylides are compatible with the less expensive Zn-catalysis. Further detailed mechanistic studies with computational analysis are currently underway and will be reported in due course.

Experimental section

General procedure for the Zn(π)-catalyzed [3 + 2] azomethine ylide cycloaddition

Under nitrogen atmosphere, ligand (S, S, S_p) -4d $(11.5 \text{ mol%)}$ and $Zn(OTf)_{2}$ (10 mol%) was added to a flame-dried 10 mL Schlenk tube. Then dry CH_2Cl_2 (0.5 mL) was added and stirred

for 60 min. After cooling to 0 \degree C, acyclic enone 1 (0.15 mmol, 1.0 equiv.) in CH_2Cl_2 (0.4 mL), α-iminoester 2 (0.3 mmol, 2 equiv.) in CH₂Cl₂, (0.4 mL) and DABCO (10 mol%) in CH₂Cl₂ (0.2 mL) were added sequentially. The reaction mixture was stirred at the same temperature for 48 h. After completion, the reaction mixture was directly purified by silica gel column chromatography (30% ethyl acetate in cyclohexane) to afford the desired product 3. The endo/exo ratio was determined by ¹H-NMR before purification and the enantioselectivity was determined by HPLC analysis of the isolated product. The configuration of the products was assigned by comparison with the optical rotation values of prepared compounds and the reported optical rotation of the cycloadducts.^{12a,16} (Note: the reaction temperature (0 °C) was maintained using a cryocooler). Paper

For 80 min. After colling to 0 "C, accelie enone 1 (0.15 mmol,

14 quay) in Origin, (0.4 min) in Origin, 2024. This article is limit in Accelie and K. C. This article. The spin of \sqrt{R} Creative Commons are the s

Data availability

The data supporting this article have been included as part of the ESI.† CDC 2299375 contains the supplementary crystallographic data for Zinc complex 9.†

Author contributions

PG supervised the research; S. V. K. and P. G. conceived the idea; S. V. K. and J. O. carried out all the synthetic experimental work and data acquisition. S. V. K. prepared the manuscript with contributions from J. O. and P. G. All authors approved the manuscript.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

S. V. K. thanks the Irish Research Council (IRC) for the award of a Postdoctoral Scholarship (GOIPD/2019/918) and UCD School of Chemistry for the award of a Teaching Fellowship. We acknowledge the facilities provided by the Centre for Synthesis and Chemical Biology (CSCB), funded by the Higher Education Authority's Programme for Research in Third-Level Institutions. We thank Dr Yannick Ortin, Dr Patricia Fleming and Dr Guanghua Jin for NMR spectroscopy, and Dr Jimmy Muldoon for mass spectrometric analysis (supported by a Science Foundation Ireland Infrastructure Award (18/RI/5702)).

References

1 (a) W. J. Olivier, J. A. Smith and A. C. Bissember, Chem. Rec., 2022, 22, e202100277; (b) M. T. Islam and M. S. Mubarak, Adv. Tradit. Med., 2020, 20, 13;

- (c) A. K. Chattopadhyay and S. Hanessian, Chem. Rev., 2017, 117, 4104; (d) C. Bhat and S. G. Tilve, RSC Adv., 2014, 4, 5405; (e) J. P. Michael, Nat. Prod. Rep., 2008, 25, 139; (f) D. O'Hagan, Nat. Prod. Rep., 2000, 17, 435.
- 2 (a) S. Poyraz, H. A. Döndaş, N. Y. Döndaş and J. M. Sansano, Front. Pharmacol., 2023, 14, 1239658; (b) G. L. Petri, M. V. Raimondi, V. Spanò, R. Holl, P. Barraja and A. Montalbano, Top. Curr. Chem., 2021, 379, 34.
- 3 Selected examples see;(a) R. Zhang, M. Sun, Q. Yan, X. Lin, X. Li, X. Fang, H. H. Y. Sung, I. D. Williams and J. Sun, Org. Lett., 2022, 24, 2359; (b) J.- B. Qiao, Y.-Q. Zhang, Q.-W. Yao, Z.-Z. Zhao, X. Peng and X.-Z. Shu, J. Am. Chem. Soc., 2021, 143, 12961; (c) Y.-F. Yu, C. Shu, T.-D. Tan, L. Li, S. Rafique and L.-W. Ye, Org. Lett., 2016, 18, 5178; (d) L. Liu, M. Leutzsch, Y. Zheng, M. W. Alachraf, W. Thiel and B. List, J. Am. Chem. Soc., 2015, 137, 13268; (e) G.-H. Hou, J.-H. Xie, P.-C. Yan and Q.-L. Zhou, J. Am. Chem. Soc., 2009, 131, 1366; (f) R. L. LaLonde, B. D. Sherry, E. J. Kang and F. D. Toste, J. Am. Chem. Soc., 2007, 129, 2452.
- 4 Selected recent examples $see; (a)$ S. V. Kumar and P. J. Guiry, Chem. – Eur. J., 2023, 29, e202300296; (b) J. Adrio and J. C. Carretero, Chem. Commun., 2019, 55, 11979; (c) X. Fang and C.-J. Wang, Org. Biomol. Chem., 2018, 16 , 2591; (d) T. Hashimoto and K. Maruoka, Chem. Rev., 2015, 115, 5366; (e) J. Adrio and J. C. Carretero, Chem. Commun., 2014, 50, 12434.
- 5 J. Corpas, A. Ponce, J. Adrio and J. C. Carretero, Org. Lett., 2018, 20, 3179.
- 6 V. A. Motornov, A. A. Tabolin, Y. V. Nelyubina, V. G. Nenajdenko and S. L. Ioffe, Org. Biomol. Chem., 2021, 19, 3413.
- 7 X. Chang, X.-T. Liu, F. Li, Y. Yang, L. W. Chung and C.-J. Wang, Chem. Sci., 2023, 14, 5460.
- 8 J. Hernández-Toribio, R. G. Arrayás, B. Martín-Matute and J. C. Carretero, Org. Lett., 2009, 11, 393.
- 9 C. Zhang, S.-B. Yu, X.-P. Hu, D.-Y. Wang and Z. Zheng, Org. Lett., 2010, 12, 5542.
- 10 I. Oura, K. Shimizu, K. Ogata and S. Fukuzawa, Org. Lett., 2010, 12, 1752.
- 11 X.-F. Bai, Z. Xu, C.-G. Xia, Z.-J. Zheng and L. W. Xu, ACS Catal., 2015, 5, 6016.
- 12 (a) G. S. Caleffi, O. Larrañaga, M. Martín-Rodríguez, P. R. R. Costa, C. Nájera, A. Cózar, F. P. Cossío and J. M. Sansano, J. Org. Chem., 2019, 84, 10593; (b) A. Cayuelas, O. Larrañaga, V. Selva, C. Nájera, T. Akiyama, J. M. Sansano, A. Cózar, J. I. Miranda and F. P. Cossío, Chem. – Eur. J., 2018, 24, 8092; (c) C. Nájera, M. G. Retamosa, M. Martín-Rodríguez, J. M. Sansano, A. Cózar and F. P. Cossío, Eur. J. Org. Chem., 2009, 5622.
- 13 J.-Y. Li, H. Y. Kim and K. Oh, Adv. Synth. Catal., 2016, 358, 984.
- 14 M. Martín-Rodríguez, C. Nájera, J. M. Sansano, A. Cózar and F. P. Cossío, Chem. – Eur. J., 2011, 17, 14224.
- 15 Zn-catalyzed iminoester-derived $[3 + 2]$ azomethine ylide cycloaddition see; (a) A. S. Gothelf, K. V. Gothelf, R. G. Hazell and K. A. Jørgensen, Angew. Chem., Int. Ed.,

2002, 41, 4236; (b) O. Dogan, H. Koyuncu, P. Garner, A. Bulut, W. J. Youngs and M. Panzner, Org. Lett., 2006, 8, 4687; For Zn-catalyzed [3+2] cycloadditions with other azomethine ylides see; (c) S. V. Kumar and P. J. Guiry, Angew. Chem., 2022, 134, e202205516; (d) Y. Yi, Y. Z. Hua, H.-J. Lu, L.-T. Liu and M.-C. Wang, Org. Lett., 2020, 22, 2527; (e) R.-L. Wang, S.-K. Jia, Y.-J. Guo, Y. Yi, Y.-Z. Hua, H.-J. Lu Organic & Blomolecular Chemistry

2002, 11, 2236; (*i)* 0 . Dogan, H. Koyuncu, P. Game;

A. This article. Published on 16 June 2024. Downloaded on 5.000. A. (*i) P. Zhun, N. Y*, Y. Hun, N. J. Hund M. Wome Commons Attributi

and M.-C. Wang, Org. Biomol. Chem., 2021, 19, 8492; (f) P. Zhou, Y. Yi, Y. Hua, S. Jia and M. Wang, Chem. - Eur. J., 2022, 28, e202103688.

- 16 Y. Zhang, J. Su, T. Lin, Z. Lin, Z. Zhou and Y. Li, J. Chem. Res., 2019, 43, 90.
- 17 E. García-Mingüens, V. Selva, O. Larrañaga, C. Nájera, J. M. Sansano and A. Cózar, ChemCatChem, 2020, 12, 2014.