



Cite this: *Org. Biomol. Chem.*, 2024, **22**, 8811

Received 20th August 2024,
Accepted 2nd October 2024

DOI: 10.1039/d4ob01373e

rsc.li/obc

Syntheses of bottromycin derivatives *via* Ugi-reactions and Matteson homologations†

Etienne Bickel and Uli Kazmaier *

New bottromycin derivatives have been prepared using flexible Ugi and Matteson reactions. The Ugi reaction allows the fast and direct assembly of sterically hindered peptide fragments, while the Matteson homologation is excellently suited for the stereoselective synthesis of unusual amino acids like β -methylphenylalanine. Some of the new compounds show excellent activity against *Streptococcus pneumoniae*.

Introduction

The rise of antimicrobial resistance (AMR) causes severe healthcare problems and, based on urgency and the need for new antibiotics, multidrug-resistant pathogens are classified as the highest priority by the World Health-Organisation (WHO).¹ Such pathogens can develop resistance towards almost all currently used antibiotics, so the development of new antimicrobial agents, with new modes of action, is highly desired.²

In 1957, Waisvisz *et al.* reported the isolation of a new peptide from the fermentation broth of *Streptomyces bottropensis* and called it bottromycin.³ Biosynthetically, bottromycins are formed from ribosomally synthesized peptides *via* post-translationally modifications.⁴ Mode of action (MoA) studies indicated that bottromycin inhibits protein biosynthesis by binding to the aminoacyl-tRNA binding site (A site) of the 50S ribosome.⁵ This biological target would avoid the cross-resistance issue but, to date, has not been addressed by any other antibiotic. Therefore, bottromycin is effective against problematic bacterial strains such as vancomycin-resistant *Enterococci* (VRE) and methicillin-resistant *Staphylococcus aureus* (MRSA).^{4b}

Although, the bottromycins were discovered and described as early as the late 1950's, only one total synthesis of bottromycin A₂ and analogues by Ōmura and Sunazuka *et al.* has been reported.⁶ A reason for this may be several incorrect structures proposed during these early years.⁷ The latest, and correct, proposal, as confirmed by the first total synthesis, was described by Shipper in 1983 (Fig. 1).⁸

Several research groups,⁹ including our own,¹⁰ have carried out synthetic studies towards the various bottromycin building blocks and proposed structures. Researchers at AiCuris GmbH (Germany) saponified the C-terminal methyl ester of biotechnologically produced bottromycin and converted the acid into several amides.¹¹ Further modifications on this C-terminal unusual amino acid were carried out by Ōmura and Sunazuka based on their previously developed total synthesis.⁶ Structure–activity relation (SAR) studies on these different derivatives indicate that especially the β -substituted phenylalanine is required for high activity. Obviously, the methyl group on the phenylalanine influences the conformation of the side chain and probably of the whole molecule as indicated by ¹H-NMR.⁶ Interestingly, while the C-terminal thiazolyl amino acid ester is not required for activity, the free amino acid or its complete removal render the derivatives almost inactive. However, replacing the amino acid by a simple benzylamide had no significant effect on the activity. Clearly, only an amide bond is required, and an aromatic substituent has a positive effect on the activity.

In the context of our interest in the total synthesis of natural products with anticancer¹² or antibiotic¹³ activities, we were motivated to develop a flexible synthetic approach towards bottromycin derivatives (Fig. 2) which should allow us to readily modify the core structure of the molecule.




Fig. 1 Bottromycin A₂.

Institute of Organic Chemistry, Saarland University, P.O. Box 151150, 66041 Saarbrücken, Germany. E-mail: u.kazmaier@mx.uni-saarland.de

† Electronic supplementary information (ESI) available: Experimental details, compound characterization, copies of ¹H and ¹³C NMR spectra. See DOI: <https://doi.org/10.1039/d4ob01373e>






Fig. 2 Modification of bottromycin.

Results and discussion

For a flexible synthesis of the side chain, we investigated two different approaches. First, a synthesis of the β -branched phenylalanine should give us the freedom to modify this amino acid. Because β -methylphenylalanine (MePhe) is incorporated not only in bottromycin but also several other natural products,¹⁴ various syntheses for this building block already exist, although many are specific for this amino acid. Therefore, we decided to apply the Matteson homologation¹⁵ which should give us a versatile and stereoselective means to modify every carbon atom of the amino acid side chain.


Starting from the known methyl boronic ester **1**,¹⁵ addition of lithiated dichloromethane (DCM) (generated *in situ* by deprotonating DCM with LDA) provided the α -chloroboronic ester **2** in a highly stereoselective fashion (Scheme 1). Addition of phenylmagnesium bromide generated a boronate complex which underwent a 1,2-phenyl shift replacing the chlorine in

an S_N2 fashion. The desired α -phenyl substituted boronic ester **3** was obtained as a single stereoisomer. Repeating the sequence using vinylmagnesium bromide in the second step gave access to allylboronate **4**.


According to a protocol described by Morken, **4** was reacted with deprotonated *O*-methylhydroxylamine resulting in replacement of the boronic ester by an NH_2 group,¹⁶ which was directly Boc-protected (**5**). Subsequent ozonolysis and Pinnick oxidation¹⁷ gave the *N*-protected amino acid **6** which was converted into benzylamide **7**. Cleavage of the Boc protecting group, peptide coupling, and further *N*-deprotection gave the desired dipeptide **8**. In principle, by using other starting boronic esters or other aryl Grignard reagents, a wide range of derivatives should be accessible.

As a complementary and powerful protocol, we took advantage of the Ugi reaction¹⁸ to generate the entire side chain in only one or two steps. In principle, enantiomerically pure 2-phenylpropionaldehyde can be used as the aldehyde component in this 4-component coupling for direct access to the bottromycin side chain. However, we decided to use achiral 2-methyl-2-phenylpropionaldehyde, generating a quaternary β -carbon (Scheme 2). Ugi reaction with Boc-*t*-leucine (Boc-Tle), ammonia, and benzyl isocyanide in trifluoroethanol generated the desired dipeptide **9** in high yield as a mixture of diastereomers and rotamers. Purification by flash chromatography provided a 1:1 diastereomeric mixture in 60% yield, which was subjected to *N*-Boc deprotection. The diastereomers of dipeptide **10** with the free amino terminus could be separated by reversed-phase chromatography or crystallisation. The crystals obtained were suitable for X-ray structure analysis which allowed us to determine the absolute configuration of the two diastereomers.

The Ugi reaction was also used to generate the sterically demanding peptide fragment applying our previously developed thio-Ugi approach (Scheme 3).¹⁹ Alloc-protected valine was activated with carbonyldiimidazol (Im₂CO) to the corresponding imidazolide which was directly reacted with H₂S to afford the corresponding thioacid. Without purification, **11** was reacted with pivalaldehyde, methylamine, and the isocyanide obtained from glycine methyl ester. The reaction proceeded cleanly and the two diastereomers of **12** could be separated.




Scheme 1 Synthesis of dipeptide **8** via Matteson homologation. Abbreviations: NMM: *N*-methylmorpholine; HATU: [O-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium-hexafluorophosphat].



Scheme 2 Synthesis and separation of diastereomers of dipeptide **10**.





Scheme 5 Synthesis of bottromycin-derivatives **23**.

any significant activity ($\text{MIC} > 64 \mu\text{g mL}^{-1}$), either against the *Mycobacterium* strains or any other Gram-positive or Gram-negative bacteria or yeasts and fungi. In contrast, some activity was observed for the two derivatives **23** with the unmodified peptide ring (Table 1). Although the activity against Mtb was

Table 1 Biological activities of new bottromycins **23**

Bacterial strain	MIC ($\mu\text{g mL}^{-1}$)		
	BotA ₂	23a	23b
<i>M. tuberculosis</i> H37Ra	1	16	32
<i>M. smegmatis</i> mc ² 155	8–16	2	4
<i>E. faecalis</i> 20478		8	>64
<i>E. faecium</i> DSM-17050		32	>64
<i>E. faecium</i> 20477		2	16
<i>S. pneumoniae</i> DSM-11865		<0.03	0.5
<i>S. pneumoniae</i> 20566		<0.03	0.25

weaker than that of bottromycin A₂, a higher activity was observed against Msm. This led us to investigate the activity against other strains that had not yet been tested. Derivative **23a** with the natural β -methylphenylalanine was approximately 10-fold more active than the double β -methylated derivative **23b**. The compounds proved to be particularly active against *Streptococcus pneumoniae*.

Conclusions

In conclusion, we have shown that bottromycins are readily accessible when applying Ugi reactions to generate larger fragments. Unfortunately, in SAR studies, no significant structural changes are tolerated on the peptide ring. *N*-Methylation facilitates cyclisation but, unfortunately, does not provide active derivatives. Matteson homologation was found to be a suitable tool for the stereoselective synthesis of β -methylated phenylalanine.

Data availability

The data supporting this article (copies of ¹H, ¹³C NMR spectra and experimental details) have been included as part of the ESI.†

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Financial support from the Saarland University and the DFG (Ka 880/13-1 and Bruker Neo 500 - 447298507) is gratefully acknowledged.

References

- 1 E. Tacconelli, E. Carrara, A. Savoldi, S. Harbarth, M. Mendelson, D. L. Monnet, C. Pulcini, G. Kahlmeter, J. Kluytmans, Y. Carmeli, M. Ouellette, K. Outtersson,



- J. Patel, M. Cavaleri, E. M. Cox, C. R. Houchens, M. L. Grayson, P. Hansen, N. Singh, U. Theuretzbacher, N. Magrini and the WHO Pathogens Priority List Working Group, Discovery, research, and development of new antibiotics: the WHO priority list of antibiotic-resistant bacteria and tuberculosis, *Lancet Infect. Dis.*, 2018, **18**, 318–327.
- 2 (a) B. A. Cunha, Antibiotic resistance: A historical perspective, *Semin. Respir. Crit. Care Med.*, 2000, **21**, 3–8; (b) Z. Lin, T. Yuan, L. Zhou, S. Cheng, X. Qu, P. Lu and Q. Feng, Impact factors of the accumulation, migration and spread of antibiotic resistance in the environment, *Environ. Geochem. Health*, 2021, **43**, 1741–1758.
- 3 (a) J. M. Waisvisz, M. G. van der Hoeven, J. van Peppen and W. C. M. Zwennis, Bottromycin. I. A new sulfurcontaining antibiotic, *J. Am. Chem. Soc.*, 1957, **79**, 4520–4521; (b) J. M. Waisvisz, M. G. van der Hoeven, J. F. Hölscher and B. te Nijenhuis, Bottromycin. II. Preliminary degradation studies, *J. Am. Chem. Soc.*, 1957, **79**, 4522–4523; (c) J. M. Waisvisz, M. G. van der Hoeven and B. te Nijenhuis, The structure of the sulfur-containing moiety of bottromycin, *J. Am. Chem. Soc.*, 1957, **79**, 4524–4527.
- 4 (a) T. H. Eyles, N. M. Vion and A. W. Truman, Rapid and Robust Yeast-Mediated Pathway Refactoring Generates Multiple New Bottromycin-Related Metabolites, *ACS Synth. Biol.*, 2018, **7**, 1211–1218; (b) L. Franz, U. Kazmaier, A. W. Truman and J. Koehnke, Bottromycins - biosynthesis, synthesis and activity, *Nat. Prod. Rep.*, 2021, **38**, 1659–1683; (c) P. H. Krushnamurthy, K. S. Subramanya, D. Simita, G. Dhananjaya and M. Nilkamal, Recent Advancements in Bottromycin Biosynthesis, *Synlett*, 2023, 793–806.
- 5 (a) T. Otaka and A. Kaji, Mode of action of bottromycin A2. Release of aminoacyl or peptidyl tRNA from ribosomes, *J. Biol. Chem.*, 1976, **251**, 2299–2306; (b) T. Otaka and A. Kaji, Mode of action of bottromycin A2: Effect of bottromycin A2 on polysomes, *FEBS Lett.*, 1983, **153**, 53–59; (c) T. Otaka and A. Kaji, Mode of action of bottromycin A2: effect on peptide bond formation, *FEBS Lett.*, 1981, **123**, 173–176.
- 6 (a) H. Shimamura, H. Gouda, K. Nagai, T. Hirose, M. Ichioka, Y. Furuya, Y. Kobayashi, S. Hirono, T. Sunazuka and S. Ōmura, Structure determination and total synthesis of bottromycin A2: a potent antibiotic against MRSA and VRE, *Angew. Chem., Int. Ed.*, 2009, **48**, 914–917; (b) Y. Kobayashi, M. Ichioka, T. Hirose, K. Nagai, A. Matsumoto, H. Matsui, H. Hanaki, R. Masuma, Y. Takahashi, S. Ōmura and T. Sunazuka, Bottromycin derivatives: Efficient chemical modifications of the ester moiety and evaluation of anti-MRSA and anti-VRE activities, *Bioorg. Med. Chem. Lett.*, 2010, **20**, 6116–6120; (c) T. Yamada, M. Yagita, Y. Kobayashi, G. Sennari, H. Shimamura, H. Matsui, Y. Horimatsu, H. Hanaki, T. Hirose, S. Ōmura and T. Sunazuka, Synthesis and evaluation of antibacterial activity of bottromycins, *J. Org. Chem.*, 2018, **83**, 7135–7149.
- 7 (a) S. Nakamura, T. Chikaike, H. Yonehara and H. Umezawa, Structures of bottromycins A and B, *J. Antibiot., Ser. A*, 1965, **18**, 60–61; (b) S. Nakamura, N. Tanaka and H. Umezawa, Bottromycin A1, A2 and their structures, *J. Antibiot.*, 1966, **19**, 10–12; (c) Y. Takahashi, H. Naganawa, T. Takita, H. Umezawa and S. Nakamura, The revised structure of bottromycin A2, *J. Antibiot.*, 1976, **29**, 1120–1123.
- 8 D. Schipper, The revised structure of Bottromycin-A2, *J. Antibiot.*, 1983, **36**, 1076–1077.
- 9 U. Kazmaier, The long, long way to Bottromycin, *Isr. J. Chem.*, 2021, **61**, 308–321.
- 10 S. Ackermann, H.-G. Lerchen, D. Haebich, A. Ullrich and U. Kazmaier, Synthetic studies towards bottromycin, *Beilstein J. Org. Chem.*, 2012, **8**, 1652–1656.
- 11 H.-G. Lerchen, G. Schiffer, H. Broetz-Oesterhelt, A. Mayer-Bartschmid, S. Eckermann, C. Freiberg, R. Endermann, J. Schuhmacher, H. Meier, N. Svenstrup, S. Seip, M. Gehling and D. Haebich, Fermentative and chemical synthesis of bottromycin derivative analogs for use in treatment or prevention of bacterial infections in humans or animals, WO2006103010A1, 2006.
- 12 (a) A. Ullrich, Y. Chai, D. Pistorius, Y. A. Elnakady, J. E. Herrmann, K. J. Weissman, U. Kazmaier and R. Müller, Pretubulysin, a potent and chemically-accessible tubulysin precursor from *Angiococcus disciformis*, *Angew. Chem., Int. Ed.*, 2009, **47**, 4422–4425; (b) L. Karmann, K. Schulz, J. Herrmann, R. Müller and U. Kazmaier, Total syntheses and biological evaluation of miuraenamides, *Angew. Chem., Int. Ed.*, 2015, **54**, 4502–4507; (c) J. Gorges and U. Kazmaier, Matteson homologation-based total synthesis of lagunamide A, *Org. Lett.*, 2018, **20**, 2033–2036; (d) O. Andler and U. Kazmaier, Total synthesis of apratoxin A using Matteson's homologation approach, *Org. Biomol. Chem.*, 2021, **19**, 4866–4870.
- 13 (a) P. Barbie and U. Kazmaier, Total synthesis of cyclo- marin A, a marine cycloheptapeptide with anti-tuberculosis and anti-malaria activity, *Org. Lett.*, 2016, **18**, 204–207; (b) T. Kinsinger and U. Kazmaier, C-H-Functionalization of N-methylated amino acids and peptides as tool in natural product synthesis – Synthesis of abyssenine A and mucronine E, *Org. Lett.*, 2018, **20**, 7726–7730; (c) J. Greve, A. Mogk and U. Kazmaier, Total synthesis and biological evaluation of modified ilamycin derivatives, *Mar. Drugs*, 2022, **20**, 632; (d) R. Priester and U. Kazmaier, A straightforward synthesis of emericellamide A using Matteson's homologation approach, *Synlett*, 2023, 2159–2164; (e) O. Andler and U. Kazmaier, Matteson homologation-based total synthesis of meliponamycin A, *Org. Lett.*, 2024, **26**, 148–152.
- 14 (a) S. Fuse, H. Koinuma, A. Kimbara, M. Izumikawa, Y. Mifune, H. He, K. Shin-ya, T. Takahashi and T. Doi, Total synthesis and stereochemistry revision of mannopeptimycin aglycone, *J. Am. Chem. Soc.*, 2014, **136**, 12011–12017; (b) C. S. V. Houge-Frydrych, M. L. Gilpin, P. W. Skett and J. W. Tyler, SB-203207 and SB-203208, two novel isoleucyl tRNA synthetase inhibitors from a *Streptomyces sp.* II. Structure determination, *J. Antibiot.*, 2000, **53**, 364–372.
- 15 (a) D. S. Matteson, Boronic esters in stereodirected synthesis, *Tetrahedron*, 1989, **7**, 1859–1885; (b) D. S. Matteson, Boronic esters in asymmetric synthesis, *J. Org. Chem.*, 2013,



- 78, 10009–10023; (c) S. Kirupakaran, H.-S. Kort and C. Hirschhäuser, A complementary toolbox of iterative methods for the stereoselective synthesis of heteroatom-rich motives from C₁-building blocks, *Synthesis*, 2018, 2307–2322; (d) D. S. Matteson, B. S. L. Collins and V. K. Aggarwal, The Matteson Reaction, *Org. React.*, 2021, **105**, 427–860.
- 16 (a) S. N. Mlynarski, A. S. Karns and J. P. Morken, Direct stereospecific amination of alkyl and aryl pinacol boronates, *J. Am. Chem. Soc.*, 2012, **134**, 16449–16451; (b) E. K. Edelstein, A. C. Grote, M. D. Palkowitz and J. P. Morken, A protocol for direct stereospecific amination of primary, secondary, and tertiary alkylboronic esters, *Synlett*, 2018, 1749–1752; (c) P. Xu, M. Zhang, B. Ingoglia, C. Allais, A.-M. R. Dechert-Schmitt, R. A. Singer and J. P. Morken, Construction of azacycles by intramolecular amination of organoboronates and organobis (boronates), *Org. Lett.*, 2021, **23**, 3379–3383.
- 17 (a) B. O. Lindgren, T. Nilsson, S. Husebye, Ø. Mikalsen, K. Leander and C.-G. Swahn, Preparation of carboxylic acids from aldehydes (including hydroxylated benzaldehydes) by oxidation with chlorite, *Acta Chem. Scand.*, 1973, **27**, 888–890; (b) B. S. Bal, W. E. Childers and H. W. Pinnick, Oxidation of α,β -unsaturated aldehydes, *Tetrahedron*, 1981, **37**, 2091–2096.
- 18 (a) I. Ugi, *Isonitrile Chemistry*, Academic press, New York, 1971; (b) A. Dömling and I. Ugi, Multicomponent reactions with isocyanides, *Angew. Chem., Int. Ed.*, 2000, **39**, 3168–3210; (c) A. Ullrich and U. Kazmaier, *Organic Reactions, Vol 112a: A half-century of the Ugi reaction: classic variant*, John Wiley & Sons, Hoboken, New Jersey, 2023, pp. 1–1050; (d) C. Liu, L. G. Voskressensky and E. V. Van der Eycken, Recent Advances in the Synthesis of Peptidomimetics via Ugi Reactions, *Chem. – Eur. J.*, 2024, **30**, e202303597.
- 19 (a) S. Heck and A. Dömling, A versatile multi-component one-pot thiazole synthesis, *Synlett*, 2000, 424–426; (b) M. Umkehrer, J. Kolb, C. Burdack and W. Hiller, 2, 4, 5-Trisubstituted thiazole building blocks by a novel multi-component reaction, *Synlett*, 2005, 79–82; (c) A. Dömling and K. Illgen, 1-Isocyano-2-dimethylamino-alkenes: Versatile reagents in diversity-oriented organic synthesis, *Synthesis*, 2005, 662–667; (d) U. Kazmaier and S. Ackermann, A straightforward approach towards thiazoles and endothiopeptides via Ugi reaction, *Org. Biomol. Chem.*, 2005, **3**, 3184–3187; (e) U. Kazmaier and A. Persch, A straightforward approach towards 5-substituted thiazolyl-peptides via thio-Ugi-reaction, *Org. Biomol. Chem.*, 2010, **8**, 5442–5447.
- 20 (a) H. Kessler, R. Gratias, G. Hessler, M. Gurrath and G. Müller, Conformation of cyclic peptides. Principle concepts and the design of selectivity and superactivity in bioactive sequences by ‘spatial screening’, *Pure Appl. Chem.*, 1996, **68**, 1201–1205; (b) N. Loiseau, J.-M. Gomis, J. Santolini, M. Delaforge and F. André, Predicting the conformational states of cyclic tetrapeptides, *Biopolymers*, 2003, **69**, 363–385.

