

RESEARCH ARTICLE

View Article Online
View Journal | View IssueCite this: *Org. Chem. Front.*, 2024, **11**, 4442

One-pot multistep synthesis of 1-fluoroalkylisoquinolines and fused fluoroalkylpyridines from *N*-fluoroalkyl-1,2,3-triazoles†

Anna Kubičková,^{a,b} Svatava Voltrová,^a Adam Kleman,^a Blanka Klepetářová^a and Petr Beier^{a*}

An efficient one-pot microwave-assisted potassium fluoride-mediated synthesis of 1-fluoroalkyl-3-fluoroisoquinolines and fused fluoroalkylpyridines from *N*-fluoroalkylated 1,2,3-triazoles was developed. The reaction has a wide scope and allows the preparation of structurally diverse 3-fluoroisoquinolines with a fluoroalkyl group in position 1, a substituent in position 4 and a substituent on the fused benzene (or heteroaromatic) ring. *N*-Fluoroalkylated ketenimines, which undergo stereoselective formal 1,3-fluorine shift to difluoroazadienes, were identified as intermediates in the reaction sequence. The presence of fluorine in position 3 and a halogen in position 4 of the resulting isoquinolines allowed for further modification by nucleophilic aromatic substitution and cross-coupling reactions, respectively. The developed methodologies were utilized for the synthesis of derivatives of drug candidates.

Received 19th April 2024,
Accepted 17th June 2024

DOI: 10.1039/d4qo00713a

rsc.li/frontiers-organic

Introduction

The isoquinoline core is present in a variety of drugs and a large number of naturally occurring alkaloids, which in many cases possess compelling biological activities (Fig. 1).^{1–4} Numerous synthetic approaches exist leading to these benzopyridines, such as multistep sequences of reactions including the well-known Bischler–Napieralski,⁵ Pomeranz–Fritsch⁶ and Pictet–Spengler⁷ reactions, as well as processes involving transition metal catalysis.⁸

The introduction of fluorine atoms or fluoroalkyl groups into a lead molecule is a widely used strategy to enhance the pharmacologically relevant properties⁹ and several 1-trifluoromethylisoquinolines exploit this trend (Fig. 2), for example valigilurax¹⁰ – a positive allosteric modulator of mGlu4 receptors and a candidate for the treatment of Parkinson's disease. Yet, the procedures for their preparation remain underdeveloped, substrate-specific, low-yielding, or require expensive, non-selective and atom non-economical fluoroalkylation methods or transition metal catalysis.^{13,14}

The first multi-step approach leading to 1-perfluoroalkylisoquinolines with the Bischler–Napieralski type cyclization was demonstrated by Pastor¹⁵ in 1979. A similar approach was also used in 2019 by Lindsley (Scheme 1A) for the preparation of valigilurax, which allowed the construction of the isoquinoline core in an overall 21% yield.¹⁰ Another possible approach involves direct C–H bond perfluoroalkylation of isoquinolines¹⁶ (Scheme 1B) or isoquinoline-*N*-oxides.^{17,18} Trifluoromethylation *via* a coupling reaction of iodoisoquinolines with copper^{19–22} or palladium²³ catalysts was also reported. However, the most common strategy towards 1-fluoroalkylated isoquinolines involves the insertion of fluoroalkyl radicals into isonitriles, followed by radical cyclization (Scheme 1C).^{13,24–29}

Herein, we report a high-yielding and novel one-pot synthetic strategy to prepare substituted 1-fluoroalkyl-3-fluoroiso-




Fig. 1 Examples of an isoquinoline containing alkaloid (papaverine) and synthetic drugs (fasudil and ripasudil).

^aInstitute of Organic Chemistry and Biochemistry of the Czech Academy of Sciences, Flemingovo nám. 2, 16610 Prague 6, Czechia. E-mail: beier@uochb.cas.cz

^bUniversity of Chemistry and Technology, Technická 5, 166 28 Prague 6, Czechia

† Electronic supplementary information (ESI) available. CCDC 2336080–2336082.

For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d4qo00713a>







Fig. 2 Examples of bioactive 1-trifluoromethylisoquinolines.^{10–12}



Scheme 1 Selected literature syntheses of 1-fluoroalkylated isoquinolines (A–C) and our new approach from triazoles (D).

quinolines **2** based on thermal decomposition of *N*-fluoroalkyl-1,2,3-triazoles **1**, formal 1,3-fluorine shift, and cyclization (Scheme 1D). The presence of fluorine in position 3 and a halogen in position 4 of the isoquinoline ring enabled further modifications by nucleophilic aromatic substitution and/or cross-coupling reactions, respectively. The procedure is applicable also to heteroaryl substituted *N*-fluoroalkyl-1,2,3-triazoles affording heteroarenes with fused fluoroalkylated pyridine rings. The methodology thus allows the expansion of known chemical space to new selectively substituted fluoroalkylated isoquinoline-type structures with potential applications in life sciences.


Results and discussion

Recently, we reported thermal rearrangement of *N*-fluoroalkyl-1,2,3-triazoles^{30–33} **1** leading to *N*-fluoroalkylated ketenimines **3**.³⁴ We noticed that a prolonged heating of **3** led to new products identified by HRMS and NMR as isoquinolines **2** and

enamides **5**. Their formation can be explained by a thermally induced 1,3-fluorine shift of ketenimines **3** to two geometric isomers of azadienes **4**. Although four isomers of **4** can be theoretically formed by the fluorine shift, only the formation of two isomers was observed. The isomer (*Z,E*)-**4** cyclized to isoquinolines **2** while the isomer (*Z,Z*)-**4** only hydrolysed to enamides **5** (Scheme 2) (see the ESI† for three examples of isolated enamides **5**). A high-temperature NMR kinetic study revealed the time course of intermediate and product formation (Fig. 3).

A related transformation was briefly reported by Lermontov in 2002.³⁵ Thermal Huisgen cycloaddition of diphenylacetylene with ethyl 3-azido-2,2,3,3-tetrafluoropropanoate afforded isoquinolines and enamides in low yields. The authors wrongly assumed antiaromatic 1*H*-azirines to be the reactive intermediates (Scheme 3A), which we disproved with *ab initio* calculations in our previous study.³⁴ In another report, Molina showed the formation of an isoquinoline by ring closure of an *N*-styryl-substituted ketenimine (Scheme 3B).^{36–38}

In order to develop a general synthesis of 1-fluoroalkylated-2-fluoroisoquinolines **2** from triazoles **1** or ketenimines **3** we studied the effect of additives on the formation of **4**. Ideally, the formation of **4** should proceed stereoselectively to the *Z,E*-isomer. Therefore, the influence of additives on the stereoselectivity of the formal 1,3-fluorine shift of **3a** at room temperature was studied. While the addition of Et₃N, DBU, or BF₃·OEt₂ did not lead to efficient formation of **4a**, the addition of other basic additives or fluoride salts proved beneficial (Table 1). Carbonates induced the stereoselective transformation to the required (*Z,E*)-isomer of **4a** with Cs₂CO₃ reacting much faster than K₂CO₃ (entries 2 and 3) and Na₂CO₃ being unreactive (presumably due to its low solubility). However, decomposition of **4a** was observed in the basic conditions over time. A similar trend was observed in the case of inorganic fluorides with NaF being unreactive and CsF inducing the formation much faster than KF, but product decomposition and isomerization precluded its use in preparative experiments (entries 4–6; see the ESI† for the isomerization study of **4a** with CsF). Therefore, mildly basic KF was used as the additive of choice, accelerating the formal 1,3-fluorine shift of ketenimines **3** and providing a high selectivity to the required isomer of **4** for further cyclization. The origin of the stereoselectivity



Scheme 2 Proposed reaction mechanism of thermal additive-free decomposition of ketenimines **3** to isoquinolines **2** and side-products enamides **5**.








Fig. 3 Conversion of **3a** ($R = H$, $R_f = CF_3$) vs. reaction time for the formation of intermediates **4a** and product **2a** determined by 1H NMR ($C_2D_4Cl_2$; $140^\circ C$).

A: Thermal Huisgen cycloaddition and cyclization



B: Ring closure of *N*-styryl ketenimines



Scheme 3 Published preparations of isoquinolines by thermal Huisgen cyclization (A) and from *N*-styryl-substituted ketenimines (B).

Table 1 Influence of additives on the formation of isomers of **4a**


Entry	Additive	^{19}F NMR yield 3a / (Z,E)-4a / (Z,Z)-4a (%)		
		1 h	5 h	24 h
1	KHCO ₃	78 : 22 : 0	49 : 45 : 0	23 : 62 : 0
2	K ₂ CO ₃	83 : 17 : 0	57 : 38 : 0	0 : 83 : 0
3	Cs ₂ CO ₃	0 : 86 : 0	0 : 43 : 0	0 : 0 : 0
4	NaF	98 : 0 : 0	98 : 0 : 0	98 : 0 : 0
5	KF	83 : 17 : 0	50 : 50 : 0	13 : 80 : 0
6	CsF	0 : 89 : 0	0 : 79 : 0	0 : 50 : 12

of the thermal or heterogeneous additive-mediated 1,3-fluorine shift is unknown; however, we propose the steric factor to be dominant with fluoride addition to the central sp carbon atom of the ketenimine proceeding *trans* to the large aryl group followed by fluoride elimination from the CF₂ group (Scheme 4).

Difluoroazadiene (*Z,E*)-**4a** was prepared using CsF (Scheme 5). The structure of its derivative **4m** was confirmed by X-ray crystallography. Furthermore, addition of sodium acetate to **3a** efficiently afforded acetate **6a**, confirming that



Scheme 4 Fluoride-mediated formation of azadienes **4a** from ketenimine **3a**.



Scheme 5 Characterized azadienes **4** and **6**.

indeed a suitable nucleophile can add to the sp carbon of ketenimine **3a**, followed by fluoride elimination. Attempts to use chloride or iodide salts were unsuccessful.


An optimization study revealed that under microwave heating conditions a slight excess of KF afforded the formation of isoquinoline **2a** directly from triazole **1a** without the need to isolate the intermediates ketenimine **3a** or difluoroazadiene **4a** (Table 2). As for the solvent effect, we previously reported that the formation of ketenimines **3** from triazoles **1** works best in DCE but other solvents (chloroform, THF, toluene, cyclohexane, acetone) can also be used.³⁴ In this study we chose DCE as the optimal solvent.

With the optimized set of conditions, we expanded the multistep one-pot process to diversely substituted *N*-fluoroalkylated 1,2,3-triazoles **1** (Table 3). The method tolerated various functional groups on the aryl moiety, including electron-neutral, electron-rich and electron-poor substituents on the phenyl group with slightly decreased yields in the last

Table 2 Optimization of potassium fluoride-accelerated synthesis of isoquinoline **2a** from triazole **1a**

Entry	KF (equiv.)	^{19}F NMR ratio 3a / (Z,E)-4a / 2a
1	0.05	70 : 20 : 10
2	0.2	48 : 28 : 24
4	0.7	7 : 5 : 88
5	1.0	3 : 3 : 94
6	1.1	0 : 0 : 99
7	2.0	0 : 0 : 99



Table 3 Substrate scope of KF-mediated synthesis of isoquinolines 2

case. Different substitution positions on the aryl group were also well tolerated with differently substituted isoquinolines being produced from *o*-, *m*- or *p*-substituted aryls. In the case of *m*-substituted aryls, two isomers of the products were formed (2i) with good regioselectivity. In another case, the reaction was regioselective (2j). Substrates with large (11) or heteroaromatic (1m) groups also underwent the reaction to afford

unique isoquinolines or fused pyridines; however, the pyridyl-substituted triazole (1n) was found to be unreactive and only decomposition to a complex mixture of products was observed.

Position 5 of the starting triazole ring can be substituted with a halogen or alkyl group, which introduced these functions into position 4 of the final isoquinoline with various degrees of efficiency. The observed trend can be explained by steric factors where bulky substituents on the ketenimine sp^2 carbon atom hindered the attack of the fluoride ion to form the productive isomer of azadienes 4.

The methodology was found to display an excellent robustness with regards to the fluoroalkyl substituent in position 1 of the products. Not only the trifluoromethyl group, but also difluoromethyl, substituted difluoromethylene and ethoxycarbonyl substituents can be introduced efficiently. Under solvent free conditions difluoroisoquinoline 2x was prepared in low yield due to its high volatility and some side reactions.

The presence of a fluorine substituent in isoquinolines 2 in the activated position called for the investigation of substitution with various nucleophiles by S_NAr which expanded the diversity of accessible 1-fluoroalkylated isoquinolines. Thus, the fluorine atom of isoquinolines 2 was readily substituted with various oxygen, sulfur, and nitrogen nucleophiles in polar solvents to obtain heteroatom-substituted 1-trifluoromethyl isoquinolines 7 (Table 4).


Furthermore, isoquinolines 2 were used for the preparation of a small library of nine analogues of the TRPM8 antagonist shown in Fig. 2. Compounds 8 were easily accessed by nucleophilic sulfonamidation of 2 (Scheme 6), demonstrating the value of our approach in the synthesis of fluorinated and fluoroalkylated isoquinolines and their structurally diverse derivatives in drug development.

Other investigated follow-up derivatizations of compounds 2 were the cross-coupling reactions. Suzuki–Miyaura coupling of arylboronic acids with chloroisoquinoline 2p afforded coupling products 9a–c in high yields (Scheme 7). Heck, Sonogashira and Buchwald–Hartwig reactions of bromoisoqui-


Table 4 S_NAr of isoquinolines 2

Entry	Nucleophile (equiv.)	X	Solvent	Temp. (°C)	7, yield (%)
1	NaOH (15)	H	H ₂ O	155	7a, 88
2	EtONa (12)	H	EtOH	80	7b, 96
3	EtONa (12)	Cl	EtOH	155	7c, 99
4	<i>t</i> -BuOK (1.2)	H	<i>t</i> -BuOH	80	7d, 80
5	PhONa (1.5)	H	DMA	80	7e, 89
6	MeSNa (5)	H	DMA	20	7f, 85
7	MeSNa (2)	Ph	DMF	20	7g, 91
8	<i>p</i> -Tol-SNa (1)	H	DMA	80	7h, 91
9	<i>p</i> -Tol-SO ₂ Li (2.5)	H	DMSO	155	7i, 58
10	NH ₂ NH ₂ (20)	H	<i>i</i> -PrOH	100	7j, 95
11	<i>p</i> -Tol-CH ₂ NH ₂ (2)	H	DMSO	155	7k, 42







Scheme 6 Analogues of the TRPM8 antagonist.



Scheme 7 Suzuki–Miyaura coupling reactions with 2p.

noline **2q** also worked well giving the coupling products **10a–c** (Scheme 8).

The developed method for the synthesis of fluorinated isoquinolines was used for the preparation of 3-fluoro analogue **12** of the drug candidate valigluxax. The brominated isoquinoline **2j** was used for Pd-catalyzed amination, followed by protecting group removal to give analogue **12** in high yields (Scheme 9).

Scheme 8 Heck, Sonogashira and Buchwald–Hartwig coupling reactions of **2r** (see the ESI† for detailed conditions).Scheme 9 Synthesis of valigluxax analogue **12**.

Conclusions

In conclusion, microwave heating of *N*-fluoroalkyl-1,2,3-triazoles in the presence of potassium fluoride led to a series of events involving triazole ring opening, nitrogen molecule elimination, rearrangement, stereoselective formal 1,3-fluorine shift, and finally cyclization to produce diverse 1-fluoroalkylated 3-fluoroisoquinolines in good yields and with excellent substrate scope. Nucleophilic aromatic substitution of the fluorine atom in position 3 with heteroatom nucleophiles afforded 1-fluoroalkylated 3-substituted isoquinolines. Cross-coupling reactions of halogen atoms in position 4 of the isoquinolines gave derivatives with aryl, alkenyl, alkynyl or alkylamino groups. This synthetic approach to novel selectively fluorinated isoquinolines was applied in the synthesis of analogues of two families of drug candidates.

Data availability

The data supporting this article have been included as part of the ESI.†

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was financially supported by the Czech Academy of Sciences (research plan RVO: 61388963) and the Czech Science Foundation (project 23-04659S). This work was supported by the grant of Specific University Research – grant no. A2_FCHT_2024_001.

References

- 1 E. Plazas, M. M. C. Avila, D. R. Muñoz and S. L. E. Cuca, Natural Isoquinoline Alkaloids: Pharmacological Features and Multi-Target Potential for Complex Diseases, *Pharmacol. Res.*, 2022, **177**, 106126, DOI: [10.1016/j.phrs.2022.106126](https://doi.org/10.1016/j.phrs.2022.106126).



- 2 X.-F. Shang, C.-J. Yang, S. L. Morris-Natschke, J.-C. Li, X.-D. Yin, Y.-Q. Liu, X. Guo, J.-W. Peng, M. Goto, J.-Y. Zhang and K.-H. Lee, Biologically Active Isoquinoline Alkaloids Covering 2014–2018, *Med. Res. Rev.*, 2020, **40**(6), 2212–2289, DOI: [10.1002/med.21703](https://doi.org/10.1002/med.21703).
- 3 G. Diaz, I. L. Miranda and M. A. N. Diaz, Quinolines, Isoquinolines, Angustureine, and Congeneric Alkaloids—Occurrence, Chemistry, and Biological Activity, in *Phytochemicals - Isolation, Characterisation and Role in Human Health*, IntechOpen, 2015. DOI: [10.5772/59819](https://doi.org/10.5772/59819).
- 4 A. Vijayakumar, M. Manod, R. B. Krishna, A. Mathew and C. Mohan, Diversely Functionalized Isoquinolines and Their Core-Embedded Heterocyclic Frameworks: A Privileged Scaffold for Medicinal Chemistry, *RSC Med. Chem.*, 2023, **14**, 2509–2534, DOI: [10.1039/D3MD00248A](https://doi.org/10.1039/D3MD00248A).
- 5 A. Bischler and B. Napieralski, Zur Kenntniss Einer Neuen Isochinolinsynthese, *Ber. Dtsch. Chem. Ges.*, 1893, **26**(2), 1903–1908, DOI: [10.1002/cber.189302602143](https://doi.org/10.1002/cber.189302602143).
- 6 C. Pomeranz, Über eine neue Isochinolinsynthese, *Monatsh. Chem.*, 1893, **14**(1), 116–119, DOI: [10.1007/BF01517862](https://doi.org/10.1007/BF01517862).
- 7 A. Pictet and T. Spengler, Über Die Bildung von Isochinolin-Derivaten Durch Einwirkung von Methylal Auf Phenyl-Äthylamin, Phenyl-Alanin Und Tyrosin, *Ber. Dtsch. Chem. Ges.*, 1911, **44**(3), 2030–2036, DOI: [10.1002/cber.19110440309](https://doi.org/10.1002/cber.19110440309).
- 8 R. Gujjarappa, N. Vodnala and C. C. Malakar, Comprehensive Strategies for the Synthesis of Isoquinolines: Progress Since 2008, *Adv. Synth. Catal.*, 2020, **362**(22), 4896–4990, DOI: [10.1002/adsc.202000658](https://doi.org/10.1002/adsc.202000658).
- 9 C. Zhang, K. Yan, C. Fu, H. Peng, C. J. Hawker and A. K. Whittaker, Biological Utility of Fluorinated Compounds: From Materials Design to Molecular Imaging, Therapeutics and Environmental Remediation, *Chem. Rev.*, 2022, **122**(1), 167–208, DOI: [10.1021/acs.chemrev.1c00632](https://doi.org/10.1021/acs.chemrev.1c00632).
- 10 J. D. Panarese, D. W. Engers, Y.-J. Wu, J. J. Bronson, J. E. Macor, A. Chun, A. L. Rodriguez, A. S. Felts, J. L. Engers, M. T. Loch, K. A. Emmitte, A. L. Castelhana, M. J. Kates, M. A. Nader, C. K. Jones, A. L. Blobaum, P. J. Conn, C. M. Niswender, C. R. Hopkins and C. W. Lindsley, Discovery of VU2957 (Valiglurax): An mGlu4 Positive Allosteric Modulator Evaluated as a Preclinical Candidate for the Treatment of Parkinson's Disease, *ACS Med. Chem. Lett.*, 2019, **10**(3), 255–260, DOI: [10.1021/acsmchemlett.8b00426](https://doi.org/10.1021/acsmchemlett.8b00426).
- 11 Y. Tsuzuki, D. Sawamoto, T. Sakamoto, T. Kato, Y. Niwa and N. Awai, Sulfonamide Compounds Having TRPM8 Antagonistic Activity. US9540360B2, 2017.
- 12 W.-B. Ho, H. Zhao, S. Deng, D. Ng, L. R. Wright, M. Wu, X. Zhou, M. P. Arend and L. A. Flippin, 4-Hydroxy-Isoquinoline Compounds as Hif Hydroxylase, *Inhibitors*, **2866556**, 2020.
- 13 K. Aradi and L. Kiss, Recent Advances in the Trifluoromethylation Reactions of Isonitriles To Construct CF₃-Substituted *N*-Heterocycles, *Synthesis*, 2023, **55**(12), 1834–1843, DOI: [10.1055/a-2020-9090](https://doi.org/10.1055/a-2020-9090).
- 14 J. C. Sloop, Advances in the Preparation of Fluorinated Isoquinolines: A Decade of Progress, *J. Chem.*, 2017, **2017**, e2860123, DOI: [10.1155/2017/2860123](https://doi.org/10.1155/2017/2860123).
- 15 P. R. Pastor and A. Cambon, Synthese d'isoquinoleines F-Alkyl Substituees, *J. Fluor. Chem.*, 1979, **13**(4), 279–296, DOI: [10.1016/S0022-1139\(00\)82079-5](https://doi.org/10.1016/S0022-1139(00)82079-5).
- 16 Y. Pan, J. Li, Z. Li, F. Huang, X. Ma, W. Jiao and H. Shao, An Efficient and Facile Strategy for Trifluoromethylation and Perfluoroalkylation of Isoquinolines and Heteroarenes, *Chem. Commun.*, 2020, **56**(56), 7813–7816, DOI: [10.1039/D0CC00963F](https://doi.org/10.1039/D0CC00963F).
- 17 C. Fan, J. Song, G. Qiu, G. Liu and J. Wu, Generation of 1-(Trifluoromethyl)Isoquinolines via a Copper-Catalyzed Reaction of Isoquinoline-*N*-Oxide with Togni Reagent, *Org. Chem. Front.*, 2014, **1**(8), 924–928, DOI: [10.1039/C4QO00173G](https://doi.org/10.1039/C4QO00173G).
- 18 D. E. Stephens, G. Chavez, M. Valdes, M. Dovalina, H. D. Arman and O. V. Larionov, Synthetic and Mechanistic Aspects of the Regioselective Base-Mediated Reaction of Perfluoroalkyl- and Perfluoroarylsilanes with Heterocyclic *N*-Oxides, *Org. Biomol. Chem.*, 2014, **12**(32), 6190–6199, DOI: [10.1039/C4OB01088D](https://doi.org/10.1039/C4OB01088D).
- 19 M. Chen and S. L. Buchwald, Rapid and Efficient Trifluoromethylation of Aromatic and Heteroaromatic Compounds Using Potassium Trifluoroacetate Enabled by a Flow System, *Angew. Chem., Int. Ed.*, 2013, **52**(44), 11628–11631, DOI: [10.1002/anie.201306094](https://doi.org/10.1002/anie.201306094).
- 20 Z. Gonda, S. Kovács, C. Wéber, T. Gáti, A. Mészáros, A. Kotschy and Z. Novák, Efficient Copper-Catalyzed Trifluoromethylation of Aromatic and Heteroaromatic Iodides: The Beneficial Anchoring Effect of Borates, *Org. Lett.*, 2014, **16**(16), 4268–4271, DOI: [10.1021/ol501967c](https://doi.org/10.1021/ol501967c).
- 21 T. Uchikura, N. Kamiyama, T. Ishikawa and T. Akiyama, Catalytic Trifluoromethylation of Iodoarenes by Use of 2-Trifluoromethylated Benzimidazole as Trifluoromethylating Reagent, *Beilstein J. Org. Chem.*, 2020, **16**(1), 2442–2447, DOI: [10.3762/bjoc.16.198](https://doi.org/10.3762/bjoc.16.198).
- 22 Y. Liu, X. Shao, P. Zhang, L. Lu and Q. Shen, Trifluoromethyl-Substituted Sulfonium Ylide: Rh-Catalyzed Carbenoid Addition to Trifluoromethylthioether, *Org. Lett.*, 2015, **17**(11), 2752–2755, DOI: [10.1021/acs.orglett.5b01170](https://doi.org/10.1021/acs.orglett.5b01170).
- 23 K. Aikawa, H. Serizawa, K. Ishii and K. Mikami, Palladium-Catalyzed Negishi Cross-Coupling Reaction of Aryl Halides with (Difluoromethyl)Zinc Reagent, *Org. Lett.*, 2016, **18**(15), 3690–3693, DOI: [10.1021/acs.orglett.6b01734](https://doi.org/10.1021/acs.orglett.6b01734).
- 24 B. Zhang and A. Studer, 1-Trifluoromethylated Isoquinolines via Radical Trifluoromethylation of Isonitriles, *Org. Biomol. Chem.*, 2014, **12**(48), 9895–9898, DOI: [10.1039/C4OB02145B](https://doi.org/10.1039/C4OB02145B).
- 25 L. Wang and A. Studer, 1-Trifluoromethylisoquinolines from α -Benzylated Tosylmethyl Isocyanide Derivatives in a Modular Approach, *Org. Lett.*, 2017, **19**(20), 5701–5704, DOI: [10.1021/acs.orglett.7b02882](https://doi.org/10.1021/acs.orglett.7b02882).
- 26 Y. Liu, K. Zhang, W. Jiang, Y. Yang, Y. Jiang, X. Liu, Y. Xie, J. Wu, J. Cai and X.-H. Xu, Synthesis of 1-Difluoroalkylated Isoquinolines via Palladium-Catalyzed Radical Cascade



- Difluoroalkylation–Cyclization of Vinyl Isocyanides with Bromodifluoroacetic Derivatives, *Chem. – Asian J.*, 2017, **12**(5), 568–576, DOI: [10.1002/asia.201601645](https://doi.org/10.1002/asia.201601645).
- 27 F. Ding, Y. Jiang, K. Lin and L. Shi, Tandem Radical Cyclization for the Construction of 1-Difluoroalkylated Isoquinolines via Cu Catalyzed and Visible Light-Promoted Pathways, *Org. Biomol. Chem.*, 2018, **16**(11), 1812–1815, DOI: [10.1039/C8OB00232K](https://doi.org/10.1039/C8OB00232K).
- 28 Y. Wang, Z. Cao, Q. He, X. Huang, J. Liu, H. Neumann, G. Chen and M. Beller, Activation of Perfluoroalkyl Iodides by Anions: Extending the Scope of Halogen Bond Activation to C(Sp³)–H Amidation, C(Sp²)–H Iodination, and Perfluoroalkylation Reactions, *Chem. Sci.*, 2023, **14**(7), 1732–1741, DOI: [10.1039/D2SC06145G](https://doi.org/10.1039/D2SC06145G).
- 29 R. Ruzziconi and F. Buonerba, Recent Advances in the Synthesis of Regioselectively Fluorinated Homo- and Heterocyclic Compounds by Complementary Cyclization Methods, *J. Fluor. Chem.*, 2013, **152**, 12–28, DOI: [10.1016/j.jfluchem.2013.05.001](https://doi.org/10.1016/j.jfluchem.2013.05.001).
- 30 Z. E. Blastik, S. Voltrová, V. Matoušek, B. Jurásek, D. W. Manley, B. Klepetářová and P. Beier, Azidoperfluoroalkanes: Synthesis and Application in Copper (i)-Catalyzed Azide-Alkyne Cycloaddition, *Angew. Chem., Int. Ed.*, 2017, **56**, 346–349, DOI: [10.1002/anie.201609715](https://doi.org/10.1002/anie.201609715).
- 31 S. Voltrová, I. Putovný, V. Matoušek, B. Klepetářová and P. Beier, Reintroducing Azidodifluoromethane: Synthesis, Stability and [3 + 2] Cycloadditions, *Eur. J. Org. Chem.*, 2018, 5087–5090, DOI: [10.1002/ejoc.201800650](https://doi.org/10.1002/ejoc.201800650).
- 32 E. Shaitanova, V. Matoušek, T. Herentin, M. Adamec, R. Matyáš, B. Klepetářová and P. Beier, Synthesis and Cycloaddition Reactions of 1-Azido-1,1,2,2-tetrafluoroethane, *J. Org. Chem.*, 2023, **88**, 14969–14977, DOI: [10.1021/acs.joc.3c01346](https://doi.org/10.1021/acs.joc.3c01346).
- 33 M. Ziabko, B. Klepetářová and P. Beier, Synthesis of Azidodifluoromethyl Phenyl Sulfone and Its Use as a Synthetic Equivalent of the Azidodifluoromethyl Anion, *J. Org. Chem.*, 2023, **88**, 6939–6946, DOI: [10.1021/acs.joc.3c00256](https://doi.org/10.1021/acs.joc.3c00256).
- 34 A. Kubičková, A. Markos, S. Voltrová, A. Marková, J. Filgas, B. Klepetářová, P. Slavíček and P. Beier, Aza-Wolff Rearrangement of *N*-Fluoroalkyl Triazoles to Ketenimines, *Org. Chem. Front.*, 2023, **10**(13), 3201–3206, DOI: [10.1039/D3QO00618B](https://doi.org/10.1039/D3QO00618B).
- 35 S. A. Lermontov, S. V. Shkavrov, A. N. Pushin and V. V. Tkachev, Unusual Fluorination of Diphenylacetylene with Methyl 3-Azidotetrafluoropropionate, *Russ. J. Gen. Chem.*, 2002, **72**(8), 1289–1290, DOI: [10.1023/A:1020804502390](https://doi.org/10.1023/A:1020804502390).
- 36 P. Molina, M. Alajarín, A. Vidal, J. Feneau-Dupont and J. P. Declercq, One-Pot Preparation of Derivatives of the Unknown Fluoreno[2,3,4-*ij*]Isoquinoline Ring from Conjugated Ketenimines by a Consecutive Electrocyclic Ring-Closure/Claisen Rearrangement/Intramolecular Diels–Alder Cycloaddition/Double Aromatization Process, *J. Chem. Soc., Chem. Commun.*, 1990, (no. 11), 829–831, DOI: [10.1039/C39900000829](https://doi.org/10.1039/C39900000829).
- 37 P. Molina, M. Alajarín, A. Vidal, J. Feneau-Dupont and J. P. Declercq, Domino Reactions. One-Pot Preparation of Fluoreno[2,3,4-*ij*]Isoquinoline Derivatives from Conjugated Ketene Imines, *J. Org. Chem.*, 1991, **56**(12), 4008–4016, DOI: [10.1021/jo00012a039](https://doi.org/10.1021/jo00012a039).
- 38 P. Molina, A. Vidal and I. Barquero, Regiospecific Electrocyclization of β -Arylvinyl Ketenimines. Formal Syntheses of the Alkaloid from Marine Origin Aaptamine, *Synthesis*, 1996, (10), 1199–1202, DOI: [10.1055/s-1996-4367](https://doi.org/10.1055/s-1996-4367).

