

## RESEARCH ARTICLE

View Article Online  
View Journal | View IssueCite this: *Org. Chem. Front.*, 2024,  
11, 7147Electrochemically promoted diamidation of  
alkenes to dihydroimidazole skeleton†Guo-Ao Wang, Kai-Wei Chen, Yu-Da Huang, Yu-Yuan Zhang, Xiu-Jin Meng,\*  
Fei-Hu Cui,\* Ying-Ming Pan and Hai-Tao Tang \*

The dihydroimidazole framework is a valuable heterocyclic compound widely found in natural products, ligands, molecular catalysts, pesticides, etc. The construction of a dihydroimidazole framework from alkenes is an attractive synthesis approach. Herein, we report a simple and environmentally friendly electrocatalytic diamination reaction of alkenes. The reaction involves constructing a dihydroimidazole framework from readily available alkenes, CH<sub>3</sub>CN, and H<sub>2</sub>O. CH<sub>3</sub>CN serves as a solvent and a nitrogen source, whereas H<sub>2</sub>O provides an oxygen source without the need for catalysts and oxidants. The synthesized 3,4-dihydroimidazole compound can be further hydrolyzed into vicinal diamines. As far as we know, the synthesis of dihydroimidazole frameworks through electrochemical Ritter-type diamination of alkenes has not been reported yet.

Received 5th August 2024,  
Accepted 11th October 2024

DOI: 10.1039/d4qo01445f

rsc.li/frontiers-organic

## Introduction

The 1,2-diamine motif is widely present in natural products, pharmaceutical compounds, and molecular catalysts (Fig. 1).<sup>1</sup> Difunctionalization of alkenes<sup>2</sup> with amination reagents, which is one of the most straightforward and practical strategies<sup>3</sup> for assembling scaffolds, has achieved remarkable progress<sup>4</sup> in the past few decades. However, these reported methods often require the utilization of transition metal reagents and even stoichiometric amounts of osmium or cobalt.<sup>4a,5</sup> Moreover, the use of stoichiometric amounts of chemical oxidants (e.g., iodine(III) reagents or organic peroxides)<sup>4d,6</sup> or azide reagents<sup>4b,c,7</sup> raises costs and environmental and safety issues, whereas the use of strong oxidizers may produce environmentally hazardous by-products and pose an explosion hazard when used with azide sources, particularly for large-scale applications.<sup>8</sup> In addition, they are often limited in substrate scope and sometimes require special amination reagents (e.g., diaziridinone and its analogs<sup>4e,9</sup> or azido-iodine compounds).<sup>4c</sup> Therefore, developing highly efficient, mild, and green synthetic strategies is still desirable, and an

additional substrate combination for the 1,2-diamine motif should be found to address the aforementioned drawbacks.

Electrochemistry offers an environmentally friendly, gentle, and efficient alternative to traditional chemical methods of redox conversion.<sup>10</sup> In the alternative method, common organic starting materials can lose or gain electrons at the electrode surface without the need for an external redox agent; thus, highly reactive intermediates are readily produced. Electrochemistry allows the precise external control of chemoselectivity and flux of reactive intermediates by regulating the applied current or potential. Compared with the energy efficiency in traditional chemical methods, that in the alternative method is maximized because of the latter's precise external control. Recently, electrochemical anodic oxidation has




Fig. 1 Dihydroimidazole-based drugs and bioactive molecules.

State Key Laboratory for Chemistry and Molecular Engineering of Medicinal Resources, Key Laboratory for Chemistry and Molecular Engineering of Medicinal Resources (Ministry of Education of China), School of Chemistry and Pharmaceutical Sciences of Guangxi Normal University, Guilin 541004, People's Republic of China. E-mail: xiujinmeng@mailbox.gxnu.edu.cn, cuifeihuao@163.com, httang@gxnu.edu.cn

† Electronic supplementary information (ESI) available. CCDC 2368464. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d4qo01445f>

shown great potential in the construction of a 1,2-diamine motif.

For instance, Xu and his colleagues<sup>11</sup> reported an electrochemical reaction of synthesizing 1,2-diamine compounds with diastereoselectivity using aryl alkenes and sulfonamides mediated by tris (2,4-dibromophenyl) amine. Lin's group<sup>7b</sup> has also made remarkable progress in the electrocatalytic diazidation of alkenes (Scheme 1). Despite the remarkable progress made above, issues such as limited types of nitrogen sources and 1,2-diamine structures may exist. Dihydroimidazole frameworks are a class of important nitrogen-containing heterocyclic compounds<sup>12</sup> with the structural characteristics of two nitrogen atoms and two sp<sup>3</sup> C–H bonds, which can be easily synthesized by 1,2-diamination of alkenes in theory. However, electrochemical synthesis of dihydroimidazole has not been reported because of the lack of well-matched nitrogen sources, alkenes, and electrochemical conditions.

Herein, we report a simple and environmentally friendly electrocatalytic diamination reaction of alkenes. The reaction involves constructing a dihydroimidazole framework from readily available alkenes, CH<sub>3</sub>CN, and H<sub>2</sub>O. CH<sub>3</sub>CN serves as a solvent and a nitrogen source,<sup>13</sup> whereas H<sub>2</sub>O provides an oxygen source without the need for catalysts and oxidants. The synthesized 3,4-dihydroimidazole compound can be further hydrolyzed into vicinal diamines. As far as we know, the synthesis of dihydroimidazole frameworks through electrochemical Ritter-type diamination of alkenes has not been reported yet.


## Results and discussion

We used 2-phenyl-1-propene (**1a**) as the model substrate to establish the optimal conditions for this reaction by optimizing the reaction conditions. Under the constant current condition of 8 mA, the reaction proceeded smoothly in a three-necked flask with Et<sub>4</sub>NBF<sub>4</sub> as the electrolyte, reticulated vitr-

eous carbon (RVC) electrode as the anode, and Ni foam electrode as the cathode. The required product **2a** (Table 1, entry 1) was obtained with an 82% yield. The yield decreased when the cathode was replaced with platinum or nickel sheets (Table 1, entries 2 and 3), whereas the yield of **2a** was 59% when the cathode and anode were replaced with platinum sheets (Table 1, entry 4). However, the yield of **2a** was only 35% when the cathode and anode were replaced with carbon rods (Table 1, entry 5). Increasing or decreasing the current would decrease the reaction activity (Table 1, entries 6 and 7). Only trace amounts of product **2a** were detected when Et<sub>4</sub>NCl, Et<sub>4</sub>NBr, and Et<sub>4</sub>NI were used as electrolytes (Table 1, entry 8). The reaction could not proceed without current (Table 1, entry 9). In summary, the optimal conditions established are as follows: RVC as anode, foam nickel electrode as cathode, Et<sub>4</sub>NBF<sub>4</sub> (4 equiv.) as electrolyte, extra dry CH<sub>3</sub>CN and H<sub>2</sub>O as solvent (extra dry CH<sub>3</sub>CN = 6 mL; H<sub>2</sub>O = 5 μL), and reaction at room temperature with a constant current of 8 mA for 3–6 h.

Under optimal conditions, the substrate range was explored. The results are shown in Schemes 2 and 3. First, the model substrate 2-phenyl-1-propene (**1a**) presented the target product in 82% yield, and it could obtain the target product with 76% (**2b**), 71% (**2c**), 73% (**2d**), 72% (**2e**), and 61% (**2f**) yields when the benzene ring was *para*-carried with groups such as fluorine, chlorine, bromine, trifluoromethyl, and nitro groups. When halogen groups such as fluorine and bromine groups exist in the *meta* position of the aromatic ring, the target product can be obtained with yields of 83% (**2g**) and 66% (**2h**), respectively. When a methyl group exists in the *meta* position of the aromatic ring, the target product can be obtained with a yield of 55% (**2j**). However, when the CH<sub>3</sub>CN


**Table 1** Optimization of the reaction conditions<sup>a</sup>




Entry	Variation from standard conditions	Yield <sup>b,c</sup> (%)
1	None	82
2	RVC (+) Pt (-) instead of RVC (+) Ni foam (-)	68
3	RVC (+) Ni (-) instead of RVC (+) Ni foam (-)	61
4	Pt (+) Pt (-) instead of RVC (+) Ni foam (-)	59
5	C (+) C (-) instead of RVC (+) Ni foam (-)	35
6	10 mA instead of 8 mA	64
7	6 mA instead of 8 mA	62
8	Et <sub>4</sub> NCl, Et <sub>4</sub> NBr, and Et <sub>4</sub> NI instead of Et <sub>4</sub> NBF <sub>4</sub>	Trace
9	No electricity	0

<sup>a</sup> Standard conditions: RVC (100 PPI, 1 cm × 1 cm × 1.2 cm) anode, Ni foam (1 cm × 2 cm × 0.5 cm) cathode, extra dry CH<sub>3</sub>CN and H<sub>2</sub>O as solvent (extra dry CH<sub>3</sub>CN = 6 mL, H<sub>2</sub>O = 5 μL), undivided cell, constant current = 8 mA, **1a** (0.2 mmol, 1 equiv.), Et<sub>4</sub>NBF<sub>4</sub> (0.8 mmol, 4 equiv.), and room temperature. <sup>b</sup> Isolated yields. <sup>c</sup> Unless otherwise specified, extra dry CH<sub>3</sub>CN with a water content of ≤10 ppm was used.

**a. Previous work:** Electrochemical and catalyst-promoted 1,2-diamination of alkenes




**b. This work:** First electrochemical synthesis of dihydroimidazole



- Metal and catalyst free
- H<sub>2</sub>O as an oxygen source
- Electrochemical multicomponent reaction
- CH<sub>3</sub>CN as a nitrogen source


**Scheme 1** 1,2-Diamination of alkenes.



**Scheme 2** The range of alkenes with substituents at the  $\alpha$ -position. Reaction conditions: RVC (100 PPI, 1 cm  $\times$  1 cm  $\times$  1.2 cm) anode, Ni foam (1 cm  $\times$  2 cm  $\times$  0.5 cm) cathode, undivided cell, constant current = 8 mA, **1a** (0.2 mmol, 1 equiv.), Et<sub>4</sub>NBF<sub>4</sub> (0.8 mmol, 4 equiv.), extra dry CH<sub>3</sub>CN and H<sub>2</sub>O as solvent (extra dry CH<sub>3</sub>CN = 6 mL, H<sub>2</sub>O = 5  $\mu$ L); electrolysis was carried out at room temperature under air atmosphere.

was replaced with butyronitrile, the target product (**2k**) could be obtained in 61% yield. When halogen groups such as chlorine are found in the adjacent position of the aromatic ring, the target product can be obtained with a yield of 65% (**2i**). When the R<sup>1</sup> group is replaced with ethyl, propyl, or phenyl, the yield of the target product gradually decreases as the steric hindrance increases. The target product can be obtained with yields of 74% (**2l**), 66% (**2m**), and 61% (**2n**). In addition, electrochemical methods can be used to modify natural products, such as *L*-menthol, 1-adamantane methanol, and 1-cyclododecanol, affording the corresponding derivatives (**2o–2q**) in 40%–55% yields.


When we replace 2-phenyl-1-propene (**1a**) with styrene (**3a**), the target product can still be obtained with a 75% (**4a**) yield. On the benzene ring of styrene, a series of different substitutions can be successfully carried out for electrochemical 1,2-diamination reactions, including fluorine, chlorine, bromine, trifluoromethyl, and methyl. Alkenes with 2,6-disubstituted



**Scheme 3** The range of alkenes without substituents at the  $\alpha$ -position. Reaction conditions: RVC (100 PPI, 1 cm  $\times$  1 cm  $\times$  1.2 cm) anode, Ni foam (1 cm  $\times$  2 cm  $\times$  0.5 cm) cathode, undivided cell, constant current = 8 mA, **3a** (0.2 mmol, 1 equiv.), Et<sub>4</sub>NBF<sub>4</sub> (0.8 mmol, 4 equiv.), extra dry CH<sub>3</sub>CN and H<sub>2</sub>O as solvent (extra dry CH<sub>3</sub>CN = 6 mL, H<sub>2</sub>O = 5  $\mu$ L); electrolysis was carried out at room temperature under air atmosphere.

benzene rings are also suitable substrates (**4i**). We also confirmed the structure of (**4j**) through X-ray diffraction analysis. In addition, the electrochemical reaction of 1,2-diamination is compatible with chloromethyl groups (**4e**). After switching to styrene, the yield is generally slightly lower than that of alkenes with substituents at the  $\alpha$ -position. The possible reason is that when substituents exist at the  $\alpha$ -position, stabilizing carbocations is easy, resulting in a high electrochemical reaction activity for 1,2-diamination. This finding is consistent with our speculated mechanism.


We conducted a gram-scale conversion of the electrocatalytic diamination reaction of alkenes under constant current conditions of 80 mA to demonstrate further the synthetic application of this electrochemical method. The reaction of alkene **1a** with CH<sub>3</sub>CN and H<sub>2</sub>O was completed at a scale of grams, and the target product **2a** (0.71 g) was obtained with a 41% yield (Scheme 4a). In addition, 1,2-diamine compounds are important synthetic units, such as drug intermediates, cat-




Scheme 4 Electrochemical applications.

alysts, and ligands. Therefore, synthesizing 1,2-diamine compounds from readily available alkenes is important. We dissolved 0.2 mmol **2a** in potassium hydroxide, methanol, and ethylene glycol, and it was heated under reflux. After the reaction was completed, the diamine compound **2aaa** was obtained with a 67% yield. In addition, **4e** can be used to modify BOC Glycine, providing the corresponding derivative with a yield of 45% (**4ee**).

We conducted several control experiments to gain a deep understanding of the reaction mechanism (Scheme 5). When 5 equiv. BHT and TEMPO are added to the reaction, only trace amounts of product **2a** are detected, indicating that the reaction may involve the free radical pathway (Scheme 5a). We conducted an  $\text{H}_2^{18}\text{O}$  labeling experiment to verify the source of oxygen in product **2a**. When  $\text{H}_2^{18}\text{O}$  (5  $\mu\text{L}$ ) is added to the reaction, the results show that the  $^{18}\text{O}$  labeled product **2aa** can be detected by HRMS analysis (Scheme 5b and Fig. S2†). The



Scheme 5 Control experiments.



Scheme 6 Possible reaction mechanism.

results indicate that  $\text{H}_2\text{O}$  may provide oxygen atoms for the formation of the target product **2a**. Finally, we conducted radical clock experiments using compounds **5a** and **6a**. Products **5aa** and **6aa** can be detected by HRMS analysis, indicating that the reaction is undergoing a free radical pathway (see ESI† for additional details).

We also conducted cyclic voltammetry analysis on compound **1a**. The test results show that an irreversible oxidation peak is observed at 2.45 V (relative to Ag/AgCl) in the cyclic voltammogram of **1a**, indicating that **1a** may be oxidized at the anode (RVC).

On the basis of the above experimental results and previous literature reports,<sup>14</sup> we proposed a possible mechanism (Scheme 6). 2-Phenyl-1-propene **1a** is oxidized at the anode (RVC) to form alkene radical cation **I**, which is captured by the  $\text{CH}_3\text{CN}$  to form intermediate **II**. Then, intermediate **II** loses an electron to intermediate **III**. Intermediate **III** is captured by a molecular  $\text{CH}_3\text{CN}$  to form intermediate **IV**, which then forms a carbocation intermediate. After hydrolysis, it forms amide product **VIII**, which is then cyclized to obtain the target product **2a**. This process is essentially a Ritter reaction,<sup>15</sup> with the difference being the methods and synthetic precursors for generating carbocations.

## Conclusions

We have developed a simple and environmentally friendly electrochemically driven 1,2-diamination reaction of alkenes, which is an effective method for constructing 1,2-diamine framework compounds from readily available alkenes,  $\text{CH}_3\text{CN}$ , and  $\text{H}_2\text{O}$ . The reaction does not require additional catalysts

and oxidants, and the raw materials are readily available. We hope that this protocol can provide a complementary way for dihydroimidazole compounds and be widely applied in modern synthetic chemistry and pharmaceutical research.

## Data availability

The data supporting this article have been included as part of the ESI.†

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

We thank the Guangxi Natural Science Foundation of China (2021GXNSFFA220005), National Natural Science Foundation of China (22061003, 22301048), the Natural Science Foundation of Guangxi Normal University (No. 2021JC004) for financial support.

## References

- (a) D. Lucet, T. L. Gall and C. Mioskowski, The Chemistry of Vicinal Diamines, *Angew. Chem., Int. Ed.*, 1998, **37**, 2580–2627; (b) S. R. S. S. Kotti, C. Timmons and G. Li, Vicinal Diamino Functionalities as Privileged Structural Elements in Bio-logically Active Compounds and Exploitation of their Synthetic Chemistry, *Chem. Biol. Drug Des.*, 2006, **67**, 101–114; (c) F. Cardona and A. Goti, Metal-Catalysed 1,2-Diamination Reactions, *Nat. Chem.*, 2009, **1**, 269–275.
- (a) A. Corma, Introduction: Chemicals from Coal, Alkynes, and Biofuels, *Chem. Rev.*, 2014, **114**, 1545–1546; (b) M. Crespo-Quesada, F. Cárdenas-Lizana, A.-L. Dessimoz and L. Kiwi-Minsker, Modern Trends in Catalyst and Process Design for Alkyne Hydrogenations, *ACS Catal.*, 2012, **2**, 1773–1786; (c) M. Eggersdorfer, D. Laudert, U. Létinois, T. McClymont, J. Medlock, T. Netscher and W. Bonrath, One Hundred Years of Vitamins—a Success Story of the Natural Sciences, *Angew. Chem., Int. Ed.*, 2012, **51**, 12960–12990; (d) R. Bai, G. He, L. Li, T. Zhang, J. Li, X. Wang, X. Wang, Y. Zou, D. Mei, A. Corma and J. Yu, Encapsulation of Palladium Carbide Subnanometric Species in Zeolite Boosts Highly Selective Semihydrogenation of Alkynes, *Angew. Chem., Int. Ed.*, 2023, **62**, e202313101; (e) B. Dong, J. Shen and L.-G. Xie, Recent Developments on 1,2-Difunctionalization and Hydrofunctionalization of Unactivated Alkenes and Alkynes Involving C–S Bond Formation, *Org. Chem. Front.*, 2023, **10**, 1322–1345; (f) K. Yang, J. Liu, D. Fu, L. Niu, S.-J. Li and Y. Lan, A Theoretical Study of Selective Radical Relay and Coupling Reactions for Alkene Difunctionalization, *Org. Chem. Front.*, 2023, **10**, 4336–4341.
- (a) R. Fu, M. Xu, Y. Wang, X. Wu and X. Bao, Organo-Photocatalytic Anti-Markovnikov Hydroamidation of Alkenes with Sulfonyl Azides: A Combined Experimental and Computational Study, *Angew. Chem., Int. Ed.*, 2024, **63**, e202406069; (b) L. Song, N. Fu, B. G. Ernst, W. H. Lee, M. O. Frederick, R. A. DiStasio and S. Lin, Dual Electrocatalysis Enables Enantioselective Hydrocyanation of Conjugated Alkenes, *Nat. Chem.*, 2020, **12**, 747–754.
- (a) P. N. Becker, M. A. White and R. G. Bergman, A New Method for 1,2-Diamination of Alkenes Using Cyclopentadienyl-nitrosylcobalt Dimer/NO/LiAlH<sub>4</sub>, *J. Am. Chem. Soc.*, 1980, **102**, 5676–5677; (b) B. Zhang and A. Studer, Copper-Catalyzed Intermolecular Aminoazidation of Alkenes, *Org. Lett.*, 2014, **16**, 1790–1793; (c) G. Fumagalli, P. T. G. Rabet, S. Boyd and M. F. Greaney, Three-Component Azidation of Styrene-Type Double Bonds: Light-Switchable Behavior of a Copper Photoredox Catalyst, *Angew. Chem., Int. Ed.*, 2015, **54**, 11481–11484; (d) S.-J. Shen, C.-L. Zhu, D.-F. Lu and H. Xu, Iron-Catalyzed Direct Olefin Diazidation via Peroxyester Activation Promoted by Nitrogen-Based Ligands, *ACS Catal.*, 2018, **8**, 4473–4482; (e) Y. Wen, B. Zhao and Y. Shi, Cu(I)-Catalyzed Diamination of Disubstituted Terminal Olefins: An Approach to Potent NK1 Antagonist, *Org. Lett.*, 2009, **11**, 2365–2368.
- A. O. Chong, K. Oshima and K. B. Sharpless, Synthesis of Dioxobis(tert-alkylimido)osmium(VIII) and Oxotris(tert-alkylimido)osmium(VIII) Complexes. Stereospecific Vicinal Diamination of Olefins, *J. Am. Chem. Soc.*, 1977, **99**, 3420–3426.
- (a) D. E. Olson, J. Y. Su, D. A. Roberts and J. D. Bois, Vicinal Diamination of Alkenes under Rh-Catalysis, *J. Am. Chem. Soc.*, 2014, **136**, 13506–13509; (b) K. Muñoz, L. Barreiro, R. M. Romero and C. Martínez, Catalytic Asymmetric Diamination of Styrenes, *J. Am. Chem. Soc.*, 2017, **139**, 4354–4357; (c) Y.-A. Yuan, D.-F. Lu, Y.-R. Chen and H. Xu, Iron-Catalyzed Direct Diazidation for a Broad Range of Olefins, *Angew. Chem., Int. Ed.*, 2016, **55**, 534–538.
- (a) W. E. Fristad, T. A. Brandvold, J. R. Peterson and S. R. Thompson, Conversion of Alkenes to 1,2-Diazides and 1,2-Diamines, *J. Org. Chem.*, 1985, **50**, 3647–3649; (b) N. Fu, G. S. Sauer, A. Saha, A. Loo and S. Lin, Metal-Catalyzed Electrochemical Diazidation of Alkenes, *Science*, 2017, **357**, 575–579; (c) N. Fu, G. S. Sauer and S. Lin, A General, Electrocatalytic Approach to the Synthesis of Vicinal Diamines, *Nat. Protoc.*, 2018, **13**, 1725–1743; (d) J. C. Siu, J. B. Parry and S. Lin, Aminoxyl-Catalyzed Electrochemical Diazidation of Alkenes Mediated by a Metastable Charge-Transfer Complex, *J. Am. Chem. Soc.*, 2019, **141**, 2825–2831.
- (a) S. Caron, R. W. Dugger, S. G. Ruggeri, J. A. Ragan and D. H. Brown, Ripin. Large-Scale Oxidations in the Pharmaceutical Industry, *Chem. Rev.*, 2006, **106**, 2943–2989; (b) H.-T. Zhu, L. Arosio, R. Villa, M. Nebuloni and H. Xu,

- Process Safety Assessment of the Iron-Catalyzed Direct Olefin Diazidation for the Expedient Synthesis of Vicinal Primary Diamines, *Org. Process Res. Dev.*, 2017, **21**, 2068–2072.
- 9 (a) Y. Zhu, R. G. Cornwall, H. Du, B. Zhao and Y. Shi, Catalytic Diamination of Olefins via N–N Bond Activation, *Acc. Chem. Res.*, 2014, **47**, 3665–3678; (b) B. Zhao, W. Yuan, H. Du and Y. Shi, Cu(i)-Catalyzed Intermolecular Diamination of Activated Terminal Olefins, *Org. Lett.*, 2007, **9**, 4943–4945.
- 10 (a) Z. Zhang, X.-J. Meng, F.-H. Cui, H.-T. Tang, Y.-C. Wang, G.-B. Huang and Y.-M. Pan, Electrochemically Promoted Three-Component Reaction to *N*-Sulfonyl Amidines, *Org. Lett.*, 2024, **26**, 193–197; (b) Z.-L. Lei, T.-C. Liu, F.-H. Cui, Y.-M. Pan, S.-H. Li and H.-T. Tang, Electrochemical Promoted Three-Component Trifluoromethylation/Spirocyclization Reaction of *N*-Arylsulfonylacrylamides to 4-Azaspiro[4.5]Decanes, *Org. Lett.*, 2024, **25**, 6001–6005; (c) C. Yin, S. Tang, J. Mei, X. Hu and H. Zhang, Electrochemical Synthesis and Transformation of Organoboron Compounds, *Org. Chem. Front.*, 2023, **10**, 3361–3377; (d) Q. Wan, Z. Zhang, Z.-W. Hou and L. Wang, Recent Advances in the Electrochemical Generation of 1,3-Dicarbonyl Radicals from C–H Bonds, *Org. Chem. Front.*, 2023, **10**, 2830–2848; (e) H.-T. Tang, Y.-Z. Pan and Y.-M. Pan, Research Progress in Electrochemical/Photochemical Utilization of Methanol as a C1 Source, *Green Chem.*, 2023, **25**, 8313–8327; (f) X.-Q. Zhou, H.-T. Tang, F.-H. Cui, Y. Liang, S.-H. Li and Y.-M. Pan, Electrocatalytic Three-Component Reactions: Synthesis of Tellurium-Containing Oxazolidinone for Anticancer Agents, *Green Chem.*, 2023, **25**, 5024–5029; (g) T.-K. Xiong, Q. Xia, X.-Q. Zhou, S.-H. Li, F.-H. Cui, H.-T. Tang, Y.-M. Pan and Y. Liang, Electrochemically Mediated fixation of CO<sub>2</sub>: Synthesis of Functionalized Oxazolidine-2,4-Diones by Three-Component Reactions, *Adv. Synth. Catal.*, 2023, **365**, 2183–2187; (h) X.-Y. Wang, Y.-F. Zhong, Z.-Y. Mo, S.-H. Wu, Y.-L. Xu, H.-T. Tang and Y.-M. Pan, Synthesis of Seleno Oxindoles via Electrochemical Cyclization of *N*-arylacrylamides with Diorganyl Diselenides, *Adv. Synth. Catal.*, 2021, **363**, 208–214.
- 11 C.-Y. Cai, X.-M. Shu and H.-C. Xu, Practical and Stereoselective Electrocatalytic 1,2-Diamination of Alkenes, *Nat. Commun.*, 2019, **10**, 4953.
- 12 (a) L. T. Vassilev, B. T. Vu, B. Graves, D. Carvajal, F. Podlaski, Z. Filipovic, N. Kong, U. Kammlott, C. Lukacs, C. Klein, N. Fotouhi and E. A. Liu, In Vivo Activation of the p53 Pathway by Small-Molecule Antagonists of MDM2, *Science*, 2004, **303**, 844–848; (b) K. Murai, S. Fukushima, A. Nakamura, M. Shimura and H. Fujioka, C<sub>3</sub>-Symmetric Chiral Trisimidazoline: The Role of a Third Imidazoline and its Application to the Nitro Michael Reaction and the  $\alpha$ -Amination of  $\beta$ -Ketoesters, *Tetrahedron*, 2011, **67**, 4862–4868; (c) C. Dardonville and I. Rozas, Imidazoline Binding Sites and their Ligands: An Overview of the Different Chemical Structures, *Med. Res. Rev.*, 2004, **24**, 639–661.
- 13 (a) X.-Y. Su and P.-Q. Huang, Tf<sub>2</sub>O-Promoted Morgan–Walls Reaction: From a Flexible Approach to Functionalized Phenanthridines and Quinazolines to the Short and Divergent Total Syntheses of Alkaloids, *Synthesis*, 2023, **55**, 877–891; (b) B. H. Hoff, Acetonitrile as a Building Block and Reactant, *Synthesis*, 2018, **50**, 2824–2852.
- 14 (a) P. Xiong, H. Long, J. Song, Y. Wang, J.-F. Li and H.-C. Xu, Electrochemically Enabled Carbohydroxylation of Alkenes with H<sub>2</sub>O and Organotrifluoroborates, *J. Am. Chem. Soc.*, 2018, **140**, 16387–16391; (b) Y. Yu, Y. Yuan and K.-Y. Ye, Electrochemical Synthesis of Vicinal Azidoacetamides, *Chem. Commun.*, 2023, **59**, 422–425.
- 15 (a) J. J. Ritter and P. P. Minieri, A New Reaction of Nitriles. I. Amides from Alkenes and Mononitriles, *J. Am. Chem. Soc.*, 1948, **70**, 4045–4048; (b) J. J. Ritter and J. Kalish, A New Reaction of Nitriles. II. Synthesis of *t*-Carbinamines, *J. Am. Chem. Soc.*, 1948, **70**, 4048–4050; (c) A. Guérinot, S. Reymond and J. Cossy, Ritter Reaction: Recent Catalytic Developments, *Eur. J. Org. Chem.*, 2012, 19–28; (d) D. Jiang, T. He, L. Ma and Z. Wang, Recent Developments in Ritter Reaction, *RSC Adv.*, 2014, **4**, 64936–64946; (e) M.-E. Chen, X.-W. Chen, Y.-H. Hu, R. Ye, J.-W. Lv, B. Li and F.-M. Zhang, Recent Advances of Ritter Reaction and its Synthetic Applications, *Org. Chem. Front.*, 2021, **8**, 4623–4664; (f) L. Bao, B.-B. Zhang, Z.-X. Wang and X.-Y. Chen, Photocatalytic Dehydrations for the Ritter Reaction, *Org. Chem. Front.*, 2023, **10**, 1375–1379; (g) G. M. Ziarani, F. S. Hasankiadeh and F. Mohajer, Recent Applications of Ritter Reactions in Organic Syntheses, *ChemistrySelect*, 2020, **5**, 14349–14379; (h) J. Y. Becker, L. R. Byrd, L. L. Miller and Y.-H. So, Remote Anodic Substitution of Ketones, *J. Am. Chem. Soc.*, 1975, **97**, 853–856; (i) Y. Xu, Q. Li, R. Ye, B. Xu and X. Zhou, Electrochemical Oxidative C–H Amination through a Ritter-Type Reaction, *J. Org. Chem.*, 2023, **88**, 9518–9522; (j) T. Shen and T. H. Lambert, C-H Amination via Electrophotocatalytic Ritter-Type Reaction, *J. Am. Chem. Soc.*, 2021, **143**, 8597–8602; (k) R. Bishop, *Comprehensive Organic Synthesis*, ed. P. Knochel and G. A. Molander, 2014, pp. 239–295.