

RESEARCH ARTICLE

View Article Online
View Journal | View IssueCite this: *Org. Chem. Front.*, 2024,
11, 6760Received 30th August 2024,
Accepted 4th October 2024

DOI: 10.1039/d4qo01607f

rsc.li/frontiers-organic

Electrochemical 1,2-hydrogen atom transfer
functionalizations of *N*-(benzyloxy)phthalimides†Xin Fu,^a Tingting Ran^a and Jie Liu ^{*a,b}




We report a straightforward, catalyst-free method for the cathodic benzylic C–H hydroxyalkylation/arylation of *N*-(alkyloxy)phthalimides with ketones or cyanopyridines. The reaction is driven by an electrochemically induced 1,2-hydrogen atom transfer involving alkoxy radicals generated by O–N bond cleavage. This protocol provides an efficient approach for the synthesis of a wide range of valuable unsymmetrical 1,2-diols and diarylmethanols.

Hydrogen atom transfer (HAT) has emerged as a practical and powerful strategy for the C(sp³)-H radical functionalization, which has been widely applied in synthetic organic chemistry.¹ Among the various HAT reactions, heteroatom-centered radicals, such as O[•], N[•], are recognized for their versatility as intermediates in HAT, facilitating the generation of C(sp³) radicals through intermolecular or intramolecular 1,2- or 1,5-HAT reactions (Scheme 1a).^{2,3} For instance, intramolecular 1,5-HAT can effectively produce δ-C(sp³) radicals through the preferred six-membered ring transition-state,⁴ which has been employed in a variety of functionalizations, including heteroarylation,⁵ alkylation,⁶ and allylation.⁷ In contrast, the intramolecular 1,2-HAT reaction is less reported, with only a few examples observed to form C(sp³)-C bonds. This is partly due to the higher activation energy required for the 1,2-HAT processes.⁸ To the best of our knowledge, the initiation of 1,2-HAT processes has been achieved using transition-metal catalysts, including Ir, Ag, Cu, and Ni catalysts,⁹ or through photocatalytic reactions (Scheme 1b).¹⁰ Despite these advances, the development of a mild and sustainable approach to initiate 1,2-HAT processes, especially for the formation of α-C(sp³)-C bonds adjacent to heteroatoms in organic chemistry, is still a worthwhile avenue for investigation.

Organic electrochemistry has been regarded as a sustainable and safe redox reagent that can generate radical species, thereby obviating the need for stoichiometric amounts of oxidants and reductants.¹¹ Recently, a number of chemists have employed an electrochemically driven HAT strategy to achieve

the functionalization of C(sp³)-H bonds.¹² For example, several kinds of different HAT reagents, such as *N*-hydroxyphthalimide (NHPI) derivatives,¹³ quinuclidine,¹⁴ sulfonamide,¹⁵ benzimidazole,¹⁶ and azide,¹⁷ have been reported to promote the intermolecular HAT reactions through anodic oxidation. In addition, several studies have revealed the 1,5-HAT reaction of heteroatom-centered radicals under anodic oxidation.¹⁸ A notable recent example is the electrochemical allylation/alkylation of ethers *via* intermolecular HAT of alkoxy radicals under cathodic reduction, as described by the Bertuzzi and Bandini group.¹⁹ However, the 1,2-HAT reaction of oxygen-centered radicals remains less investigated.²⁰ Therefore, the exploration of electrochemical functionalization of oxygen-

a) 1,5-HAT (well studied) and 1,2-HAT (less studied)

b) The C(sp³)-H functionalization *via* HAT processc) This work: electrochemical cross-coupling *via* 1,2-HATScheme 1 The functionalization of C(sp³)-H *via* HAT process.

^aCollege of Chemistry and Chemical Engineering, State Key Laboratory of Chemo/Biosensing and Chemometrics, Hunan University, 410082 Changsha, China.
E-mail: jieliu@hnu.edu.cn

^bGreater Bay Area Institute for Innovation, Hunan University, 511300 Guangzhou, China

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4qo01607f>

centered α -C(sp³)-H bonds through the 1,2-HAT would significantly broaden the synthetic applications of alkoxy radicals in organic electrochemistry. Herein, we present a simple and metal-free cathodic benzylic C-H hydroxyalkylation/arylation of *N*-(alkoxy)-phthalimides through 1,2-HAT of alkoxy radicals (Scheme 1c). This electrochemically driven HAT strategy offers mild reaction conditions and a wide substrate scope, resulting in the synthesis of diverse 1,2-diols and diarylmethanols.

We initially explored the reaction parameters using benzylic alcohol-derived *N*-alkoxyphthalimide **1a** and benzophenone **2a** as the model substrates (Table 1). Gratifyingly, the reaction yielded the desired 1,2-diol product **3aa** in 78% when conducted in DMAc at a constant current of 8 mA, employing TBAB as the electrolyte in an undivided cell with a magnesium anode and a carbon felt cathode (entry 1). Increasing or decreasing the constant current was found to be less effective and gave slightly lower yields of desired product **3aa** (entry 2). Reducing the reaction time to 3 hours did not improve the yield of product **3aa**, whereas extending it to 5 hours led to a 57% yield (entry 3). Other sacrificial anodes, including Zn, Al and stainless steel, proved less effective than the Mg anode (entry 4). When ⁿBu₄NI, ⁿBu₄NBr or ⁿBu₄NCl were used as electrolytes, significant decreased yields of the desired product **3aa** were obtained (entry 5). Increasing the amount of **1a** to 3.0 equivalents did not improve the yield (entry 6). Notably, no product **3aa** was observed under air conditions, indicating the reaction was air-sensitive (entry 7). Finally, a control experiment confirmed that electricity was essential for the synthesis of 1,2-diols (entry 8).

With the optimal reaction conditions in hand, we probed the substrate scope of *N*-(benzyloxy)phthalimides **1** and ketones **2** (Scheme 2). Using benzophenone **2a** as a coupling

partner, a series of *para*-substituted *N*-(benzyloxy)phthalimides (such as methyl, *i*-propyl, *t*-butyl, phenyl, methoxy, phenoxy, and fluoro substitutions) underwent the electrochemical cross-coupling reaction smoothly, and the desired products **3ba–3ha** were isolated in 27–75% yields. The *ortho*-substituted methyl and fluoro *N*-(benzyloxy)phthalimides were also compatible and gave the desired products **3ia** and **3ja** in 63% and 24% yields, respectively. Moreover, *meta*-substituted *N*-(benzyloxy)phthalimides were efficiently converted to the corresponding 1,2-diols **3ka–3ma**, with yields ranging from 32–70% for substrates containing methyl, methoxy, and fluorine groups. Additionally, *N*-(benzyloxy)phthalimides bearing multiple substituents could also give the desired products **3na–3pa** in 21–46% yields. Notably, the substrate derived from pepper alcohol smoothly participated in the reaction, affording the corresponding product **3qa** in 30% yield. 2-Naphthalene- and thiophen-substituted substrates were also tested and gave the




Table 1 Optimization of reaction conditions^a

Entry	Variation from standard conditions	Yield of 3aa (%)
1	None	78 ^b
2	6 mA or 10 mA instead of 8 mA	66/62
3	3 h or 5 h instead of 3.75 h	52/57
4 ^c	Zn, Al or SS instead of Mg anode	29/0/0
5	ⁿ Bu ₄ NI, ⁿ Bu ₄ NBr, ⁿ Bu ₄ NCl instead of ⁿ Bu ₄ NBF ₄	47/30/37
6	3.0 equivalents of 1a	68
7	Air instead of N ₂	Trace
8	No electricity	0

^a Standard conditions: undivided cell, magnesium (Mg) anode, carbon felt (CF) cathode, constant current = 8 mA, **1a** (2 equiv.), **2a** (0.2 mmol), ⁿBu₄NBF₄ (2 equiv.), DMAc (4 mL), N₂, r.t., 3.75 h. Yield was determined by ¹H NMR analysis using dibromomethane as the internal standard. ^b Isolated yield. ^c Reactions were performed at 10 mA.

Scheme 2 Scope of the electrochemical 1,2-HAT of *N*-(benzyloxy)phthalimides with ketones. Reaction conditions: undivided cell, magnesium (Mg) anode, carbon felt (CF) cathode, constant current = 8 mA, **1** (2 equiv.), **2** (0.2 mmol), ⁿBu₄NBF₄ (2 equiv.), DMAc (4 mL), N₂, r.t., 3.75 h.


corresponding coupling products **3ra** (35%) and **3sa** (49%). Subsequently, we turned our attention to the scope of diarylketone derivatives. Symmetrical diarylketone derivatives, including 4,4'-dimethylbenzophenone, 4,4'-difluorobenzophenone, 9-fluorenone, xanthone, and thioxanthone-9-one, successfully coupled with *N*-(benzyloxy)phthalimide to result the corresponding 1,2-diols **3ab–3af** in 35–72% yields. Furthermore, unsymmetrical diarylketones were employed to form 1,2-diols **3ag–3ak** in 38–68% yields and nearly 1 : 1 diastereoselectivity.

To further demonstrate the diversity of this electrochemical method, we evaluated the feasibility of cyanopyridines as coupling partners in the electrochemical 1,2-HAT coupling, aiming to synthesize heteroaryl-containing diarylmethanols. We were pleased to find that this method could effectively yield a range of heteroaryl-containing diarylmethanols (Scheme 3). For example, the reaction of *N*-(benzyloxy)phthalimide (**1a**) and 4-cyanopyridine (**4a**) with DABCO as the sacrificial reagent led to the desired diarylmethanol product **5aa** in 76% yield. In addition to the simple *N*-(benzyloxy)phthalimide, this process was applicable to a variety of *N*-(benzyloxy)phthalimides bearing electron-rich and electron-poor substituents, affording the desired products **5ba–5ia** in 40–78% yields. Both *meta*- and *ortho*-substituted *N*-(benzyloxy)phthalimides were well-tolerated,


and gave products **5ja–5na** in 23–41% yields. The application of multi-substituted substrates resulted in the corresponding alcohol products **5oa–5ra** in 44–52% yields. Furthermore, the use of naphthalene- or heterocyclic-substituted *N*-(benzyloxy)phthalimides as coupling partners delivered products **5sa–5va** in 36–65% yields. On the other hand, a series of cyanopyridines smoothly underwent the electroreductive arylation, providing the desired products **5ab–5af** in 38–83% yields. Additionally, 1,4-dicyanobenzene could also take part in this reaction to form the product **5ag**, albeit in a lower yield of 20%.

Next, we evaluated the scalability of the electroreductive cross-coupling strategy by conducting the reaction on a 5 mmol scale, employing **1a** and **2a**, or **4a** as substrates. The corresponding products **3a** and **5a**, were successfully obtained with yields of 52% and 46%, respectively (Scheme 4). These results highlight the practicality and efficacy of this method for the synthesis of 1,2-diols and diarylmethanols under electrochemical conditions.


To understand this reaction, a series of mechanistic experiments were investigated (Scheme 5). The addition of a free radical inhibitor (TEMPO) to the standard reaction led to significant decreased or no yields of the desired product (Scheme 5a). These results indicated that the electrochemical hydroxyalkylation and arylation processes proceed *via* a radical pathway. In addition, the application of benzyl alcohol **6a** instead of *N*-alkoxyphthalimide **1a** led to no arylation product **5aa**, suggesting the alcohol is not related in this transformation. Meanwhile, the reaction of benzaldehyde **6b** and cyanopyridine **4a** resulted in a 32% yield of dialymethanol **5aa**, suggesting a similar reaction pathway through a ketyl radical intermediate (Scheme 5b).²¹ Moreover, the EPR analysis of **1a** with DMPO under electroreduction conditions indicated the production of a DMPO-captured alkoxy radical **7a** and ketyl radical **7b** (Scheme 5c).^{10d} The use of the *N*-alkoxyphthalimide salt derivative also afforded the α -C–H arylation adduct in 38% yield (see ESI, Scheme S1†).^{5c,d,22} Furthermore, the kinetic isotope effect (KIE) experiments of hydroxyalkylation and arylation were observed with **1a** and **D-1a**, in which the k_H/k_D ratios were 0.76 and 1.14, respectively (see ESI, Fig. S7 and S8†). These results confirm the intermediacy of radicals species and intramolecular 1,2-HAT process in this transformation.



Scheme 3 Scope of the electrochemical 1,2-HAT of *N*-(benzyloxy)phthalimides with cyanopyridines. Reaction conditions: undivided cell, platinum plate (Pt) anode, carbon felt (CF) cathode, constant current = 10 mA, **1** (0.2 mmol), **4** (2 equiv.), DABCO (3 equiv.), ^tBu₄NBF₄ (2 equiv.), DMSO (4 mL), air, r.t., 3 h.




Scheme 4 Scale-up experiments.



Scheme 5 Mechanistic experiments.

Based on these mechanistic studies and previous reports,^{10b–e,23} we propose a possible mechanism for this electrochemical α -C(sp³)-H hydroxyalkylation/arylation of *N*-(alkoxy)phthalimides (Scheme 6). The cathodic reduction of *N*-(benzyloxy)phthalimide **1a** results in the formation of the phthalimide anion and a key benzyloxy radical **A**. This radical subsequently undergoes 1,2-HAT to generate the ketyl radical **B**. In the hydroxyalkylation process, the benzophenone **2a** gains a single electron to form radical anion **C** at the cathode. Subsequently, intermediate **B** and **C** undergo radical–radical cross-coupling to produce anion intermediate **D**, followed by



Scheme 6 Proposed mechanism.

protonation to form the final product 1,2-diol **3aa**. Meanwhile, Mg is oxidized to Mg²⁺ at the anode. In the arylation process, the 4-cyanopyridine **4a** is reduced *via* a single-electron transfer (SET) process to generate a radical anion **E** on the cathode. This intermediate **E** couples with the above intermediate **B** to produce anion intermediate **F** and then loss of CN[−] to obtain the desired diarylmethanol **5aa**. In this transformation, DABCO works as a sacrificing agent at the anode.

Conclusions

In conclusion, we have described a straightforward and efficient electrochemical approach for the cathodic benzylic C–H hydroxyalkylation and arylation of *N*-(alkoxy)phthalimides, mediated by the 1,2-HAT of alkoxy radicals. This method provides access to a range of valuable 1,2-diols and diarylmethanols in moderate to good yields under mild and straightforward electrolysis conditions. We anticipate that this versatile and efficient electrochemical strategy will broaden the synthetic applications of alkoxy radicals in organic chemistry.

Data availability

The data supporting this article have been included as part of the ESI.†

Conflicts of interest

The authors declare no competing financial interest.

Acknowledgements

The authors are grateful for the generous financial support provided by the Natural Science Foundation of China (22301073) and the Natural Science Foundation of Hunan Province (2021RC3056) and the Fundamental Research Funds for the Central Universities.

References

- For reviews about hydrogen atom transfer, see: (a) J. M. Mayer, Understanding hydrogen atom transfer: from bond strengths to marcus theory, *Acc. Chem. Res.*, 2011, **44**, 36–46; (b) S. Sarkar, K. P. S. Cheung and V. Gevorgyan, C–H functionalization reactions enabled by hydrogen atom transfer to carbon-centered radicals, *Chem. Sci.*, 2020, **11**, 12974–12993; (c) L. Capaldo, D. Ravelli and M. Fagnoni, Direct photocatalyzed hydrogen atom transfer (HAT) for aliphatic C–H bonds elaboration, *Chem. Rev.*, 2021, **122**, 1875–1924; (d) D. L. Golden, S.-E. Suh and S. S. Stahl, Radical C(sp³)-H functionalization and cross-coupling reactions, *Nat. Rev. Chem.*, 2022, **6**, 405–427;

- (e) J. Zhang and M. Rueping, Metallaphotoredox catalysis for sp^3 C–H functionalizations through hydrogen atom transfer (HAT), *Chem. Soc. Rev.*, 2023, **52**, 4099–4120.
- 2 For reviews see: (a) M. Salamone and M. Bietti, Tuning reactivity and selectivity in hydrogen atom transfer from aliphatic C–H bonds to alkoxy radicals: role of structural and medium effects, *Acc. Chem. Res.*, 2015, **48**, 2895–2903; (b) K. Kwon, R. T. Simons, M. Nandakumar and J. L. Roizen, Strategies to generate nitrogen-centered radicals that may rely on photoredox catalysis: development in reaction methodology and applications in organic synthesis, *Chem. Rev.*, 2022, **122**, 2353–2428; (c) X. Wu and C. Zhu, Radical functionalization of remote $C(sp^3)$ –H bonds mediated by unprotected alcohols and amides, *CCS Chem.*, 2020, **2**, 813–828; (d) W. Guo, Q. Wang and J. Zhu, Visible light photoredox-catalysed remote C–H functionalisation enabled by 1,5-hydrogen atom transfer (1,5-HAT), *Chem. Soc. Rev.*, 2021, **50**, 7359–7377; (e) K. Jia and Y. Chen, Visible-light-induced alkoxy radical generation for inert chemical bond cleavage/functionalization, *Chem. Commun.*, 2018, **54**, 6105–6112; (f) J.-J. Guo, A. Hu and Z. Zuo, Photocatalytic alkoxy radical-mediated transformations, *Tetrahedron Lett.*, 2018, **59**, 2103; (g) L. Kong, Y. Zhou, F. Luo and G. Zhu, Recent advances on oxidative radical addition to aldehydes, *Chin. J. Org. Chem.*, 2018, **38**, 2858–2865; (h) X. Wu and C. Zhu, Recent advances in radical-mediated C–C Bond fragmentation of non-strained molecules, *Chin. J. Chem.*, 2019, **37**, 171–182; (i) X. Wu and C. Zhu, Recent advances in alkoxy radical-promoted C–C and C–H bond functionalization starting from free alcohols, *Chem. Commun.*, 2019, **55**, 9747–9756; (j) C. Liang, Q. An, L. Duan, K. Feng and Z. Zuo, Alkoxy radicals see the light: new paradigms of photochemical synthesis, *Chem. Rev.*, 2021, **122**, 2429–2486.
- 3 For representative papers, see: (a) A. E. Dorigo and K. N. Houk, The relationship between proximity and reactivity. An ab initio study of the flexibility of the OH \cdots H + CH₄ hydrogen abstraction transition state and a force-field model for the transition states of intramolecular hydrogen abstractions, *J. Org. Chem.*, 1988, **53**, 1650–1664; (b) J. Hartung, Stereoselective construction of the tetrahydrofuran nucleus by alkoxy radical cyclizations, *Eur. J. Org. Chem.*, 2001, 619–632; (c) S. Chiba and H. Chen, sp^3 C–H oxidation by remote H-radical shift with oxygen- and nitrogen-radicals: a recent update, *Org. Biomol. Chem.*, 2014, **12**, 4051–4060; (d) H. Zhu, J. G. Wickenden, N. E. Campbell, J. C. Leung, K. M. Johnson and G. M. Sammis, Construction of carbo- and heterocycles using radical relay cyclizations initiated by alkoxy radicals, *Org. Lett.*, 2009, **11**, 2019–2022; (e) A. Hu, J.-J. Guo, H. Pan and Z. Zuo, Selective functionalization of methane, ethane, and higher alkanes by cerium photocatalysis, *Science*, 2018, **361**, 668–672; (f) A. Hu, J. J. Guo, H. Pan, H. Tang, Z. Gao and Z. Zuo, δ -Selective functionalization of alkanols enabled by visible-light-induced ligand-to-metal charge transfer, *J. Am. Chem. Soc.*, 2018, **140**, 1612–1616; (g) H. Guan, S. Sun, Y. Mao, L. Chen, R. Lu, J. Huang and L. Liu, Iron(II)-Catalyzed site-selective functionalization of unactivated $C(sp^3)$ –H bonds guided by alkoxy radicals, *Angew. Chem., Int. Ed.*, 2018, **57**, 11413–11417.
- 4 (a) A. W. Hofmann, Ueber die einwirkung des broms in alkalischer lösung auf die amine, *Ber. Dtsch. Chem. Ges.*, 1883, **16**, 558–560; (b) D. H. R. Barton, J. M. Beaton, L. E. Geller and M. M. Pechet, A new photochemical reaction, *J. Am. Chem. Soc.*, 1960, **82**, 2640–2641; (c) D. Moran, R. Jacob, G. P. F. Wood, M. L. Coote, M. J. Davies, R. O'Hair, C. J. Easton and L. Radom, Rearrangements in model peptide-type radicals via intramolecular hydrogen-atom transfer, *Helv. Chim. Acta*, 2006, **89**, 2254–2272.
- 5 For recent selected examples of heteroarylation, see: (a) X. Wu, M. Wang, L. Huan, D. Wang, J. Wang and C. Zhu, Tertiary-alcohol-directed functionalization of remote $C(sp^3)$ –H bonds by sequential hydrogen atom and heteroaryl migrations, *Angew. Chem., Int. Ed.*, 2018, **57**, 1640–1644; (b) X. Wu, H. Zhang, N. Tang, Z. Wu, D. Wang, M. Ji, Y. Xu, M. Wang and C. Zhu, Metal-free alcohol-directed regioselective heteroarylation of remote unactivated $C(sp^3)$ –H bonds, *Nat. Commun.*, 2018, **9**, 3343–3350; (c) I. Kim, B. Park, G. Kang, J. Kim, H. Jung, H. Lee, M. Baik and S. Hong, Visible-light-induced pyridylation of remote $C(sp^3)$ –H bonds by radical translocation of *N*-alkoxy-pyridinium salts, *Angew. Chem., Int. Ed.*, 2018, **57**, 15517–15522; (d) X. Bao, Q. Wang and J. Zhu, Remote $C(sp^3)$ –H arylation and vinylation of *N*-alkoxy-pyridinium salts to δ -aryl and δ -vinyl alcohols, *Chem. – Eur. J.*, 2019, **25**, 11630–11634; (e) G.-X. Li, X. Hu, G. He and G. Chen, Photoredox-mediated remote $C(sp^3)$ –H heteroarylation of free alcohols, *Chem. Sci.*, 2019, **10**, 688–693; (f) H. Chen, W. Fan, X.-A. Yuan and S. Yu, Site-selective remote $C(sp^3)$ –H heteroarylation of amides via organic photoredox catalysis, *Nat. Commun.*, 2019, **10**, 4743–4751; (g) C. Wang, Z. Chen, J. Sun, L. Tong, W. Wang, S. Song and J. Li, Sulfonamide-directed site-selective functionalization of unactivated $C(sp^3)$ –H enabled by photocatalytic sequential electron/proton transfer, *Nat. Commun.*, 2024, **15**, 5087–5097.
- 6 For recent selected examples of alkylation, see: (a) S. M. Thullen, S. M. Treacy and T. Rovis, Regioselective alkylation cross-coupling of remote unactivated $C(sp^3)$ –H bonds, *J. Am. Chem. Soc.*, 2019, **141**, 14062–14067; (b) L.-J. Zhong, G.-F. Lv, X.-H. Ouyang, Y. Li and J.-H. Li, Copper-catalyzed fluoroamide-directed remote benzylic C–H olefination: facile access to internal alkenes, *Org. Chem. Front.*, 2022, **9**, 4309–4315; (c) Z. Du, S. Liu, Y. Li, J. Peng, Y. Sun, Y. Song, Y. Liu and X. Zeng, Fluoroamide-directed regiodivergent C-alkylation of nitroalkanes, *Org. Lett.*, 2023, **25**, 4632–4637.
- 7 For recent selected examples of allylation, see: (a) J. Zhang, Y. Li, F. Zhang, C. Hu and Y. Chen, Generation of alkoxy radicals by photoredox catalysis enables selective $C(sp^3)$ –H functionalization under mild reaction conditions, *Angew. Chem., Int. Ed.*, 2016, **55**, 1872–1875; (b) B. Xu and

- U. K. Tambar, Remote allylation of unactivated C(sp³)-H bonds triggered by photogenerated amidyl radicals, *ACS Catal.*, 2019, **9**, 4627–4631; (c) K. Wu, L. Wang, S. Colón-Rodríguez, G.-U. Flechsig and T. Wang, Amidyl radical directed remote allylation of unactivated sp³ C–H bonds by organic photoredox catalysis, *Angew. Chem., Int. Ed.*, 2019, **58**, 1774–1778; (d) Y. Deng, M. D. Nguyen, Y. Zou, K. N. Houk and A. B. Smith III, Generation of dithianyl and dioxolanyl radicals using photoredox catalysis: application in the total synthesis of the danshenspiroketallactones via radical relay chemistry, *Org. Lett.*, 2019, **21**, 1708–1712.
- 8 (a) S. Saebo, L. Radom and H. F. Schaefer, The weakly exothermic rearrangement of methoxy radical (CH₃O[•]) to the hydroxymethyl radical (CH₂OH[•]), *J. Chem. Phys.*, 1983, **78**, 845–853; (b) L. A. Curtiss, L. D. Kock and J. A. Pople, Energies of CH₂OH, CH₃O, and related compounds, *J. Chem. Phys.*, 1991, **95**, 4040–4043.
- 9 (a) C. Che, Q. Huang, H. Zheng and G. Zhu, Copper-catalyzed cascade annulation of unsaturated α -bromocarbonyls with enynals: a facile access to ketones from aldehydes, *Chem. Sci.*, 2016, **7**, 4134–4139; (b) Z. Liu, Y. Bai, J. Zhang, Y. Yu, Z. Tan and G. Zhu, Copper-catalyzed acyltrifluoromethylation of alkenes: rapid access to trifluoroethyl indanones and related compounds, *Chem. Commun.*, 2017, **53**, 6440–6443; (c) D. Lu, Y. Wan, L. Kong and G. Zhu, Visible-Light-Induced Tandem Radical Addition–Cyclization of Alkenyl Aldehydes Leading to Indanones and Related Compounds, *Org. Lett.*, 2017, **19**, 2929–2932; (d) C. Che, Z. Qian, M. Wu, Y. Zhao and G. Zhu, Intermolecular oxidative radical addition to aromatic aldehydes: direct access to 1,4- and 1,5-diketones via silver-catalyzed ring-opening acylation of cyclopropanols and cyclobutanols, *J. Org. Chem.*, 2018, **83**, 5665–5673; (e) L. Lombardi, A. Cerveri, R. Giovanelli, M. C. Reis, C. S. Lopez, G. Bertuzzi and M. Bandini, Direct synthesis of α -aryl- α -trifluoromethyl alcohols via nickel catalyzed cross-electrophile coupling, *Angew. Chem., Int. Ed.*, 2022, **61**, e202211732; (f) Y. Jiang, D. Liu, M. E. Rotella, G. Deng, Z. Liu, W. Chen, H. Zhang, M. C. Kozłowski, P. J. Walsh and X. Yang, Net-1,2-hydrogen atom transfer of amidyl radicals: toward the synthesis of 1,2-diamine derivatives, *J. Am. Chem. Soc.*, 2023, **145**, 16045–16057.
- 10 (a) C. Wang, Y. Yu, W.-L. Liu and W.-L. Duan, Site-Tunable Csp³-H Bonds Functionalization by Visible-Light-Induced Radical Translocation of *N*-Alkoxyphthalimides, *Org. Lett.*, 2019, **21**, 9147–9152; (b) L. Niu, J. Liu, X. Liang, S. Wang and A. Lei, Visible light-induced direct α C–H functionalization of alcohols, *Nat. Commun.*, 2019, **10**, 467; (c) L. J. Zhong, H. Y. Wang, X. H. Ouyang, J. H. Li and D. L. An, Benzylic C-H heteroarylation of *N*-(benzyloxy) phthalimides with cyanopyridines enabled by photoredox 1,2-hydrogen atom transfer, *Chem. Commun.*, 2020, **56**, 8671–8674; (d) J. Zhang, D. Liu, S. Liu, Y. Ge, Y. Lan and Y. Chen, Visible-light-induced alkoxy radicals enable α -C(sp³)-H bond allylation, *iScience*, 2020, **23**, 100755; (e) D. Liu, J. Zhang and Y. Chen, Investigations on the 1,2-hydrogen atom transfer reactivity of alkoxy radicals under visible-light-induced reaction conditions, *Synlett*, 2021, 356–361; (f) F. Chen, X.-H. Xu, L. Chu and F.-L. Qing, Visible-light-induced nickel-catalyzed radical cross-couplings to access α -aryl- α -trifluoromethyl alcohols, *Org. Lett.*, 2022, **24**, 9332–9336; (g) B. Chen, Q. Chen, Y. Liu, J. Chen, X. Zhou, H. Wang, Q. Yan, W. Wang, Z. Cai and F.-E. Chen, Visible-light-induced defluorinative α -C(sp³)-H alkylation for the synthesis of gem-difluoroallylated α -trifluoromethylamines, *Org. Lett.*, 2023, **25**, 9124–9129; (h) J. J. Monteith and S. A. L. Rousseaux, A dual Ni/photoredox cross-coupling approach toward mandelic acids, *Org. Lett.*, 2024, **26**, 4566–4570; (i) Z. Wang, E. Tang, Q. Zhou and J.-P. Wan, Metal-free photocatalyzed homo- and cross-dimerizations of α -hydroxyl ketones via dual C(sp³)-H functionalization: synthesis of 2,3-dihydroxy-1,4-butanediones, *Org. Chem. Front.*, 2024, **11**, 1157–1162; (j) E. Tang, Q.-Q. Zhou and J.-P. Wan, Visible-light-induced Ritter-type amidation of α -hydroxy ketones in the selective synthesis of α,α -diamido and monoamido ketones, *Chem. Commun.*, 2024, **60**, 7471–7474; (k) Z. Wang, L. Gan, Z. Song, Y. Liu and J.-P. Wan, ^tBuOK/TMSOK-mediated “alkyl halide to alkyl free radical” transformation for transition-metal-free benzoin α -C–H alkylation, *Chin. J. Chem.*, 2024, **42**, 3041–3046; (l) J. Caner, A. Matsumoto and K. Maruoka, Facile synthesis of 1,2-aminoalcohols via α -C–H aminoalkylation of alcohols by photoinduced hydrogen-atom transfer catalysis, *Chem. Sci.*, 2023, **14**, 13879–13884.
- 11 (a) A. Jutan, Contribution of electrochemistry to organometallic catalysis, *Chem. Rev.*, 2008, **108**, 2300–2347; (b) B. A. Frontana-Urbe, R. D. Little, J. G. Ibanez, A. Palma and R. Vasquez-Medrano, Organic electrosynthesis: a promising green methodology in organic chemistry, *Green Chem.*, 2010, **12**, 2099–2119; (c) H. J. Schäfer, Contributions of organic electrosynthesis to green chemistry, *C. R. Chim.*, 2011, **14**, 745–765; (d) R. Francke and R. D. Little, Redox catalysis in organic electrosynthesis: basic principles and recent developments, *Chem. Soc. Rev.*, 2014, **43**, 2492–2521; (e) E. J. Horn, B. R. Rosen and P. S. Baran, Synthetic organic electrochemistry: an enabling and innately sustainable method, *ACS Cent. Sci.*, 2016, **2**, 302–308; (f) M. Yan, Y. Kawamata and P. S. Baran, Synthetic organic electrochemical methods since 2000: on the verge of a renaissance, *Chem. Rev.*, 2017, **117**, 13230–13319; (g) S. Tang, Y. Liu and A. Lei, Electrochemical oxidative cross-coupling with hydrogen evolution: a green and sustainable way for bond formation, *Chem*, 2018, **4**, 27–45; (h) Y. Jiang, K. Xu and C. Zeng, Use of electrochemistry in the synthesis of heterocyclic structures, *Chem. Rev.*, 2018, **118**, 4485–4540; (i) Q.-L. Yang, P. Fang and T.-S. Mei, Recent advances in organic electrochemical C–H functionalization, *Chin. J. Chem.*, 2018, **36**, 338–352; (j) L. Zeng, Q. Yang, J. Wang, X. Wang, P. Wang, S. Wang, S. Lv, S. Muhammad, Y. Liu, H. Yi and A. Lei, Programmed alternating current optimization of Cu-catalyzed C–H bond transformations, *Science*, 2024, **385**, 216–223.

- 12 (a) M. D. Kärkäs, Electrochemical strategies for C–H functionalization and C–N bond formation, *Chem. Soc. Rev.*, 2018, **47**, 5786–5865; (b) P. R. D. Murray, J. H. Cox, N. D. Chiappini, C. B. Roos, E. A. McLoughlin, B. G. Hejna, S. T. Nguyen, H. H. Ripberger, J. M. Ganley, E. Tsui, N. Y. Shin, B. Koronkiewicz, G. Qiu and R. R. Knowles, Photochemical and electrochemical applications of proton-coupled electron transfer in organic synthesis, *Chem. Rev.*, 2022, **122**, 2017–2291; (c) S. Zhang and M. Findlater, Electrochemically driven hydrogen atom transfer catalysis: a tool for C(sp³)/Si–H functionalization and hydrofunctionalization of alkenes, *ACS Catal.*, 2023, **13**, 8731–8751; (d) Z. Yang, W. Shi, H. Alhumade, H. Yi and A. Lei, Electrochemical oxidative C(sp³)-H cross-coupling with hydrogen evolution, *Nat. Synth.*, 2023, **2**, 217–230; (e) P. Xu, P.-Y. Chen and H.-C. Xu, Scalable photoelectrochemical dehydrogenative cross-coupling of heteroarenes with aliphatic C–H bonds, *Angew. Chem., Int. Ed.*, 2020, **59**, 14275–14280; (f) Y. Jiang, K. Xu and C. Zeng, Electrophotocatalytic Si–H activation governed by polarity-matching effects, *CCS Chem.*, 2021, **4**, 1796–1805; (g) P.-F. Zhong, J.-L. Tu, Y. Zhao, N. Zhong, C. Yang, L. Guo and W. Xia, Photoelectrochemical oxidative C(sp³)-H borylation of unactivated hydrocarbons, *Nat. Commun.*, 2023, **14**, 6530–6540; (h) L. Zou, X. Wang, S. Xiang, W. Zheng and Q. Lu, Paired oxidative and reductive catalysis: breaking the potential barrier of electrochemical C(sp³)-H alkenylation, *Angew. Chem., Int. Ed.*, 2023, **62**, e202301026.
- 13 (a) F. Wang and S. S. Stahl, Electrochemical oxidation of organic molecules at lower overpotential: accessing broader functional group compatibility with electron–proton transfer mediators, *Acc. Chem. Res.*, 2020, **53**, 561–574; (b) E. J. Horn, B. R. Rosen, Y. Chen, J. Tang, K. Chen, M. D. Eastgate and P. S. Baran, Scalable and sustainable electrochemical allylic C–H oxidation, *Nature*, 2016, **533**, 77–81; (c) M. A. Hoque, J. Twilton, J. Zhu, M. D. Graaf, K. C. Harper, E. Tuca, G. A. DiLabio and S. S. Stahl, Electrochemical PINOylation of methylarenes: improving the scope and utility of benzylic oxidation through mediated electrolysis, *J. Am. Chem. Soc.*, 2022, **144**, 15295–15302; (d) H. Liang, L.-J. Wang, Y.-X. Ji, H. Wang and B. Zhang, Selective electrochemical hydrolysis of hydrosilanes to silanols *via* anodically generated silyl cations, *Angew. Chem., Int. Ed.*, 2021, **60**, 1839–1844; (e) J. Ke, W. Liu, X. Zhu, X. Tan and C. He, Electrochemical radical silyloxygenation of activated alkenes, *Angew. Chem., Int. Ed.*, 2021, **60**, 8744–8749.
- 14 Y. Kawamata, M. Yan, Z. Liu, D.-H. Bao, J. Chen, J. T. Starr and P. S. Baran, Scalable, electrochemical oxidation of unactivated C–H bonds, *J. Am. Chem. Soc.*, 2017, **139**, 7448–7451.
- 15 Y. Liu, B. Shi, Z. Liu, R. Gao, C. Huang, H. Alhumade, S. Wang, X. Qi and A. Lei, Time-resolved EPR revealed the formation, structure, and reactivity of *N*-centered radicals in an electrochemical C(sp³)-H arylation reaction, *J. Am. Chem. Soc.*, 2021, **143**, 20863–20872.
- 16 Y. Liang, X. Zhan, F. Li, H. Bi, W. Fan, S. Zhang and M.-B. Li, Using a nitrogen-centered radical as a selective mediator in electrochemical C(sp³)-H amination, *Chem. Catal.*, 2023, **3**, 100582.
- 17 (a) H. Li, J. W. Tong, Y. Zhu, C. Jiang, P. Liu and P. Sun, Electrochemical Minisci reaction *via* HAT-driven α -C(sp³)-H functionalization of alcohols, *Green Chem.*, 2022, **24**, 8406–8411; (b) H. Li, C. Qiang, Y. Liao, Q. Chu, P. Liu and P. Sun, HAT-mediated electrochemical cross dehydrogenative coupling reaction of quinoxalin-2(1*H*)-ones with ethers, *Adv. Synth. Catal.*, 2023, **365**, 3149–3154.
- 18 (a) S. Herold, D. Bafaluy and K. Muñiz, Anodic benzylic C(sp³)-H amination: unified access to pyrrolidines and piperidines, *Green Chem.*, 2018, **20**, 3191–3196; (b) X. Hu, G. Zhang, F. Bu, L. Nie and A. Lei, Electrochemical oxidation-induced site-selective intramolecular C(sp³)-H amination, *ACS Catal.*, 2018, **8**, 9370–9375; (c) P. Nikolaienko, M. Jentsch, A. P. Kale, Y. Cai and M. Rueping, Electrochemical and scalable dehydrogenative C(sp³)-H amination *via* remote hydrogen atom transfer in batch and continuous flow, *Chem. – Eur. J.*, 2019, **25**, 7177–7184; (d) J. E. Hong, J. Yoon, W. Baek, K. Kim, J.-H. Kwak and Y. Park, Electrochemical C(sp³)-H lactonization of 2-alkylbenzoic acids toward phthalides, *Org. Lett.*, 2023, **25**, 298–303; (e) X. Pang, H. He, X. Meng, L. Zhang, S. Ni, M. Li and W. Guo, Electrochemical remote C(sp³)-H thiocyanation, *Org. Chem. Front.*, 2024, **11**, 2283–2288.
- 19 L. Rapisarda, A. Fermi, P. Ceroni, R. Giovanelli, G. Bertuzzi and M. Bandini, Electrochemical C(sp³)-H functionalization of ethers *via* hydrogen-atom transfer by means of cathodic reduction, *Chem. Commun.*, 2023, **59**, 2664–2667.
- 20 A. A. M. A. El Gehani, H. A. Maashi, J. Harnedy and L. C. Morrill, Electrochemical generation and utilization of alkoxy radicals, *Chem. Commun.*, 2023, **59**, 3655–3664.
- 21 (a) Á. Péter, S. Agasti, O. Knowles, E. Pye and D. J. Procter, Recent advances in the chemistry of ketyl radicals, *Chem. Soc. Rev.*, 2021, **50**, 5349–5365; (b) X. Zhang, C. Yang, H. Gao, L. Wang, L. Guo and W. Xia, Reductive arylation of aliphatic and aromatic aldehydes with cyanoarenes by electrolysis for the synthesis of alcohols, *Org. Lett.*, 2021, **23**, 3472–3476; (c) L.-J. Wang, P. Ye, N. Tan and B. Zhang, Electroreductive cross-coupling between aldehydes and ketones or imines *via* cathodically generated dianions, *Green Chem.*, 2022, **24**, 8386–8392.
- 22 (a) A. L. Barthelemy, B. Tuccio, E. Magnier and G. Dagousset, Alkoxy radicals generated under photoredox catalysis: a strategy for anti-Markovnikov alkoxylation reactions, *Angew. Chem., Int. Ed.*, 2018, **57**, 13790–13794; (b) X. Bao, Q. Wang and J. Zhu, Dual photoredox/copper catalysis for the remote C(sp³)-H functionalization of alcohols and alkyl halides by *N*-alkoxy pyridinium salts, *Angew. Chem., Int. Ed.*, 2019, **58**, 2139–2143; (c) Y. Kim, K. Lee, G. R. Mathi, I. Kim and S. Hong, Visible-light-induced cascade radical ring-closure and pyridylation for the synthesis of tetrahydrofurans, *Green Chem.*, 2019, **21**, 2082–2087.

- 23 (a) B. C. Gilbert, R. G. G. Holmes, H. A. H. Laue and R. O. Norman, Electron spin resonance studies. Part L. Reactions of alkoxy radicals generated from alkyl hydroperoxides and titanium(III) ion in aqueous solution, *J. Chem. Soc., Perkin Trans. 2*, 1976, 1047–1052; (b) P. E. Elford and B. P. Roberts, EPR studies of the formation and transformation of isomeric radicals $[C_3H_5O]^{\cdot}$. Rearrangement of the allyloxy radical in non-aqueous solution involving a formal 1,2-hydrogen-atom shift promoted by alcohols, *J. Chem. Soc., Perkin Trans. 2*, 1996, 2247–2256; (c) E. Baciocchi, M. Bietti and S. Steenken, Base-catalyzed C–H deprotonation of 4-methoxybenzyl alcohol radical cations in water: evidence for a carbon-to-oxygen 1,2-H-shift mechanism, *J. Am. Chem. Soc.*, 1997, **119**, 4078–4079; (d) G. Wang, J. Cao, L. Gao, W. Chen, W. Huang, X. Cheng and S. Li, Metal-free synthesis of C-4 substituted pyridine derivatives using pyridine-boryl radicals via a radical addition/coupling mechanism: a combined computational and experimental study, *J. Am. Chem. Soc.*, 2017, **139**(10), 3904–3910; (e) W. Ding, M. Li, J. Fan and X. Cheng, Palladium-catalyzed asymmetric allylic 4-pyridinylation via electroreductive substitution reaction, *Nat. Commun.*, 2022, **13**, 5642; (f) W. Ding, J. Sheng, J. Li and X. Cheng, Electroreductive 4-pyridylation of unsaturated compounds using gaseous ammonia as a hydrogen source, *Org. Chem. Front.*, 2022, **9**, 2634–2639.