


 Cite this: *RSC Adv.*, 2024, 14, 5542

 Received 23rd January 2024
 Accepted 31st January 2024

DOI: 10.1039/d4ra00589a

rsc.li/rsc-advances

First total synthesis of caerulomycin K: a case study on selective, multiple C–H functionalizations of pyridines†

 Alessandro Dimasi,[‡] Mattia Failla,[‡] Arianna Montoli,^a Andrea Citarella,^a Paolo Ronchi,[‡] Daniele Passarella,^a and Valerio Fasano^{*,a}

Caerulomycins, natural alkaloids with antimicrobial properties, have been previously synthesized starting with highly pre-functionalized building blocks or requiring many functional group manipulations. In this work, we report the first total synthesis of caerulomycin K, a diversely trifunctionalized pyridine readily assembled in three steps exploiting the recent advancements in the C–H activation of N-heterocycles.

Pyridines are ubiquitous in many natural products and drugs, often with a wide selection of functionalities decorating these aromatic rings.^{1,2} While classical pyridine syntheses (*e.g.* Bohlmann–Rahtz reaction, Hantzsch condensation, *etc.*) allow the introduction of substituents in the final ring, the functionalization of existing pyridines using C–H activation is usually a better option to avoid the *de novo* synthesis of complex pyridines.³ However, despite pyridines exhibiting a clear similarity to benzenes, they present distinct challenges when it comes to their C–H functionalization.^{4,5} As a result, relatively simple pyridines may require several steps to be synthesized, especially if the substituents around the aromatic ring are different in nature. This is the case for caerulomycins (and related collismycins), a class of natural alkaloids produced by *Streptomyces caeruleus* and endowed with antimicrobial properties (Scheme 1A).^{6–11} For instance, taking caerulomycin E as the prototype of this type of bioactive compounds, different routes have been designed to decorate the core pyridine ring with common substituents such as a carbonyl group (*ortho*-aldehyde), an alkoxide (*para*-MeO), and an aromatic ring (*ortho*-pyridine).^{12–21} Yet, the installation of these functionalities *via* C–H activation is not straightforward since it requires several functional group interconversions (Scheme 1B).


Specifically, the installation of the methoxy group can require four steps: formation of the N-oxide with an oxidant, nitration with concentrated H₂SO₄, nucleophilic aromatic

substitution (S_NAr) with MeONa, and removal of the oxide with Ac₂O.^{18,19} The insertion of an *ortho*-pyridine group is usually more rapid but requires the use of Grignard reagents or pre-functionalized 2-bromopyridines.^{15,16} Finally, the insertion of the carbonyl group is achieved by oxidation of a methyl group whose installation has been obtained only with a halogen (Cl or Br) already placed in *ortho*-position.^{18,21} Given the recent advancements in selective C–H functionalizations of pyridines,^{4,5} it would be expected that alternative strategies should now allow a faster synthesis of caerulomycins. Herein, we report our efforts to rapidly convert a cheap monosubstituted pyridine into caerulomycin K, a recently isolated alkaloid whose total synthesis has never been reported before.


A) Caerulomycins' structures

	Name	Ar	R	R ₁	X
	Caerulomycin A	Py	OMe	-	N-OH
	Caerulomycin B	Py	OMe	OH	N-OH
	Caerulomycin E	Py	OMe	-	O
	Caerulomycin K	Ph	OMe	-	N-OH
	Collismycin A	Py	OMe	SMe	N-OH
	Collismycin D	Py	OH	SMe	N-OH

B) Previously reported approaches to caerulomycin E



C) First total synthesis of caerulomycin K (this work)



Scheme 1 Structures and synthetic routes to caerulomycins.

^aDepartment of Chemistry, Università degli Studi di Milano, Via Camillo Golgi, 19, 20133 Milano, Italy. E-mail: valerio.fasano@unimi.it; Web: <https://www.fasanolab.com>

^bMedicinal Chemistry and Drug Design Technologies Department, Global Research and Preclinical Development, Chiesi Farmaceutici S.p.A, Largo Francesco Belloli 11/a, 43126 Parma, Italy


† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4ra00589a>

‡ Equal contribution.



Finally, attempts to use a more nucleophilic phosphine (*i.e.* (4-anisyl)₃P) did not improve the C–P bond formation, and neither did the use of it as the first installed phosphine (see ESI†). Indeed, the use of (4-anisyl)₃P mainly resulted in the formation of the corresponding phosphine oxide, as expected for electron-rich phosphines. Given the problematic separation of salts **2** and **3** and the modest conversion observed in the second step, we decided to perform one C–P bond formation at a time. Treatment of **2** with MeONa in dichloromethane gave disubstituted pyridine **7** in 53% ¹H NMR yield (Scheme 5), although its isolation was complicated by co-eluting Ph₃PO (the product, together with **1**, of competitive protodephosphination). Before performing the second C–P bond formation on **7**, the installation of a methyl group using **2** as a model compound was attempted.


Indeed, while the replacement of Ph₃P with chalcogens/pnictogen nucleophiles (–OR, –SR, –NR₂) is relatively straightforward,^{27,30,31} the installation of alkyl or aryl groups *via* ligand-coupling requires additional manipulations.^{32–34} However, the direct use of organolithium has been shown successful in a couple of cases (*e.g.* ArLi), hence we hoped the use of MeLi or MeMgBr would avoid extra steps.²⁷ Unfortunately, treatment of **2** with these organometallics provided equimolar amounts of 2,4-diphenylpyridine **8** and Ph₂P(O)Me (Scheme 5), the latter observable in the ¹H NMR spectrum (2.01 ppm, d, ²J_{P–H} = 13.2 Hz, 3H, Me). This result highlights how, in contrast to alkoxides, the phenyl ring has a higher migration aptitude than a methyl group during the ligand-coupling of the phosphorane intermediate,^{35,36} thus leading to **8** and Ph₂PMe (then oxidized during the workup). An alternative approach to the use of phosphonium salts would be an *ortho*-halogenation followed by Negishi coupling with MeZnCl, in analogy with the reported synthesis of caeruleomycin E.¹⁸ These halogenations (Reissert–Henze reactions) require the use of N-oxides, easily made upon treatment of pyridines with an oxidant such as H₂O₂ or *m*-chloroperbenzoic acid (*m*CPBA).³⁷ Initially, using **1-O** as a model substrate, activation with Tf₂O and bromination with tetra-*n*-butylammonium bromide (TBAB) was attempted, in analogy



Scheme 5 Ligand-coupling reactions with phosphonium salt **2**. ¹H NMR yield determined using CH₂Br₂ as internal standard (in brackets, isolated yield).


with the *ortho*-bromination of quinolines reported by Baran and co-workers (Scheme 6).³⁸ However, no desired product was observed, but only a mixture of brominated pyridines, probably due to some Br₂ formed from the residual oxidant (*m*CPBA) still present in **1-O**. In contrast to pyridines, the successful *ortho*-bromination observed by Baran for quinolines reflects the lower loss in resonance stabilization typical of bicyclic aromatics (*i.e.* naphthalene *vs.* benzene). Chlorination of **1-O** with POCl₃ gave better results (**9** isolated in 55% yield), but the need for harsh conditions (neat POCl₃ refluxing at 106 °C) somehow defeated our original purpose for a short and mild synthesis, thus a completely different strategy was considered.

Minisci-type chemistry is an excellent method for *ortho*-functionalizations of pyridines, especially employing nucleophilic carbon-based radicals (ideal for the synthesis of caeruleomycin K).^{39,40} Moreover, starting with a 4-substituted pyridine would prevent regioselectivity issues (C2 *vs.* C6) typical of unsymmetrical starting materials. For the *ortho*-arylation of pyridines, Baran and others have shown how aryl boronic esters, in combination with AgNO₃, Na₂S₂O₈, and TFA, are excellent aryl radical precursors.^{41,42} To install a carbonyl group, Angeles, Yeung, and colleagues have used 1,3,5-trioxanes as an aldehyde equivalent in Minisci-type carbonylation of pyridines.⁴³ Based on this precedent, a Minisci arylation of 4-chloropyridine **10** was performed (Scheme 7). In this case, an excess of phenylboronic acid (1.5 equiv.) was needed to compensate for competitive protodeboronation, whereas a higher loading of AgNO₃ allowed the isolation of product **11** in 56% yield. A second Minisci reaction was then performed on this pyridine using 1,3,5-trioxane in the presence of (*n*Bu₄N)₂S₂O₈. A successful *ortho*-alkylation gave product **12** in 50% yield (a value in agreement with previous reports),^{44–47} whereas a subsequent nucleophilic aromatic substitution allowed to access trifunctionalized pyridine **13** almost quantitatively. It has to be noted that starting from 4-methoxypyridine **14** would shorten the synthesis, but the electron-donating effect of the methoxy group will negatively affect both steps since Minisci reactions are based on the addition of nucleophilic radicals. The final conversion of **13** into caeruleomycin K was achieved in a one-pot procedure by treatment with HCl (to reveal the aldehyde functionality), followed by condensation with NH₂OH. Importantly, **12** could be directly converted into caeruleomycin K



Scheme 6 *ortho*-Halogenations with **1-O**.





Scheme 7 *ortho*-Functionalization of 10 and total synthesis of caerulomycin K.

without the need for isolation of 13, further simplifying the synthesis (route in green). Therefore, this three-step total synthesis (overall yield of 10%) represents the first synthesis of caerulomycin K as well as a potential alternative to the synthesis of caerulomycins.

In conclusion, the first total synthesis of caerulomycin K has been reported. Starting from monofunctionalized pyridines, the first strategy looked at a double C–H activation by means of phosphonium chemistry. However, a poor conversion of the second C–P bond formation and a problematic *ortho*-methylation, including *via* halogenation, prompted the search for a better alternative. This was achieved by sequential Minisci *ortho*-arylation and *ortho*-alkylation, with the latter converted in one pot into the desired oxime. Compared to previously reported caerulomycin syntheses, this novel approach does not require highly pre-functionalized starting materials.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors acknowledge Miriana Guarnaccia and Marissa Panethiere for preliminary results as well as Dr Elisa Moretti and Dr Francesco Castagnetti for mass analyses. The authors acknowledge Chiesi Farmaceutici S.p.A for funding and the support of the APC central fund of the University of Milan.

Notes and references

- 1 S. De, A. Kumar, S. K. Shah, S. Kazi, N. Sarkar, S. Banerjee and S. Dey, *RSC Adv.*, 2022, **12**, 15385–15406.

- 2 Y. Ling, Z. Y. Hao, D. Liang, C. L. Zhang, Y. F. Liu and Y. Wang, *Drug Des., Dev. Ther.*, 2021, **15**, 4289–4338.
- 3 S. Maity, A. Bera, A. Bhattacharjya and P. Maity, *Org. Biomol. Chem.*, 2023, **21**, 5671–5690.
- 4 M. Failla, G. W. Lombardo, P. Orlando, D. Fiorito, E. Bombonato, P. Ronchi, D. Passarella and V. Fasano, *Eur. J. Org. Chem.*, 2023, e202300074.
- 5 C. M. Josephitis, H. M. H. Nguyen and A. McNally, *Chem. Rev.*, 2023, **123**, 7655–7691.
- 6 L. C. Vining, A. G. McInnes, A. W. McCulloch, D. G. Smith and J. A. Walter, *Can. J. Chem.*, 1988, **66**, 191–194.
- 7 A. G. McInnes, D. G. Smith, J. A. Walter, L. C. Vining and J. L. C. Wright, *Can. J. Chem.*, 1979, **57**, 3200–3204.
- 8 D. K. Chatterjee, W. Raether, N. Iyer and B. N. Ganguli, *Z. Parasitenkd.*, 1984, **70**, 569–573.
- 9 D. Chen, Q. Zhao and W. Liu, *J. Ind. Microbiol. Biotechnol.*, 2019, **46**, 459–468.
- 10 B. Pang, R. Liao, Z. Tang, S. Guo, Z. Wu and W. Liu, *Nat. Commun.*, 2021, **12**, 3124.
- 11 P. Fu, S. Wang, K. Hong, X. Li, P. Liu, Y. Wang and W. Zhu, *J. Nat. Prod.*, 2011, **74**, 1751–1756.
- 12 B. D. Alreja, S. L. Kattige, B. La and N. J. de Souza, *Heterocycles*, 1986, **24**, 1637–1640.
- 13 J. Dash and H. U. Reissig, *Chem.–Eur. J.*, 2009, **15**, 6811–6814.
- 14 D. N. Bobrov and V. Tyvorskii, *Tetrahedron*, 2010, **66**, 5432–5434.
- 15 F. Zhang and X.-F. Duan, *Org. Lett.*, 2011, **13**, 6102–6105.
- 16 A. M. Horan, V. K. Duong and E. M. McGarrigle, *Org. Lett.*, 2021, **23**, 9089–9093.
- 17 F. Mongin, F. Trécourt, B. Gervais, O. Mongin and G. Quéguiner, *J. Org. Chem.*, 2002, **67**, 3272–3276.
- 18 F. Trécourt, B. Gervais, O. Mongin, C. Le Gal, F. Mongin and G. Quéguiner, *J. Org. Chem.*, 1998, **63**, 2892–2897.
- 19 X. F. Duan, Z. Q. Ma, F. Zhang and Z. B. Zhang, *J. Org. Chem.*, 2009, **74**, 939–942.
- 20 T. Sammakia, E. L. Stangeland and M. C. Whitcomb, *Org. Lett.*, 2002, **4**, 2385–2388.
- 21 F. Mongin, F. Trécourt, B. Gervais, O. Mongin and G. Quéguiner, *J. Org. Chem.*, 2002, **67**, 3272–3276.
- 22 R. G. Anderson, B. M. Jett and A. McNally, *Angew. Chem., Int. Ed.*, 2018, **57**, 12514–12518.
- 23 R. D. Dolewski, M. C. Hilton and A. McNally, *Synlett*, 2018, **29**, 8–14.
- 24 J.-P. Finet, *Tetrahedron Org. Chem. Ser.*, 1998, **18**, 9–46.
- 25 E. Anders and F. Markus, *Chem. Ber.*, 1989, **122**, 119–122.
- 26 E. Anders and F. Markus, *Chem. Ber.*, 1989, **122**, 113–118.
- 27 M. C. Hilton, R. D. Dolewski and A. McNally, *J. Am. Chem. Soc.*, 2016, **138**, 13806–13809.
- 28 P. Du, Y. Yin, D. Shi, K. Mao, Q. Yu and J. Zhao, *Molecules*, 2022, **27**, 5694.
- 29 J. L. Koniarczyk, D. Hesk, A. Overgard, I. W. Davies and A. McNally, *J. Am. Chem. Soc.*, 2018, **140**, 1990–1993.
- 30 R. G. Anderson, B. M. Jett and A. McNally, *Tetrahedron*, 2018, **74**, 3129–3136.
- 31 C. Patel, M. Mohnike, M. C. Hilton and A. McNally, *Org. Lett.*, 2018, **20**, 2607–2610.



- 32 X. Zhang and A. McNally, *Angew. Chem., Int. Ed.*, 2017, **56**, 9833–9836.
- 33 M. C. Hilton, X. Zhang, B. T. Boyle, J. V. Alegre-Requena, R. S. Paton and A. McNally, *Science*, 2018, **362**, 799–804.
- 34 X. Zhang and A. McNally, *ACS Catal.*, 2019, **9**, 4862–4866.
- 35 S. Oae and Y. Uchida, *Acc. Chem. Res.*, 1991, **24**, 202–208.
- 36 R. Hoffmann, J. M. Howell and E. L. Muetterties, *J. Am. Chem. Soc.*, 1972, **94**, 3047–3058.
- 37 D. Wang, L. Désaubry, G. Li, M. Huang and S. Zheng, *Adv. Synth. Catal.*, 2021, **363**, 2–39.
- 38 S. E. Wengryniuk, A. Weickgenannt, C. Reiher, N. A. Strotman, K. Chen, M. D. Eastgate and P. S. Baran, *Org. Lett.*, 2013, **15**, 792–795.
- 39 R. S. J. Proctor and R. J. Phipps, *Angew. Chem., Int. Ed.*, 2019, **58**, 13666–13699.
- 40 M. A. J. Duncton, *MedChemComm*, 2011, **2**, 1135–1161.
- 41 I. B. Seiple, S. Su, R. A. Rodriguez, R. Gianatassio, Y. Fujiwara, A. L. Sobel and P. S. Baran, *J. Am. Chem. Soc.*, 2010, **132**, 13194–13196.
- 42 P. P. Singh, S. K. Aithagani, M. Yadav, V. P. Singh and R. A. Vishwakarma, *J. Org. Chem.*, 2013, **78**, 2639–2648.
- 43 J. M. Ganley, M. Christensen, Y. H. Lam, Z. Peng, A. R. Angeles and C. S. Yeung, *Org. Lett.*, 2018, **20**, 5752–5756.
- 44 H. Tian, H. Yang, C. Tian, G. An and G. Li, *Org. Lett.*, 2020, **22**, 7709–7715.
- 45 H. Zhao and J. Jin, *Org. Lett.*, 2019, **21**, 6179–6184.
- 46 H. Zhao, Z. Li and J. Jin, *New J. Chem.*, 2019, **43**, 12533–12537.
- 47 G. P. Gardini, *Tetrahedron Lett.*, 1972, **13**, 4113–4116.

