RSC Advances



PAPER

View Article Online
View Journal | View Issue



Cite this: RSC Adv., 2024, 14, 13640

Lewis acid-mediated transformations of 5-acyl-N-fluoroalkyl-1,2,3-triazoles to cyclopentenones, indenones, or oxazoles†

Lukáš Janecký o ab and Petr Beier **

Received 5th March 2024 Accepted 19th April 2024

DOI: 10.1039/d4ra01707b

rsc.li/rsc-advances

We present a transition metal-free approach to 2-N-substituted indenones, cyclopentenones, and 4-carbonyl oxazoles, based on the reaction of 5-acylated N-fluoroalkyl substituted 1,2,3-triazoles (prepared by a three-component click reaction of copper acetylides, fluoroalkyl azides, and acyl chlorides) with Lewis acids aluminium trichloride or boron trifluoride etherate, proceeding via the generation and cyclization of vinyl cations.

cyclopentenones.

(Scheme 1).29

Introduction

Multi-substituted cyclopentenones, indenones and 4carbonyl oxazoles constitute important classes of biologically active compounds known as anti-inflammatory/ anticancer agents or enzyme inhibitors (Fig. 1).1-6 Synthetic strategies for obtaining their N-alkenyl derivatives (2-Nsubstituted cyclopentenones and indenones) or fully substituted oxazoles are limited because of the low availability of the starting materials and the necessity to use transition metal complexes or harsh reaction conditions. There is no general synthetic approach leading to these structures and each type of product requires a specific methodology. Despite the availability of numerous synthetic methods for the preparation of mono- or di-substituted oxazoles,7 access to tri-substituted 4-carbonyl oxazoles is not well explored and relies mainly on the intramolecular Cu-catalyzed cyclization of (thio)enamides8 or bromo(thio) enones,9 or on a protocol starting from 2-azido enones10 or alkynyl ketones.11

2-N-substituted indenones can be accessed from ynamides, ¹² 2-alkynylbenzoyl cyanides, ¹³ or 2-hydroxy-substituted internal alkynes. ¹⁴ 2-Amino indenones were also prepared *via* Aucatalyzed intermolecular oxidation of 2-carbonyl-1-ethynyl benzenes ¹⁵ or by co-catalyzed annulation of thioamides with ynamides. ¹⁶ Approaches leading to 2-amino-substituted cyclopentenones are limited to methods starting from previously modified cyclopentenone or cyclopentane rings. ¹⁷⁻¹⁹

Other procedures, starting from ynamides, 20 α-aminoenals21 or

vinyl ketenes,²² are highly substrate-specific and do not allow for the further modification of the amino position in

We recently reported Brønsted and Lewis acid-mediated

reactions of N-fluoroalkylated 1,2,3-triazoles,23-25 as an effi-

cient approach to the generation of vinyl cation intermediates which then reacted with various nucleophiles providing *N*-

alkenyl compounds such as β-enamido triflates,²⁶ β-fluoro enamides²⁷ and β-halo alkenyl imidoyl halides,²⁸ or undergo

cyclization reactions to form multisubstituted cyclopentenes

Herein, we propose a new synthetic methodology to 2-N-

Fig. 1 Selected examples of bioactive 4-carbonyl oxazoles, cyclopentenones and indenones.

substituted cyclopentenones, indenones and 4-carbonyl oxazoles from 5-acyl-*N*-fluoroalkyl-1,2,3-triazoles by a treatment with a Lewis acid (AlCl₃ or BF₃·OEt₂), proceeding νia vinyl cations and their cyclization onto alkenes or arenes (Scheme 2). This type of cyclization is very rare.³⁰

[&]quot;The Institute of Organic Chemistry and Biochemistry of the Czech Academy of Sciences, Flemingovo nam. 2, 16610 Prague 6, Czech Republic. E-mail: beier@uochb.cas.cz

^bDepartment of Organic Chemistry, Faculty of Science, Charles University, Hlavova 2030/8, 128 43 Prague 2, Czech Republic

 $[\]dagger$ Electronic supplementary information (ESI) available. See DOI: $\label{eq:boltz} https://doi.org/10.1039/d4ra01707b$

Scheme 1 Brønsted and Lewis acid-mediated denitrogenation of N-fluoroalkyl-1,2,3-triazoles (HA = TfOH or FSO₃H, LA = BF₃·OEt₂ or AlX₃).

$$\begin{array}{c} N=N\\ R^1\\ \hline\\ N-CF_2R^3\\ \hline\\ LA (AlCl_3\\ or BF_3OEt_2) - N_2\\ \hline\\ LA \\ N-CF_2R^3\\ \hline\\ R^1\\ \hline\\ N-CF_2R^3\\ \hline\\ R^2=Alkenyl\\ \hline\\ R^3\\ \hline\\ N-CI\\ R^3\\ \hline\\ R^3\\ \hline\\ N-CI\\ N-CI\\ R^3\\ \hline\\ N-CI\\ R^3\\ \hline\\ N-CI\\ R^3\\ \hline\\ N-CI\\ R^3\\ \hline\\ N-CI\\ R$$

Scheme 2 Proposed Lewis acid-mediated transformation of 5-acyl-1,2,3-triazoles into new cyclic products.

Results and discussion

Synthesis of 5-acetyl- and 5-benzoyl-substituted 1,2,3-triazoles by an intercepted click reaction was briefly described in the literature. 31 For our study it was necessary to prepare a library of 1-(per)fluoroalkyl-4-substituted-5-acyl-1,2,3-triazoles. However, application of Wu's conditions³¹ (phenylacetylene, CuI, Et₃N and benzoyl chloride or acetyl chloride) with our fluorinated azides 23,32,33 (CF $_3N_3$ or C $_2F_5N_3$) did not afford the desired 5acylated triazoles. Therefore, we turned to the use of (phenylethynyl)copper (1a), which was shown to be reactive with azido(per)fluoroalkanes in intercepted click iodination or allylation.28 A three-component reaction of 1a, azidopentafluoroethane (2a) and methacryloyl chloride was used for the optimization of the synthesis of triazole 3a (Table 1). Initially, a mixture of 3a and triazole side-product 3a-H formed (entry 1). Increasing the amount of acyl chloride and using anhydrous conditions improved the yield of 3a and suppressed the formation of 3a-H (entries 2-4). To further increase the yield of 3a, screening for an additional base was conducted (entries 5-11), identifying DIPEA as the most efficient one (entry 12). Finally, the use of 2 equiv. of acyl chloride and 3 equiv. of DIPEA provided 3a in optimized yield (entry 14).

Having established the optimal reaction conditions for the intercepted click reaction and acylation sequence, the scope of the protocol was investigated on diverse acetylenic substrates,

Table 1 Optimization of reaction conditions leading to triazole 3a^a

$$\begin{array}{c|c} Cu & C_2F_5N_3 \\ \hline & (2a) \\ Ph & Cu \\ \end{array}$$

$$\begin{array}{c|c} N=N \\ N-C_2F_5 \\ \hline & (Base) \\ \end{array}$$

$$\begin{array}{c|c} N=N \\ N-C_2F_5 \\ \hline & H \\ \end{array}$$

Entry	RCOCl (equiv.)	Base (equiv.)	Ratio $3a/3a-H^b$	Yield 3a (%)
1^d	1	_	75:25	n.d.
2	1.2	_	99:1	51
3	2	_	100:0	53
4	5	_	100:0	33
5	1.2	Pyridine (1)	61:39	n.d.
6	1.2	NaOMe (2)	22:78	n.d.
7	1.2	Et ₃ N (2)	80:20	n.d.
8	1.2	$NaNH_2$ (3)	38:62	n.d.
9	1.2	KF (1)	100:0	45
10	1.2	DBU (2)	0:100	0
11	1.2	<i>i</i> -Pr ₂ NH (2)	22:78	n.d.
12	1.2	DIPEA (2)	100:0	63
13	2	DIPEA (2)	100:0	75
14	2	DIPEA (3)	100:0	80
15	2	DIPEA (5)	100:0	46

 a Reaction conditions: 1a (1 mmol), 2a (1.5 equiv.), THF (4 mL), 0–25 °C, then RCOCl, base, 3 Å MS (240 mg), rt, 18 h. b 19 F NMR ratio. c Isolated yield. d Without 3 Å molecular sieves. n.d. not determined.


fluorinated azides, and acid chlorides (Scheme 3). 5-Methacryloyl triazoles (3b-e) with strongly electron-poor or electronrich aryl rings in position 4 were prepared in moderate yields. To demonstrate the scalability of reaction, triazole 3a was prepared on 1.76 g (10.7 mmol) scale in high yield. 5-Benzoyl-substituted triazoles (3f-i) were also prepared in good yields. Varying the azide reagent revealed that highly fluorinated azidoalkanes afforded better product yields than tosyl azide or ethyl azidodifluoroacetate. Modification of triazoles in position 4 with aryl, alkyl and alkenyl groups and modification in position 5 with aryl, heteroaryl, alkyl, cycloalkyl and alkenyl groups afforded products mostly in satisfactory yields. Scale-up of 3f and 3y to 4–5 mmol was also successful.

Next, AlCl3 was chosen as a suitably strong and easily available Lewis acid to investigate the denitrogenative transformation of 5-acryloyl substituted triazoles. Cyclization of the formed vinyl cation intermediate onto the alkene moiety and chloride capture of the resulting carbocation led to the formation of cyclopentenone imidoyl chlorides (4a-d) in moderate yields (Scheme 4). The cation-stabilizing p-methoxyphenyl group in position 4 of triazole 3 improved the product yields. This observation, together with the necessity of triazole denitrogenation in the initial step of the reaction, speaks against an alternative reaction mechanism involving Nazarov cyclization, typically starting from a divinyl ketone. BF3 · OEt2 was used for the preparation of fluorinated cyclopentenone amides 4e-g from triazoles 3 (Scheme 4). In the case of N-CF₃ triazole 3j, the final product is not a cyclopentenone amide but cyclopentenone isocyanate 4h due to the facile HF elimination of the NHC(O)F intermediate.

$$R^{1} = -Cu \xrightarrow{R^{3}CF_{2}N_{3} (\sim 1.5 \text{ equiv.})} -Cu \xrightarrow{R^{2}COCI (2 \text{ equiv.})} -R^{1} \xrightarrow{N-CF_{2}R^{3}}$$

$$1 \qquad THF, \text{ mol. sieves} \qquad R^{2} \xrightarrow{N^{2}}$$

Scheme 3 Substrate scope of the intercepted click reaction leading to 5-acyl triazoles 3. Reaction conditions: 1 (1.0 mmol), 2 (1.5 mmol), DIPEA (3.0 mmol), acyl halide (2.0 mmol), THF (4 mL), 3 Å MS (240 mg), rt, 18 h. a 10.7 mmol scale. b 4.16 mmol scale. c 5.0 mmol scale.



Scheme 4 AlCl₃-mediated denitrogenation/cyclization of triazoles 3 leading to cyclopentenone imidoyl chlorides 4a-d and BF₃·OEt₂mediated denitrogenation/cyclization of triazoles 3 leading to cyclopentenone amides 4e-g and cyclopentenone isocyanate 4h. Reaction conditions: 3 (1.0 mmol), AlCl₃ or BF₃ OEt₂ (1.1 mmol), DCE (0.1M), 60 ° C, 1-2 h. ^a2.2 mmol scale.

Triazole 3r was a special case, as the vinyl cation intermediate induced a 1,5-hydride shift to form a tertiary carbocation and an α,β -unsaturated ketone. Cyclization and proton elimination afforded cyclopentenone 4i (Scheme S1 in the ESI†).

Subjecting 5-benzoyl triazoles 3 to a reaction with AlCl₃ led to the cyclization on the aryl ring of the benzoyl moiety, forming indenone imidoyl chlorides 5 (Scheme 5). Again, the presence of vinyl cation-stabilizing groups in position 4 of the starting triazole and electron-rich groups on the aryl ring of the substituted benzoyl moiety in position 5 of the triazole both increased product yields. Using 5-benzoyl triazoles instead of dimethoxybenzoyl triazoles afforded only low to moderate indenone yields.

The formation of 4-acyl oxazoles 6a-g was observed upon boron trifluoride-mediated transformation of 5-benzoyl triazoles (Scheme 6). Two isomers of products 6 can be formed: electron-rich or -neutral aryl groups in position 5 of the starting triazole cyclized selectively to form isomer A and triazoles with deactivated aryl groups in position 5 gave a mixture of isomers A and B with good to high selectivity for isomer A. Heating the mixture of isomers in a microwave did not lead to their interconversion, ruling out the possibility of Cornforth rearrangement in 2-trifluoromethyloxazoles. Oxazole 6f was accompanied by indenone 5f side-product. Oxazolone 6g was prepared selectively from 3m, as 2-fluorooxazoles are hydrolytically unstable.

To rationalize the formation of products 6A and 6B, we considered the following reaction mechanism (Scheme 7). As demonstrated in our earlier report,27 the coordination of BF3 to nitrogen atoms of the triazole ring (particularly the coordination to N1) led to opening of the triazole ring and the

Scheme 5 AlCl $_3$ -mediated denitrogenation/cyclization of triazoles 3 leading to indenone imidoyl chlorides 5. Reaction conditions: 3 (1.0 mmol), AlCl $_3$ (1.1 mmol), DCE (0.1M), 60 °C, 1–2 h.

5e. 56%

5d. 48%

Scheme 6 $BF_3\cdot OEt_2$ -mediated denitrogenation/cyclization of triazoles 3 leading to 4-acyl oxazoles 6. Reaction conditions: 3 (1.0 mmol), $BF_3\cdot OEt_2$ (1.1 mmol), DCE (0.1M), 60 °C, 1–2 h.

elimination of the nitrogen molecule from the diazonium intermediate C. The resulting vinyl cation D underwent elimination of tetrafluoroborate and fluoride transfer to form imidoyl fluoride E. Cyclization of the acyl oxygen led to 6A. Alternatively, hydrolysis of the vinyl fluoride moiety of E created a competitive acyl moiety (oxygen nucleophile) for cyclization to 6B. The required one molecule of water probably comes from the moisture in the solvent. Deliberate addition of a small amount of water reduced product yields,

$$\begin{array}{c} \begin{array}{c} \begin{array}{c} N=N\\ Ar^1 \end{array} \end{array} \begin{array}{c} N=N\\ Ar^2 \end{array} \begin{array}{c} N=N\\$$

Scheme 7 Mechanism of BF_3 -mediated transformation of triazoles 3 to oxazoles 6.

Scheme 8 Synthetic utilization of cyclopentenone 4a.

presumably because of competitive hydrolysis of the used Lewis acid.

The synthetic utility of cyclopentenone imidoyl chlorides was demonstrated on examples of post-functionalization of **4a** (Scheme 8). Trifluoromethylated tetrazole **7a** was easily prepared from **4a** using sodium azide. The addition of an aqueous ammonia solution afforded amidine **7b**, and the addition of aqueous hydrazine led to the cyclization of the carbonyl and imidoyl chloride functional groups to give trifluoromethylated triazine **7c**. We proposed that **7c** is formed by cyclization of hydrazine nitrogen to the six-membered ring,

Scheme 9 Proposed mechanism of the formation of 7c.

followed by the hydroxyl shift and substitution of the chlorine atom for oxygen on the cyclopentene moiety (Scheme 9).

Conclusions

In conclusion, we present the synthesis of *N*-electron-acceptor group-substituted 5-acyl-1,2,3-triazoles by an intercepted click reaction, namely a three-component cyclization of azide, copper acetylide and acyl chloride. Lewis acid-mediated triazole ring opening and nitrogen molecule elimination provided key reactive intermediates - vinyl cations, which cyclized selectively to form either cyclopentenone imidoyl chlorides, indenone imidoyl chlorides, cyclopentenone amides, or 2-trifluoromethyl oxazoles, depending on the combination of the Lewis acid used and the substitution in position 5 of the triazole ring. Postfunctionalization of cyclopentenone imidoyl chloride gave access to selectively functionalized N-alkenyl compounds (amidines) or new nitrogen heterocycles (triazine or tetrazole). The presented methodology demonstrates a Lewis acidmediated generation of vinyl cations from triazoles and their synthetic utilization in the formation of new C-C bonds.

Author contributions

PB supervised the project. LJ contributed to experiments and product characterization. LJ and PB jointly conceived the project, prepared the manuscript, and contributed to discussions.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was financially supported by the Czech Academy of Sciences (Research Plan RVO: 61388963) and by the Czech Science Foundation (Project 23-04659S).

References

- 1 A. Y. Shaw, M. C. Henderson, G. Flynn, B. Samulitis, H. Han, S. P. Stratton, H. H. S. Chow, L. H. Hurley and R. T. Dorr, Characterization of Novel Diaryl Oxazole-Based Compounds as Potential Agents to Treat Pancreatic Cancer, J. Pharmacol. Exp. Ther., 2009, 331, 636–647.
- 2 J. H. Ahn, M. S. Shin, S. H. Jung, S. K. Kang, K. R. Kim, S. D. Rhee, W. H. Jung, S. D. Yang, S. J. Kim, J. R. Woo, et al., Indenone Derivatives: A Novel Template for Peroxisome Proliferator- Activated Receptor γ (PPARγ) Agonists, J. Med. Chem., 2006, 49, 4781–4784.
- 3 Y. Hayashi, K. Ogawa, F. Inagaki and C. Mukai, First Total Synthesis of (+)-Indicanone, *Org. Biomol. Chem.*, 2012, **10**, 4747–4751.
- 4 G. Mehta and H. M. Shinde, An Approach to Seco-Prezizaane Sesquiterpenoids: Enantioselective Total Synthesis of (+)-1S-Minwanenone, *J. Org. Chem.*, 2012, 77, 8056–8070.

- 5 D. C. Harrowven, N. A. Newman and C. A. Knight, On the Identity of a Neo-Lignan from the Fruits of Virola Sebifera, *Tetrahedron Lett.*, 1998, **39**, 6757–6760.
- 6 N. Desroy, A. Denis, C. Oliveira, D. Atamanyuk, S. Briet, F. Faivre, G. Lefralliec, Y. Bonvin, M. Oxoby, S. Escaich, et al., Novel HldE-K Inhibitors Leading to Attenuated Gram Negative Bacterial Virulence, J. Med. Chem., 2013, 56, 1418–1430.
- 7 S. Joshi, M. Mehra, R. Singh and S. Kakar, Review on Chemistry of Oxazole Derivatives: Current to Future Therapeutic Prospective, *Egypt. J. Basic Appl. Sci.*, 2023, **10**, 218–239.
- 8 S. Vijay Kumar, B. Saraiah, N. C. Misra and H. Ila, Synthesis of 2-Phenyl-4,5-Substituted Oxazoles by Copper-Catalyzed Intramolecular Cyclization of Functionalized Enamides, *J. Org. Chem.*, 2012, 77, 10752–10763.
- 9 S. V. Kumar, A. Acharya and H. Ila, Synthesis of 2,4,5-Trisubstituted Oxazoles with Complementary Regioselectivity from α-Oxoketene Dithioacetals and β-(Methylthio)-β-(Het)Aryl-2-Propenones, *J. Org. Chem.*, 2018, 83, 6607–6622.
- 10 K. Rajaguru, A. Mariappan, R. Suresh, P. Manivannan and S. Muthusubramanian, Investigation on the Reactivity of α-Azidochalcones with Carboxylic Acids: Formation of α-Amido-1,3-Diketones and Highly Substituted 2-(Trifluoromethyl)Oxazoles, *Beilstein J. Org. Chem.*, 2015, 11, 2021–2027.
- 11 A. Y. Dubovtsev, D. V. Dar'in and V. Y. Kukushkin, Three-Component [2+2+1] Gold(I)-Catalyzed Oxidative Generation of Fully Substituted 1,3-Oxazoles Involving Internal Alkynes, *Adv. Synth. Catal.*, 2019, **361**, 2926–2935.
- 12 S. Golling, P. Hansjacob, N. Bami, F. R. Leroux and M. Donnard, Direct Access to 2,3-Disubstituted Amido-Indenones through Annulation of 2-Iodobenzaldehydes with Ynamides, J. Org. Chem., 2022, 87, 16860–16866.
- 13 H. Shimizu and M. Murakami, Reaction of 2-Alkynylbenzoyl Cyanides with Carboxylic Acids Producing Functionalized Indenones, *Synlett*, 2008, **12**, 1817–1820.
- 14 J. Sun, G. Zheng, T. Xiong, Q. Zhang, J. Zhao, Y. Li and Q. Zhang, Copper-Catalyzed Hydroxyl-Directed Aminoarylation of Alkynes, *ACS Catal.*, 2016, **6**, 3674–3678.
- 15 B. D. Mokar, D. B. Huple and R. S. Liu, Gold-Catalyzed Intermolecular Oxidations of 2-Ketonyl-1-Ethynyl Benzenes with N-Hydoxyanilines to Yield 2-Aminoindenones via Gold Carbene Intermediates, *Angew. Chem., Int. Ed.*, 2016, 55, 11892–11896.
- 16 S. Sau, A. Ghosh, M. Shankar, M. P. Gogoi and A. K. Sahoo, Cobalt-Catalyzed Thioamide Directed C(Arene)-H Annulation with Ynamide: Regioselective Access to 2-Amidoindenones, Org. Lett., 2022, 24, 9508–9513.
- 17 Z. Wang, B. J. Reinus and G. Dong, Catalytic Intermolecular β -C–H Alkenylation of α -Enamino-Ketones with Simple Alkynes, *Chem. Commun.*, 2014, **50**, 5230–5232.
- 18 H. Ottinger, T. Soldo and T. Hofmann, Systematic Studies on Structure and Physiological Activity of Cyclic α-Keto Enamines, a Novel Class of "Cooling" Compounds, *J. Agric. Food Chem.*, 2001, **49**, 5383–5390.

- 19 Y. Li, R. Zhang, X. Bi and J. Fu, Multifunctionalization of Unactivated Cyclic Ketones via Synergistic Catalysis of Copper and Diarylamine: Access to Cyclic α-Enaminone, *Org. Lett.*, 2018, **20**, 1207–1211.
- 20 J. Balsells, J. Vazquez, A. Moyano, M. A. Pericas and A. Riera, Low-Energy Pathway for Pauson-Khand Reactions: Synthesis and Reactivity of Dicobalt Hexacarbonyl Complexes of Chiral Ynamines, J. Org. Chem., 2000, 65, 7291–7302.
- 21 R. B. Shnigirev, I. A. Ushakov, V. A. Semenov and A. Y. Rulev, Reactions of α-Functionally Substituted Enals with Terminal Alkynes: Unexpected Assembly of 2-Amino-2-Cyclopentenones, *J. Org. Chem.*, 2023, **88**, 4886–4890.
- 22 J. H. Rigby and Z. Wang, Synthesis of Highly Substituted Cyclopentenones via the [4 + 1] Cycloaddition of Nucleophilic Carbenes and Vinyl Ketenes, *Org. Lett.*, 2003, 5, 263–264.
- 23 Z. E. Blastik, S. Voltrová, V. Matoušek, B. Jurásek, D. W. Manley, B. Klepetářová and P. Beier, Azidoperfluoroalkanes: Synthesis and Application in Copper(I)-Catalyzed Azide–Alkyne Cycloaddition, *Angew. Chem., Int. Ed.*, 2017, 56, 346–349.
- 24 O. Bakhanovich and P. Beier, Synthesis, Stability and Reactivity of α-Fluorinated Azidoalkanes, *Chem.–Eur. J.*, 2020, **26**, 773–782.
- 25 A. Markos, V. Matoušek and P. Beier, Fluoroalkyl Azides and Triazoles: Unlocking a Novel Chemical Space, *Aldrichimica Acta*, 2022, **55**, 37–44.
- 26 A. Markos, S. Voltrová, V. Motornov, D. Tichý, B. Klepetářová and P. Beier, Stereoselective Synthesis of (Z)-β-Enamido Triflates and Fluorosulfonates from N-Fluoroalkylated Triazoles, *Chem.-Eur. J.*, 2019, 25, 7640–7644.

- 27 A. Markos, L. Janecký, B. Klepetářová, R. Pohl and P. Beier, Stereoselective Synthesis of (Z)-β-Enamido Fluorides from N-Fluoroalkyl- And N-Sulfonyl-1,2,3-Triazoles, *Org. Lett.*, 2021, 23, 4224–4227.
- 28 A. Markos, L. Janecký, T. Chvojka, T. Martinek, H. Martinez-Seara, B. Klepetářová and P. Beier, Haloalkenyl Imidoyl Halides as Multifacial Substrates in the Stereoselective Synthesis of N-Alkenyl Compounds, *Adv. Synth. Catal.*, 2021, 363, 3258–3266.
- 29 L. Janecký, A. Markos, B. Klepetářová and P. Beier, Lewis-Acid-Mediated Intramolecular Cyclization of 4-Aryl-5-Allyl-1,2,3-Triazoles to Substituted Cyclopentene Derivatives, *J. Org. Chem.*, 2023, 88, 1155–1167.
- 30 J. C. Corcoran, R. Guo, Y. Xia and Y. M. Wang, Vinyl Cation-Mediated Intramolecular Hydroarylation of Alkynes Using Pyridinium Reagents, *Chem. Commun.*, 2022, **82**, 11523–11526.
- 31 Y. M. Wu, J. Deng, Y. Li and Q. Y. Chen, Regiospecific Synthesis of 1,4,5-Trisubstituted-1,2,3-Triazole via One-Pot Reaction Promoted by Copper(I) Salt, *Synthesis*, 2005, 8, 1314–1318.
- 32 S. Voltrová, M. Muselli, J. Filgas, V. Matoušek, B. Klepetářová and P. Beier, Synthesis of Tetrafluoroethylene-and Tetrafluoroethyl-Containing Azides and Their 1,3-Dipolar Cycloaddition as Synthetic Application, *Org. Biomol. Chem.*, 2017, 15, 4962–4965.
- 33 E. Shaitanova, V. Matoušek, T. Herentin, M. Adamec, R. Matyáš, B. Klepetářová and P. Beier, Synthesis and Cycloaddition Reactions of 1-Azido-1,1,2,2-Tetrafluoroethane, *J. Org. Chem.*, 2023, **88**, 14969–14977.