


 Cite this: *RSC Adv.*, 2024, 14, 34843

 Received 4th August 2024
 Accepted 22nd October 2024

DOI: 10.1039/d4ra05652c

rsc.li/rsc-advances

Clues from an ionic cocrystal structure: from catalysis to mechanochemistry†

 Bandana Sar, Mollah Rohan Ahsan  and Arijit Mukherjee *

A crystal structure is no longer conceived as a static entity; rather, it often mirrors crystallization pathways, linking crystal structures with a solution scenario. In this study, taking a clue from a previously reported ionic cocrystal structure, an *in situ* acetic acid-based catalytic protocol is developed for the *N*-acylation of amines using ester sources. In addition to better catalytic efficiency, this *in situ* approach was extended further to a mechanochemistry protocol in the context of a multicomponent reaction. This points towards a broader applicability of crystal structures that goes beyond the domain of structural chemistry and delves into catalysis and mechanosynthesis.

Crystal structures are historically viewed as static entities, mainly used for determining atomic coordinates and identifying intermolecular interactions.¹ The modern notion of crystals evolved with the advancement of supramolecular chemistry² and the description of crystals as “supermolecule par excellence”,^{3,4} has contributed immensely to the purpose.^{5–10} Despite some significant early milestones correlating crystal structures with chemical reactivity,^{3,11} the perception of some crystal structures being “crystallization snapshots” took time to emerge.^{12,13} It is only recently, with the concept of crystal structure landscape,^{5,7,12} that crystal structures are conceived as data points in a given structural landscape and can, therefore, be linked with crystallization pathways.¹³

In 2006, Davey and co-workers showed, with the example of tetrolic acid (TA), that polymorphism in a given system can be linked with solution structures.^{14–16} Later, while investigating TA crystallization from ionic liquids, Myerson and co-workers identified that the metastable (α) form could be obtained while crystallized from hydrophobic ionic liquids.¹⁷ Using a polar ionic liquid 1-ethyl-3-methylimidazolium acetate ([Emim]⁺[OAc][−]) as a crystallization medium, however, led to an ionic cocrystal which was crystallized in *P*1, with cell parameters of $a = 6.746(3) \text{ \AA}$, $b = 7.246(3) \text{ \AA}$, $c = 7.434(3) \text{ \AA}$; $\alpha = 77.801(7)^\circ$, $\beta = 83.650(8)^\circ$, $\gamma = 88.409(7)^\circ$.¹⁷ The TA molecules form a catemeric chain, and the respective C–O bond distances were found to be 1.279(2) \AA and 1.221(2) \AA with $\Delta\text{C–O} \sim 0.058 \text{ \AA}$. This lowering of $\Delta\text{C–O}$ could be attributed to the partial deprotonation in the ionic cocrystal.¹⁸ Henceforth, this ionic cocrystal is denoted as [TA]·[Emim]⁺ [TA][−] with a neutral TA,

a deprotonated TA, and a disordered [Emim]⁺ cation in the asymmetric unit (Fig. 1).¹⁷

While the disorder [Emim]⁺ may be understood from its mobile nature, the partially deprotonated TA, however, implies that the proton could be abstracted by the [OAc][−] anion of the




Fig. 1 (a) Chemical diagrams of tetrolic acid and 1-ethyl-3-methylimidazolium acetate, (b) the crystal structure of TA×[Emim]⁺ [TA][−] (protonated and deprotonated TA are marked with blue and red patches). Blue: nitrogen, red: oxygen, gray: carbon, white: hydrogen.

Department of Chemistry, Birla Institute of Science and Technology (BITS) Pilani, Hyderabad campus, Hyderabad 500078, Telangana, India. E-mail: arijit.mukherjee@hyderabad.bits-pilani.ac.in

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4ra05652c>



acid chlorides and acid anhydrides and generating acetic acid only through a dynamic equilibrium.

To investigate if the reaction was catalyzed by the *in situ* generated acetic acid, a blank reaction was first performed with 1 mmol benzylamine and 1 ml of EtOAc at 80 °C for 20 hours, only a $3.8 \pm 0.7\%$ of **1a** was produced (ESI-S3(a)†). While the reaction was carried out for 20 hours with [Emim]⁺[OAc]⁻ (10 mol% with respect to benzylamine in the absence of TA, a $17.6 \pm 0.3\%$ yield was observed (ESI-S3(b)†). On the other hand, $25.6 \pm 1.1\%$ of **1a** was obtained when TA (10 mol% with respect to benzylamine) was added to the reaction medium in the absence of [Emim]⁺[OAc]⁻ (ESI-S3(c)†). This may be attributed to the competition between formations of salts *vs.* amides, as often observed with organic acids.²⁶ These results indicate that a facile *N*-acetylation could not be carried out in the presence of either TA or [Emim]⁺[OAc]⁻, with sufficient yield. The reaction yield, however, was strikingly increased to 100% when [Emim]⁺[OAc]⁻ (10 mol%) and TA (10 mol%) were used together in the reaction mixture. When benzylamine was refluxed with EtOAc at 80 °C for 20 hours in the presence of TA and [Emim]⁺[OAc]⁻, **1a** was produced with 100% yield (ESI-S4(a)†). To investigate the role of reaction time, the reaction was carried out for different intervals ranging between 4 and 16 hours. The yield of **1a** varied between 18% and 86% during those intervals (Table 1) (ESI-S4(a)†). More significantly, **1a** was obtained in $97.5 \pm 1.8\%$ yield even when the catalytic load was reduced from 10% to 1% (ESI-S4(b)†). (Table 1). A comparison between this method and the previous method (ESI-S11†) shows a greater efficiency of the *in situ* technique over the previous methods.

Subsequently, the scope of the above approach was investigated toward variegated amines to check the generality as well as variation of yield% with electronic factors. Some of the systems chosen here are similar to the previous studies²⁴ to strike a direct comparison between the two approaches. In the presence of an electron-withdrawing group (such as -Br in **1b** and -CF₃ in **1c**), the yield of **1b** and **1c** was found to be $\sim 91 \pm 0.8\%$ and $\sim 43.7 \pm 1.8\%$ (ESI-S5(a and b)†), respectively, when the reaction was performed with 1 mol% catalytic load for 20 hours at 80 °C, indicating the effectiveness of the approach even in slightly deactivated systems.

The chemoselectivity of the reaction was also investigated, and the desired product **1d** was obtained in quantitative yield. The peak at 1633 cm^{-1} in FTIR-ATR confirmed the formation of an amide bond and proved the chemoselectivity of the reaction (ESI-S5(c)†). Since more vigorous conditions may be required for secondary amines, butyl acetate (b.p. = 126 °C) was used during the synthesis of **1e** (with 1 mol% catalyst), and the reaction was performed at 115 °C. Under these conditions, **1e** was formed with a $99.3 \pm 0.5\%$ yield after 24 hours (ESI-S5(d)†).

Further, when more deactivated aromatic amines were employed as one of the reactants, the reactions were performed at 115 °C for 24 hours using butyl acetate and with 1 mol% TA and 1 mol% [Emim]⁺[OAc]⁻. Respective products were formed in quantitative yield in most of the cases (**2a**, **2b**, **2c**, **2d**, **2e**; ESI-S6(a-e)†). In the synthesis of **2g**, however, the yield was reduced to $41 \pm 1.6\%$ due to additional deactivation by a pyridine ring (ESI-S6(g)†). This method was also successful in synthesizing

phenacetine (**2f**), an anti-analgesic API, with a $98.7 \pm 1.0\%$ yield (ESI-S6(f)†). In the next step, to check the variability of the approach, respective acetylating agents were changed from aliphatic to aromatic esters (methyl benzoate and methyl 3-nitrobenzoate). Under the same reaction conditions (with 1 mol% catalyst at 80 °C for 20 hours) using toluene as a solvent, $90.7 \pm 1.4\%$ of **3a** and $89 \pm 0.8\%$ of **3b** were obtained, respectively (ESI-S7(a and b)†) (Fig. 3).

From a mechanistic viewpoint, a weak acid like acetic acid is preferred as a catalyst in this reaction compared to strong acids (such as sulfuric acid) due to its preference for amide formation over salt formation, as shown before.^{24,27} Apart from its role in




Fig. 3 *N*-Acetylation of amines with 1 mol% catalytic load.



Table 2 Mechanochemical *N*-acetylation reaction

Reactant	Molar ratio	Catalyst (mol%)	Liquid-assisted grinding (LAG)					Product	Yield (%)
			Time (min)	Frequency (Hz)	Jar (mL)	No. of ball (10 mm)	η ($\mu\text{l mg}^{-1}$)		
A + B	(1 : 1)	0	60	30	10	1	0	NIL	0
A + B	(2 : 1)	0	60	30	10	1	1.81	NIL	0
A + B	(1 : 1)	30	60	30	10	1	1.81	1f	100
A + B	(1 : 1)	15	30	30	10	1	1.81	1f	100
A + B	(1 : 1)	5	30	30	10	1	1.81	1f	100
A + B	(2 : 1)	30	90	30	10	2	1.81	1f	100
A + B + C	(1 : 1 : 1)	5	30	30	10	1	1.81	4a	56
A + B + D	(2 : 2 : 1)	5	30	30	10	1	1.81	4b	87

acid catalysis, acetic acid also plays a pivotal role in forming the respective transition state.²⁶ Hence, it may be possible that replacing direct acetic acid catalysis with an *in situ* approach lowers the requirement of minimum catalytic load (from 10 to 1 mol%) as an *in situ* process dictates an equilibrium between the cyclic intermediate²⁶ and acetic acid formation and enables a controlled release of acetic acid. Other organic acids were also used as catalysts to test this hypothesis, *e.g.*, oxalic acid (OA) and malonic acid (MA). In the presence of these strong acids (OA ($pK_a = 1.2$) and MA ($pK_a = 2.83$)) as catalysts, the salt formation pathway was favored, and only 17% and 27% **1a** were formed, respectively (ESI-S8(a and b)†). **1a**, however, was still formed with 100% yield when these two strong acids were mixed with [Emim]⁺[OAc]⁻ (ESI-S8(c and d)†). This indicates solid organic acids with a pK_a lower than acetic acid may be used with [Emim]⁺[OAc]⁻ to produce *in situ* acetic acid. This observation opens up an opportunity to apply it to mechanochemistry.^{28,29}

In the next step, we tried to extend the above approach to a mechanochemistry protocol. When hexamethylenediamine and methyl 3-nitrobenzoate were ground together in a mixer ball mill for 60 minutes at a frequency of 30 Hz in a 10 ml jar with a single 10 mm steel ball, without a catalyst, no product was formed (ESI-S9(a and b)†). In the presence of TA and [Emim]⁺[OAc]⁻ mixture with the above conditions, **1f** was

formed with 100% yield (ESI-S9(c)†). The experimental parameters were varied further for grinding conditions and catalytic load to check the robustness of the protocol. **1f** was formed exclusively (ESI-S9(d-f)† and (Table 2)). Since **1f** formation dictates one -NH₂ group of hexamethylenediamine being converted to an amide with the other -NH₂ remaining free, we thought to further utilize this mechanochemistry protocol in a multicomponent reaction.

When methyl 3-nitrobenzoate, hexamethylenediamine, and 3,5-dichlorobenzaldehyde were ground at 30 Hz in a 10 ml jar with one 10 mm ball and 5 mol% catalyst, **4a** was formed with a significant yield (ESI-S9(g)†). Next, **4b** was synthesized when terephthalaldehyde was used instead of 3,5-dichlorobenzaldehyde in the above protocol (ESI-S9(h)†) (Fig. 4).

In conclusion, a clue from a previously reported ionic co-crystal structure leads to the development of a robust protocol for *in situ* acetic acid catalysis for amide synthesis. The protocol was generalized with many examples of varying electronic and structural factors, including modifying the acylating agent from aliphatic to aromatic esters. Apart from showing catalytic efficiency, this protocol could be extended to the mechanochemistry synthesis of amides and a mechanochemical multicomponent reaction. Given the importance of amide derivatives in organic synthesis and medicinal chemistry, this robust protocol is expected to diversify the synthesized substrates through a greener route. In the end, this study shows that crystal structures, which were once conceived as “chemical cemeteries,”^{30,31} can provide meaningful insights to develop intriguing and useful protocols relevant to the other areas of the subject, such as greener synthesis,³² or organocatalysis.³³

Data availability

The data supporting this article have been included in the ESI.†

Author contributions

BS: data curation, formal analysis, investigation, methodology; MRA: validation, visualization, software, writing – original draft preparation; AM: conceptualization, project administration, funding acquisition, writing – review & editing, supervision.




Fig. 4 Reaction adopted for mechanochemistry synthesis of amides.



Conflicts of interest

There are no conflicts to declare.

Acknowledgements

BS acknowledges SERB (SRG/2022/000259) for a junior research fellowship (JRF). MRA acknowledges BITS Pilani for an institute fellowship. AM acknowledges SERB (SRG/2022/000259) for funding. We acknowledge Central Analytical Laboratory (CAL), BITS Pilani Hyderabad campus for instrumental facilities.

References

- 1 J. M. Robertson, *Helv. Chim. Acta*, 1972, **55**, 119–127.
- 2 J. Lehn, *Angew. Chem., Int. Ed.*, 1988, **27**, 89–112.
- 3 H. B. Burgi, J. D. Dunitz and E. Shefter, *J. Am. Chem. Soc.*, 1973, **95**, 5065–5067.
- 4 J. D. Dunitz, *Pure Appl. Chem.*, 1991, **63**, 177–185.
- 5 G. R. Desiraju, *J. Am. Chem. Soc.*, 2013, **135**, 9952–9967.
- 6 G. Bolla and A. Nangia, *Chem. Commun.*, 2016, **52**, 8342–8360.
- 7 A. Mukherjee, *Cryst. Growth Des.*, 2015, **15**, 3076–3085.
- 8 C. M. Reddy, G. Rama Krishna and S. Ghosh, *CrystEngComm*, 2010, **12**, 2296.
- 9 Z. Wang, W. Han, R. Shi, X. Han, Y. Zheng, J. Xu and X.-H. Bu, *JACS Au*, 2024, **4**, 279–300.
- 10 M. R. Ahsan, L. Singh, H. Varma and A. Mukherjee, *Chem. Commun.*, 2023, **59**, 12711–12714.
- 11 H. B. Buergi and J. D. Dunitz, *Acc. Chem. Res.*, 1983, **16**, 153–161.
- 12 V. S. S. Kumar, A. Addlagatta, A. Nangia, W. T. Robinson, C. K. Broder, R. Mondal, I. R. Evans, J. A. K. Howard and F. H. Allen, *Angew. Chem., Int. Ed.*, 2002, **41**, 3848–3851.
- 13 G. R. Desiraju, *Angew. Chem., Int. Ed.*, 2007, **46**, 8342–8356.
- 14 M. R. Ahsan, L. Singh, B. Sar and A. Mukherjee, *Cryst. Growth Des.*, 2024, **24**, 1695–1704.
- 15 S. Parveen, R. J. Davey, G. Dent and R. G. Pritchard, *Chem. Commun.*, 2005, 1531.
- 16 R. J. Davey, S. L. M. Schroeder and J. H. ter Horst, *Angew. Chem., Int. Ed.*, 2013, **52**, 2166–2179.
- 17 Q. Zeng, A. Mukherjee, P. Müller, R. D. Rogers and A. S. Myerson, *Chem. Sci.*, 2018, **9**, 1510–1520.
- 18 V. G. Saraswatula, S. Bhattacharya and B. K. Saha, *New J. Chem.*, 2015, **39**, 3345–3348.
- 19 S. Aerry, A. Kumar, A. Saxena, A. De and S. Mozumdar, *Green Chem. Lett. Rev.*, 2013, **6**, 183–188.
- 20 N. O. V. Sonntag, *Chem. Rev.*, 1953, **52**, 237–416.
- 21 H. Lundberg and H. Adolffson, *ACS Catal.*, 2015, **5**, 3271–3277.
- 22 B. C. Ranu and P. Dutta, *Synth. Commun.*, 2003, **33**, 297–301.
- 23 K. Arnold, B. Davies, D. Héroult and A. Whiting, *Angew. Chem., Int. Ed.*, 2008, **47**, 2673–2676.
- 24 D. D. Sanz Sharley and J. M. J. Williams, *Chem. Commun.*, 2017, **53**, 2020–2023.
- 25 H. M. Titi, S. P. Kelley, M. E. Easton, S. D. Emerson and R. D. Rogers, *Chem. Commun.*, 2017, **53**, 8569–8572.
- 26 H. Charville, D. A. Jackson, G. Hodges, A. Whiting and M. R. Wilson, *Eur. J. Org. Chem.*, 2011, **2011**, 5981–5990.
- 27 H. S. Prasad, G. R. Srinivasa and D. Channe Gowda, *Synth. Commun.*, 2005, **35**, 1189–1195.
- 28 T. Friščić, C. Mottillo and H. M. Titi, *Angew. Chem., Int. Ed.*, 2020, **59**, 1018–1029.
- 29 W. Jones and M. D. Eddleston, *Faraday Discuss.*, 2014, **170**, 9–34.
- 30 D. Braga, *Isr. J. Chem.*, 2017, **57**, 101–108.
- 31 J. D. Dunitz, *Trans. Am. Crystallogr. Assoc.*, 1984, **20**, 1.
- 32 K. Tanaka and F. Toda, *Chem. Rev.*, 2000, **100**, 1025–1074.
- 33 B. List, *Chem. Rev.*, 2007, **107**, 5413–5415.

