Chemical Science

EDGE ARTICLE

Cite this: Chem. Sci., 2024, 15, 4824

C All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 18th December 2023 Accepted 13th February 2024

DOI: 10.1039/d3sc06816a

rsc.li/chemical-science

Introduction

Epilepsy stands as one of the prevailing neurological disorders, afflicting approximately 70 million individuals worldwide and contributing to 0.7% of the global disease burden.¹ Recurrent seizures often lead to hippocampal damage, resulting in subsequent cognitive impairment.² Despite its prevalence, the mechanism underlying the pathogenesis of epilepsy remains inadequately understood. $SO₂$, as a pivotal neurochemical transmitter in the brain, can be internally synthesized from sulfur-containing amino acids. Variations in its concentrations have been linked to a spectrum of cerebral disorders.³ Previous investigations have posited the involvement of the sulfur dioxide $(SO₂)/$ aspartate aminotransferase (AAT) pathway in the genesis of epilepsy and associated cerebral injury.⁴ An abnormal elevation of the levels of SO₂ and its derivatives $\left(\mathrm{SO}_3^{-2-}/\mathrm{HSO}_3\right)^$ has been considered as a possible basis for apoptosis of hippocampal neurons in epilepsy. Additionally, research has also delineated that SO_2 exerts a protective effect against brain

Danying Lin,^a Tao Lu,^b Xiao Wang,^a Xiaoxue Ye^{*b} and Zhihong Liu D^{*ab}

Epilepsy is considered one of the most prevalent neurological disorders, yet the precise mechanisms underlying its pathogenesis remain inadequately elucidated. Emerging evidence implicates endogenous sulfur dioxide (SO_2) in the brain as playing a significant role in epilepsy and associated neuronal apoptosis. Consequently, tracking the dynamic fluctuations in the levels of $SO₂$ and its derivatives $(SO₃^{2–}/HSO₃⁻)$ provides valuable insights into the molecular mechanisms underlying epilepsy, with potential implications for its diagnosis and therapeutic intervention. Nonetheless, the absence of reversible in vivo detection tools constitutes a formidable obstacle in the real-time monitoring of $SO₂$ dynamics in the brain. In response to this challenge, we propose a novel approach involving a photoelectrochemical (PEC) microsensor capable of reversibly detecting SO₂. This microsensor leverages a reversibly recognizing dye for $SO₂$ and upconversion nanoparticles as the modulator of the excitation source for the photoactive material, enabling modulation of the photocurrent by the target. The reversible output of PEC signals allows for the monitoring of $SO₂$ levels in real time in the brains of epileptic mice. This study reveals the patterns of $SO₂$ level changes during epilepsy and provides insights into the neuroprotective mechanism of exogenous $SO₂$. EDGE ARTICLE
 (A) Check for unduses
 EDGE ARTICLE
 A reversible photoelectrochemical microsensor for the properties of the state of the state of the control of the state of the control of the state of the state of t

injury associated with epilepsy through various molecular mechanisms.⁵ It is evidently imperative to monitor the levels of endogenous $SO₂$ in the brain to gain insights into the pathogenic mechanisms of this disorder.⁶ Regrettably, the dearth of suitable in situ analysis tools has hindered our ability to track time-resolved fluctuations in $SO₂$ levels during epilepsy.

In vivo electrochemical analysis, utilizing implantable microelectrodes, represents a robust methodology for real-time monitoring of neurochemicals in the living brain, offering exceptional spatiotemporal resolution and sensitivity.⁷ More recently, photoelectrochemical (PEC) sensing has emerged as a promising tool for in vivo analysis. Its design, predicated on the chemical recognition of a small molecule dye, has proved to be particularly adept at selectively detecting electrochemically inert neurochemicals, such as $SO₂$, which are difficult to detect directly through faradaic signals.⁸ However, a major challenge associated with this method lies in monitoring dynamic changes in target levels over time, due to the fact that the vast majority of known recognition processes are irreversible, thereby only reflecting a monotonic increase in the target concentration. To date, only a limited number of conditional reversible dyes have been reported for sulfite detection, which typically necessitate the addition of sulfur dioxide scavengers or oxidants (e.g., formaldehyde, hydrogen peroxide) to facilitate the reverse process.⁹ Consequently, these reversible dyes fall short in dynamically monitoring sulfite in living organisms. Obviously, to achieve real-time monitoring of the fluctuating

a College of Chemistry and Molecular Sciences, Wuhan University, Wuhan 430072, China. E-mail: zhhliu@whu.edu.cn

b College of Health Science and Engineering, Ministry of Education Key Laboratory for the Synthesis and Application of Organic Functional Molecules, Hubei University, Wuhan 430062, China. E-mail: yexxchem@hubu.edu.cn

[†] Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d3sc06816a>

levels of SO_2 in vivo, the dye must be capable of swiftly and spontaneously undergoing a reversible reaction in response to both increasing and decreasing $SO₂$ concentrations without the need for additional reagents, including catalysts, oxidants or SO2 scavengers. Unfortunately, as of now, there have been no reported instances of a reversible dye capable of meeting this formidable challenge.

In this work, we have successfully engineered a reversible photoelectrochemical (PEC) microsensor, employing a spontaneous reversible dye founded on the Michael addition reaction, to dynamically track the fluctuations in $SO₂$ levels in the brains of epileptic mice. Initially, three dyes were designed and synthesized to enable reversible recognition of SO_2 . The effect of the electron cloud density of the recognition site on the reversibility of the dye was investigated. To accurately detect intracerebral SO_2 concentrations in the range of 5-25 μ M, a dissociation constant (K_d) value of approximately 10 μ M is required.¹⁰ Eventually, we identified an optimized dye, boasting a desirable K_d value of 9.6 μ M for detecting SO₂ in the brain. More importantly, the addition reaction between the dye and sulfur dioxide can occur spontaneously and reversibly. Afterwards, the selected dye underwent further modification by attachment to upconversion nanoparticles (UCNPs), thereby giving rise to a fluorescence resonance energy transfer (FRET)based nanoprobe. This FRET-based nanoprobe was subsequently co-modified with photoactive materials on Ti wires to construct a reversible PEC microsensor. Specifically, the FRET process led to the quenching of upconversion luminescence (UCL) which functioned as the excitation source of the photoactive materials, resulting in a low photocurrent of the PEC sensor. The introduction of $SO₂$ caused a reduction in FRET efficiency, thereby increasing both the UCL and photocurrent. The reversible nature of the nanoprobe allowed it to Edge Article

Levela Steine

Levela Steine

Levela Steine

Top minimals are complexe are considered on the property are considered to the distinguished on 2022. The properties

med to medicinal regress Article is licensed

spontaneously form a free state as $SO₂$ levels decreased, again leading to a reduced PEC signal (Scheme 1). The proposed microsensor showed excellent reversible response both in vitro and in vivo, and successfully monitored $SO₂$ in three brain regions of epileptic mice, revealing the law of $SO₂$ fluctuations and providing insights into the neuroprotective mechanism of exogenous SO₂.

Results and discussion

Design and response of the reversible molecular recognizers

We employed the concept of the Michael addition reaction, which involves the conjugated addition between an electrophilic conjugated system and a nucleophilic electron donor, to design reversibly recognizing dyes for SO₂. The reversibility and sensitivity of the dyes, represented by K_d , can be adjusted by modifying the molecular structure to manipulate the electron density. Herein, we rationally designed and synthesized three novel SO_2 reversible dyes named rSO_2 -1, rSO_2 -2, and rSO_2 -3, respectively (Scheme 2). The benzopyran unit was selected as the recognition site, and three different electron donors including p -phenylenediamine, p -phenylamine, and p -phenylacetamido were introduced to adjust the charge density of the recognition site, thus endowing the molecules with flexible reversible properties. The successful synthesis of $rSO₂$ dyes was confirmed by 1 H NMR, 13 C NMR and HRMS (Fig. S1-S9 \dagger). To verify the reaction mechanism of $rSO₂$ toward $SO₂$, specifically taking rSO_2-2 as the representative, the reaction products were characterized by mass spectrometry and ¹H NMR. The observed peak at m/z 397.1249 [M + Na]⁺ in the mass spectrum indicated the molecular weight to be 374, corresponding to the formula $C_{19}H_{22}N_2O_4S$, as expected. Changes in the chemical shift calculated in the 1 H NMR spectrum suggested a structural

Scheme 1 Schematic illustration of the response mechanism of the PEC microsensor for reversible sensing of $SO₂$.

Scheme 2 The molecular structures of $rSO₂-1$, $rSO₂-2$, and $rSO₂-3$, as well as the response mechanism of $rSO₂$.

alteration in rSO_2-2 . The obtained results support the proposed mechanism outlined in Scheme 2. Fig. 1a illustrates the absorption spectra of rSO_2-1 , rSO_2-2 , and rSO_2-3 , showing a sequential blue-shift. This shift is consistent with the decreasing electron-donating ability of dimethylamino, amino, and acetamido connected to the conjugated system. Consequently, the electrophilic abilities of the recognition sites were enhanced due to the higher charge densities (0.02627, 0.03019, and 0.03349, as calculated by the extended Huckel method) (Fig. 1b). Since SO₂ exists in the form of HSO₃ $^{-}$ and SO₃^{2–} (1:3,

M/M) in organisms, SO $_3^{2-}$ was used for *in vitro* experiments. To evaluate the reversibility of $rSO₂$ dyes, dilution experiments were carried out. Results from the 10-fold dilution indicate a rapid and reversible response of rSO_2 -1 (Fig. 1c), as well as $rSO₂$ -2 (Fig. 1d), which is predominantly dependent on the dose of SO_2 . r SO_2 -3 (Fig. 1e) exhibited poor dose-dependent reversibility, possibly due to the strong cohesion caused by the weak electron-donating ability of the acetylamino group. These dyes demonstrated exceptional selectivity towards $SO₂$, without any interference from ROS, H_2S , 40 μ M GSH and other biothiols, whereas a high concentration of GSH (10 mM) noticeably affected their selectivity (Fig. $S11\dagger$). These findings indicate the potential use of these dyes for detecting $SO₂$ in the intercellular fluid, as the low concentration of GSH in this fluid does not compromise the selectivity of the dye.¹¹ However, it is not suitable for detecting intracellular $SO₂$ due to the substantial influence of the high concentration of GSH within the intracellular environment.¹² Fig. 1f and g depict the gradual decrease in absorption of rSO_2-1 and rSO_2-2 as the concentration of SO_2 increased. Additionally, the widely used mercaptan scavenger N-methylmaleimide (NMM) was used to explore the reversibility of the reaction. Upon NMM being added to the dye solutions pretreated with SO_2 , the UV-Vis absorption was restored to varying degrees since SO_2 was removed (Fig. 1h and i). The calculated K_d values for rSO₂-1 and rSO₂-2, based on the slope of Chemical Science

Comparison Access Article

Comparison Access Article is license article. The commons are computed to the commons are computed under a Creative Commons are commons are commons are commons are commons are

Fig. 1 (a) Absorption spectra of rSO₂-1, rSO₂-2, and rSO₂-3. (b) Charge density at the recognition site of rSO₂-1, rSO₂-2, and rSO₂-3 calculated by the extended Huckel method. (c–e) The initial mixtures of SO₂ and rSO₂-1 (rSO₂-1/SO₂) (c), rSO₂-2 (rSO₂-2/SO₂) (d), and rSO₂-3 (rSO₂-3/SO₂) (e) were diluted 10-fold into the PBS containing no or 10 μ M SO₂. (f and g) UV-vis response of rSO₂-1 (f) and rSO₂-2 (g) upon addition of SO₂ (0– 200 µM SO₂ in rSO₂-1 and 0-100 µM SO₂ in rSO₂-2). (h and i) Reversible changes in the absorption spectra of rSO₂-1/SO₂ (h) and rSO₂-2/SO₂ (i) upon addition of NMM (0–1 mM). (j and k) Absorbance of rSO₂-1 (j) and rSO₂-2 (k) as a function of SO₂ concentration (0–20 µM)

the absorbance-concentration line, were 21.3 μ M and 9.6 μ M, respectively (Fig. 1j and k). Considering the physiological level of $SO₂$ in the cerebrospinal fluid (CSF) (within the range of 5–25 μ M),¹⁰ rSO₂-2 is more suitable for the reversible analysis of SO₂ in the brain.

Properties of the $rSO₂$ -UCNPs nanoprobe

Furthermore, PAA-modified NaYF₄:Yb,Er UCNPs were synthesized with the intention of constructing a FRET-based nanoprobe in combination with rSO_2-2 . Transmission electron microscopy (TEM) images revealed that the UCNPs exhibited a uniform size distribution, with an average diameter of about 37 nm (Fig. S12†). X-ray diffraction (XRD) patterns confirmed that the obtained nanoparticles possessed a hexagonal phase (Fig. S13†). Importantly, the absorption band of rSO_2 -2 fits nicely with the emission band of UCNPs at 545 nm, facilitating efficient energy transfer from UCNPs to rSO_2 -2 and UCL quenching (Fig. 2a). FRET occurred between UCNPs and rSO_2-2 when the positively charged $rSO₂-2$ assembled on the surface of the negatively charged UCNPs via electrostatic interaction. The zeta potential shifted from -30 mV for bare UCNPs to -20 mV

for rSO_2 -UCNPs, indicating the binding of rSO_2 -2 to UCNPs (Fig. $S14\dagger$). As anticipated, the degree of fluorescence quenching increased with the loading of rSO_2-2 , reaching a maximum quenching efficiency of 95% with a dose of 100 μ M rSO₂-2 (Fig. 2b). In the presence of SO_2 , FRET was inhibited, leading to the recovery of UCL intensity (Fig. 2c). Subsequent addition of NMM removed sulfite from the solution, resulting in dissociation of sulfite from the adducts, accompanied by the reoccurrence of FRET and UCL quenching (Fig. 2d). The above results implied that $rSO₂$ -UCNPs exhibited a reversible response to sulfite, with significant changes in the UCL intensity of UCNPs at 545 nm, which can be exploited to modulate the photocurrent signal.

Characteristics of the PEC microsensor

A 70 µm-diameter titanium wire was packaged in a glass capillary with the exposed length at $600 \mu m$ (denoted as TiME, Fig. S15†), which served as the working electrode. To improve the measurement depth in vivo, a $200 \mu m$ optical fiber was inserted into the front end of the drawn glass capillary tube and encapsulated with optical glue (Fig. 2e). This configuration

Fig. 2 (a) Normalized fluorescence spectrum of UCNPs excited at 980 nm and the absorption spectra of $rSO₂$ and CdTe QDs. (b and c) Fluorescence spectra of rSO₂-UCNPs upon addition of different amounts of rSO₂ (b) and SO₂ (c). (d) Fluorescence spectra of rSO₂-UCNPs with the removal of SO₂. (e) Schematic illustration of the architecture of designed microelectrode TiME. (f) Fluorescence images of microelectrodes with different coatings after adsorption of FITC-BSA (1%) for 8 h: rSO₂-UCNPs/CdTe-MWCNTs/TiME (top) and rSO₂-UCNPs/CdTe-MWCNTs/ BSA-GA/TiME (bottom). (g) Photocurrent signals obtained at RPMS in aCSF containing 1% BSA. (h) Photocurrent of bare and modified microelectrodes excited at 980 nm in aCSF: (1) TiWE, (2) CdTe-MWNTs/TiME, (3) UCNPs/CdTe-MWNTs/TiME, and rSO₂-UCNPs/CdTe-MWNTs/TiME before (4) and after (5) incubation with 20 μ M SO₂. (i) Reversible cycles observed in the PEC response of RPMS in aCSF without or with 20 μ M SO₂. (j) Photocurrent obtained from the reversible cycles described in (i). (k) Photocurrent response of RPMS in aCSF toward varying concentrations of SO_2 . (I) Selectivity test of SO_2 with interfering substances.

Chemical Science **Edge Article**

allowed stable transfer of the light via the optical fiber and optical glue, enabling in situ excitation and detection of deep brain regions. Then, photoactive materials composed of CdTe QDs and acidied multiwall carbon nanotubes (CdTe-MWNTs) were prepared (Fig. S16-S18†), and modified on the tip of TiME through titanium-carboxyl interaction.¹³ The resulting microphotoelectrode was referred to as CdTe-MWNTs/TiME for generating photocurrent. Next, the $rSO₂$ -UCNPs nanoprobe was assembled on the surface of CdTe-MWNTs/TiME by immersion and drying to fabricate the reversible PEC sensing platform $(rSO₂-UCNPs/CdTe-MWNTs/TIME)$. It is worth noting that biofouling of implanted microelectrodes can negatively impact their sensing performance by causing passivation. To address this issue, an anti-biofouling modification was applied. Specifically, a three-dimensional bovine serum albumin-glutaraldehyde (BSA-GA) film was introduced to enhance the anti-fouling properties of the microelectrode, and the entire reversible PEC microsensor was obtained (referred to as RPMS).¹⁴ The hydrophilicity of the modified surface was confirmed through contact angle tests (Fig. S19†). Additionally, protein adsorption tests were carried out, demonstrating that the BSA-GA coating significantly reduced protein adsorption (Fig. 2f). After immersing in a BSA solution for 8 hours, the photocurrent response of RPMS remained at 91.9% of its initial response (Fig. 2g), indicating the effectiveness of the anti-fouling modi fication. Scanning electron microscopy (SEM) (Fig. S20†) confirmed the step-by-step modification process and revealed the distribution of Cd, Te, Na, Y, F, Yb, Er, O, C, N, and S on the surface of RPMS, indicating the successful modification of TiME. Then, the PEC performance of the aforementioned RPMS was characterized to validate the capability of the $rSO₂$ -UCNPs nanoprobe as an excitation source to trigger CdTe-MWNTs to generate photocurrent signals and its response to sulfite. As depicted in Fig. 2h, a minimal photocurrent response of 8 nA (curve 2) was observed for CdTe-MWNTs/TiME under 980 nm light irradiation, as CdTe-MWNTs have limited absorption in the NIR light range. In contrast, the as-prepared UCNPs/CdTe-MWNTs/TiME showed the largest photocurrent response of 90 nA (curve 3), due to the effectively excited CdTe-MWNTs by the UCL of UCNPs at 545 nm. After that, the introduction of $rSO₂$ quenched the UCL, resulting in a 65% drop in photocurrent compared to the UCNPs/CdTe-MWNTs/TiME (curve 4). Finally, the incubation of sulfite at a concentration of 20 μ M resulted in a sharp increase in photocurrent (curve 5) as the FRET process was inhibited, leading to the recovery of UCL and an increase in the photocurrent signal. Control experiments were conducted by treating UCNPs/CdTe-MWNTs/TiME in artificial cerebrospinal fluid (aCSF) with and without 50 μ M sulfite, respectively, to exclude the influence of sulfite on other electrode modification materials (Fig. S21†). These results validated the principle and effectiveness of our sensing strategy. Chemical Science

Since Article. The science of the Euck view theorem are universal for example in the science of th

Reversible response and analytical performance of the microsensor towards $SO₂$

The reversibility of the microsensor RPMS was evaluated to assess its ability for real-time monitoring of dynamic fluctuations in SO_2 levels. Fig. 2i and j show the reversibility analysis by incubating the RPMS in $SO₂$ -free aCSF and aCSF containing 20 μ M SO₂. It was observed that the photocurrent response sharply increased when the RPMS was incubated with aCSF containing SO₂, while a decreased response was observed with SO_2 -free aCSF. Additionally, the reversibility of the RPMS was maintained after 5 cycles, indicating its robust performance. Under optimal conditions (Fig. S22†), the reversibility of RPMS was further explored (Fig. S23†). As expected, the PEC signal gradually enhanced with the addition of SO_2 , and a gradual drop in the PEC signal was observed upon stepwise dilution of the SO_2 solution with aCSF. Moreover, a linear correlation between the photocurrent and $SO₂$ concentration in the range of $3-25 \mu M$ was established, with a detection limit of 0.14 μ M, ensuring quantitative detection (Fig. 2k). The fast response of 120 s for recognition and 150 s for the reversible process enabled real-time and dynamic monitoring of $SO₂$ levels (Fig. $S24\dagger$). Benefiting from the high selectivity of rSO_2 , the excellent specificity for SO_2 was demonstrated by co-incubating the RPMS with various coexisting species in the biosystem (Fig. 2l). Ascorbic acid (AA), a common electron donor in the biosystem with a low redox potential, has the potential to affect the anode photocurrent through redox reactions.¹⁵ Although the photocurrent can be affected by low AA levels, negligible effect was found when AA reached over 100 μ M (Fig. S25†). Considering the physiological levels of AA (200 \pm 50 μ M), aCSF with 200 µM AA was used in in vitro experiments. RPMS also showed good photostability after continuous off-on-off irradiation (Fig. S26 and S27†), as well as good reproducibility for independent electrodes (Fig. S28†).

Reversible monitoring of $SO₂$ in the brain

As demonstrated above, the developed RPMS exhibited satisfactory reversible properties, as well as remarkable analytical performance, providing a reliable in vivo analysis platform for dynamic monitoring of $SO₂$ in the brain of living animals. Before the in vivo application, toxicity of RPMS was assessed by evaluating the cytotoxicity of the electrode modification material. HBMEC cells were incubated in dishes coated with CdTe-MWCNTs/rSO₂-UCNPs/BSA-GA film for 48 h. Subsequently, cell viability was determined through co-staining with calcein AMpropidium iodide. The utilization of confocal microscope imaging demonstrated favorable cell growth on the surface coated with CdTe-MWCNTs/rSO₂-UCNPs/BSA-GA (Fig. S29†), thus indicating good biocompatibility of the materials.

Furthermore, the standard deviation of RPMS in five repeated measurements was less than 5% (Fig. S30†). These results further confirm the excellent performance of the RPMS for in vivo detection.

In the first instance, RPMS was used to detect exogenous $SO₂$ in living mouse brain. The mice received in situ injection of $Na₂SO₃$ in the hippocampus (CA1) followed by the scavenger NMM. PEC signals were recorded after each injection. As depicted in Fig. 3a and b, a signicantly elevated PEC signal following the in situ injection of $Na₂SO₃$ was observed, indicating a robust response towards SO_2 in vivo. Subsequently,

Fig. 3 (a) Reversible photocurrent responses recorded at RPMS in CA1 of a live mouse brain following local microinjections (1 µL min⁻¹ for 60 s) of Na₂SO₃ and NMM in a continuous manner. (b) Photocurrents obtained from the reversible responses described in (a). (c) SO₂ levels (red) and AAT activity (black) in the hippocampus of epileptic mice pretreated with HDX. (d) Time-dependent PEC signals obtained with RPMS in CA1 of acute SO₂ poisoning mice. (e) Photocurrents obtained from (d). (f) Matrix map of photocurrents in the CA1, M1 and CPU regions of epileptic mice. (g) Analysis of $SO₂$ levels and AAT activity in the CA1, M1 and CPU regions of epileptic mice.

NMM was injected to eliminate SO_2 , resulting in a noticeably attenuated PEC signal. The RPMS successfully recorded signals from three consecutive $\text{Na}_2\text{SO}_3/\text{NMM}$ injections, and the photocurrent response varied in accordance with the levels of SO2, demonstrating the excellent reversible detection capability of SO_2 in vivo by the RPMS.

Subsequently, the ability of RPMS to dynamically monitor $SO₂$ in vivo was further investigated in a mouse model of acute SO2 poisoning. The mice were randomly divided into two groups: one group received injection of $Na₂SO₃$ as the $SO₂$ source, while the other group received saline as a control. As displayed in Fig. 3d and e, an intraperitoneal injection of Na2SO3 induced an elevated photocurrent response of RPMS implanted in CA1 over time, but no remarkable shift followed the injection of saline, suggesting that the increase in $SO₂$ level in the mouse brain was specifically caused by the rapid absorption of $Na₂SO₃$. In addition, the level of $SO₂$ in the brain reached its maximum at 1.5 h after poisoning and then slowly decreased. However, upon the injection of NMM, there was a dramatic decrease in photocurrent. The above results again

validated the capability of RPMS to dynamically monitor and track real-time changes of $SO₂$ levels in vivo.

Tracking the $SO₂$ level in the brain of epileptic mice

Epilepsy is a neurological disorder accompanied by abnormal discharges, and can cause irreversible damage to the brain. While previous studies have indicated the involvement of endogenous SO_2 in epilepsy and associated neuronal apoptosis,^{5,16} limited research tools impeded the investigation of the dynamics of SO_2 fluctuations in the brain throughout the course of epilepsy. In the present study, we aimed to investigate the real-time changes in endogenous $SO₂$ levels in the extracellular fluid of the living brain during epilepsy. Firstly, the kainic acid (KA)-induced status epilepticus (S.E.) model was established according to the existing literature.¹⁷ Then, RPMS was utilized to monitor intercellular $SO₂$ levels in different brain regions including M1, CPU and CA1 (Fig. 3f, S31†). The results revealed a gradual increase in $SO₂$ levels in all three brain regions after epilepsy, with the highest levels at approximately 3 h. SO₂ levels in the CPU region reach their peak slightly earlier than those in the CA1 and M1 regions. The photocurrent response in M1, CPU and CA1 rose to 151%, 158% and 216%, respectively. Notably, CA1 exhibited the highest magnitude of photocurrent rise compared to M1 and CPU during S.E., indicating that CA1 was the most affected brain region. Moreover, the CA1 region demonstrates the most rapid elevation rate, while the elevation rates of SO_2 levels in the CPU and M1 regions are comparable but slower (Fig. S32†). The above results suggested that during epilepsy, the $SO₂$ levels in different brain regions undergo a significant increase to varying degrees with CA1 being the most prominently affected and underscored the heterogeneity in endogenous $SO₂$ alterations across distinct cerebral regions during epilepsy.

As an endogenous gaseous molecule, $SO₂$ can be endogenously generated from sulfur-containing amino acids by an AAT dependent pathway.¹⁸ To gain further insight into the $AAT/SO₂$ pathway in epilepsy, we determined AAT activity in three brain regions (CA1, M1 and CPU). As can be seen from Fig. 3g, the variation trend of AAT activity was in harmony with that of $SO₂$ levels. It was reasonable to predict that the increased $SO₂$ production is preceded by an upregulation of AAT activity. To validate this hypothesis, a second group of mice was pretreated with AAT inhibitor L-aspartate-hydroxamate (HDX). As expected, the inhibition of AAT activity led to a significant decrease in both the SO_2 levels and AAT activity in CA1 (Fig. 3c). These findings provide further evidence that the upregulation in AAT activity in epilepsy is one of the major factors contributing to the increase of $SO₂$ content in the brain of epileptic mice. The results support the involvement of the $AAT/SO₂$ pathway in epileptic processes and highlight the potential signicance of AAT as a target for therapeutic interventions aimed at modulating $SO₂$ levels in epilepsy.

Neuroprotective mechanism of exogenous sulfur dioxide

Low-concentration SO_2 pretreatment has been reported to have a neuroprotective effect against neurological injuries.⁵ In investigating this neuroprotective effect and related mechanism in the context of epilepsy for the first time, mice were subjected to an experiment involving the administration of a low concentration of SO₂ (25 µmol kg^{-1}) 30 minutes prior to KA injection. TUNEL and HE staining were performed to evaluate the neuronal injury in the hippocampus. Fig. 4a and b depict that, in contrast to the S.E. group, the SO_2 -pretreated S.E. group exhibited comparatively minor neuronal damage, indicating the protective efficacy of $SO₂$ pretreatment against epilepsy-induced neuronal injury. Furthermore, the inhibition of AAT by HDX exacerbated epilepsy-induced neuronal injury, underscoring the crucial role of SO_2 in neuronal protection from such injuries. To delve further into the neuroprotective mechanism, the recording of SO_2 levels in the CA1 region of the hippocampus was conducted using RPMS. The results demonstrated a swi increase in the SO_2 level following SO_2 pretreatment, as illustrated in Fig. 4c and S33.† Subsequent to KA-induced epilepsy Chemical Science

Open Access Article

Chemical Science Article

methods are a computed to the non-Distribution-Non-Commons are a subset of the solelow are all the solelow in the solelow in the solelow in the solelow in th

Fig. 4 (a) TUNEL staining of CA1 in the mice with seizures for 3 h pretreated with SO₂ and HDX. (b) HE staining of CA1 in the mice with seizures for 3 h pretreated with SO₂ and HDX. Yellow arrows represent inflammatory cells, red arrows represent necrotic neurons. (c) Time-dependent photocurrents obtained using RPMS in the CA1 region with different treatments: KA-induced epilepsy (S.E.) and KA-induced epilepsy with the pretreatment of SO₂ (25 µmol kg^{−1}) (SO₂ + S.E.). (d) AAT activity in the hippocampus of epileptic mice pretreated with SO₂ at different time points (*p < 0.05, **p < 0.01, and ***p < 0.001). (e) Total antioxidant capacity (T-AOC) in the hippocampus of the mice with seizures for 3 h pretreated with SO₂ and HDX. (f) MDA content in the hippocampus of the mice with seizures for 3 h pretreated with SO₂ and HDX.

induction, the SO_2 level continued to increase, albeit with reduced magnitude compared to the S.E. group. Throughout the recording duration, the average sulfur dioxide concentration was higher than that of the control group but lower than that of the S.E. group. To explore the underlying cause of this disparity, we measured AAT activity in the hippocampus at different time points (Fig. 4d). The outcomes revealed an inhibition of AAT activity post SO_2 pretreatment, with the lowest levels of activity being observed after 2 hours and progressively recovering afterwards. These findings suggest that the restrained increase in SO_2 levels in SO_2 -pretreated epileptic brains can be attributed to the inhibition of AAT activity by exogenous SO₂.

However, the question arises as to what may be the mechanisms for this neuroprotective effect of low SO₂ pretreatment. Oxidative stress is one of the primary mechanisms implicated in S.E.-induced neuronal damage.⁵ Sulfur dioxide pretreatment has been found to enhance antioxidant capacity during myocardial ischemia.¹⁹ Therefore, we hypothesize that the neuroprotective mechanism of $SO₂$ pretreatment in epilepsy may be associated with its augmentation of antioxidant capacity. To verify this hypothesis, we examined the total antioxidant capacity (T-AOC) (Fig. 4e) and MDA (a biomarker for oxidative stress) (Fig. 4f) levels in the brains of different mouse groups. The results revealed a correlation between the degree of neuronal damage and oxidative stress, indicating that oxidative stress is a major contributing factor in the model of neuronal damage induced by epilepsy. More importantly, the antioxidant capacity of the epilepsy group pretreated with $SO₂$ significantly improved, while the HDX group exhibited the lowest antioxidant capacity, aligning with the findings from TUNEL and HE staining. These results underscore the pivotal role of $SO₂$ in protecting against brain injury induced by epilepsy, highlighting that low concentrations of $SO₂$ pretreatment contribute to a neuroprotective effect by upregulating antioxidant capacity in the brain. Edge Article

includes Cheme access the control on 22 February 2022. The control of the control of the control on the second on 13 February 2024. The control of the control on the second one of the common and the second on

Conclusions

The developed reversible PEC microsensor (RPMS) based on a FRET-modulated signal has demonstrated its effectiveness for dynamic monitoring of $SO₂$ levels in the living mouse brain. The utilization of the Michael addition reaction between $rSO₂$ and SO2 enables reversible detection and ensures high selectivity of RPMS. The biocompatibility of the microsensor has been confirmed, and it has exhibited excellent analytical performance both in vitro and in vivo, making it suitable for real-time monitoring in living brains. By utilizing this microsensor, the study effectively demonstrated the variation of endogenous $SO₂$ in distinct cerebral regions during epilepsy, along with elucidating the neuroprotective mechanism of low concentration exogenous SO_2 in mitigating neuronal damage induced by epilepsy. This work introduces a novel tool for recording $SO₂$ fluctuations in living brains, which has the potential to contribute to the understanding of molecular-level mechanisms of brain function and the development of treatments for various neurological disorders.

Author contributions

D. Lin: most of the experiments and writing of the manuscript. T. Lu and X. Wang: in vivo experiments. X. Ye: supervision of experiments, discussion, and writing of the manuscript. Z. Liu: writing, review, and editing of the manuscript, funding acquisition and supervision.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors acknowledge the financial support from the National Natural Science Foundation of China (Grant No. 22276047, 22227804). All animal studies were performed according to the Guidelines for the Care and Use of Laboratory Animals of the Chinese Animal Welfare Committee and approved by the Institutional Animal Care and Use Committee, Wuhan University Center for Animal Experiment, Wuhan, China.

References

- 1 G. K. Mbizvo, K. Bennett, C. R. Simpson, S. E. Duncan and R. F. M. Chin, Epilepsy Res., 2019, 157, 106192.
- 2 (a) Y. Xiong, H. Zhou and L. Zhang, Brain Res., 2014, 1557, 190–200; (b) U. Sayin, T. P. Sutula and C. E. Stafstrom, Epilepsia, 2004, 45, 1539–1548.
- 3 H. Yan, Y. Wang, F. Huo and C. Yin, J. Am. Chem. Soc., 2023, 145, 3229–3237.
- 4 M. Niu, Y. Han, Q. Li and J. Zhang, Neurosci. Lett., 2018, 665, 22–28.
- 5 Y. Han, W. Yi, J. Qin, Y. Zhao, J. Zhang and X. Chang, Neurosci. Lett., 2014, 563, 149–154.
- 6 W. Zhang, F. Huo, Y. Yue, Y. Zhang, J. Chao, F. Cheng and C. Yin, J. Am. Chem. Soc., 2020, 142, 3262–3268.
- 7 (a) Y. Wang, Y. Qian, L. Zhang, Z. Zhang, S. Chen, J. Liu, X. He and Y. Tian, J. Am. Chem. Soc., 2023, 145, 2118–2126; (b) F. Zhu, Y. Xue, W. Ji, X. Li, W. Ma, P. Yu, Y. Jiang and L. Mao, Angew. Chem., Int. Ed., 2023, 62, e2022124; (c) C. Yang, Q. Cao, P. Puthongkham, S. T. Lee, M. Ganesana, N. V. Lavrik and B. J. Venton, Angew. Chem., Int. Ed., 2018, 57, 14255–14259; (d) X. He and A. G. Ewing, J. Am. Chem. Soc., 2022, 144, 4310–4314; (e) E. F. Megan and R. M. Wightman, Pharmacol. Rev., 2017, 69, 12; (f) C. Wang, X. Bi, X. Zhao, M. Zhou, Y. Ding and Y. Lin, ACS Sustain. Chem. Eng., 2019, 7, 3715–3721; (g) A. Meiller, E. Sequeira and S. Marinesco, Anal. Chem., 2020, 92, 1804–1810; (h) K. Zhang, H. Wei, T. Xiong, Y. Jiang, W. Ma, F. Wu, P. Yu and L. Mao, Chem. Sci., 2021, 12, 7369–7376.
- 8 X. Ye, X. Wang, Y. Kong, M. Dai, D. Han and Z. Liu, Angew. Chem., Int. Ed., 2021, 60, 11774–11778.
- 9 (a) H. Zhu, X. Zhang, C. Liu, Y. Zhang, M. Su, X. Rong, X. Wang, M. Liu, X. Zhang and W. Sheng, Sens. Actuators, B, 2022, 366, 131962; (b) Y. Zhang, L. Guan, H. Yu, Y. Yan,

L. Du, Y. Liu, M. Sun, D. Huang and S. Wang, Anal. Chem., 2016, 88, 4426–4431.

- 10 Y. Kimura, N. Shibuya and H. Kimura, Br. J. Pharmacol., 2019, 176, 571–582.
- 11 (a) M. We˛grzynowicz, W. Hilgier, A. Dybel, S. S. Oja, P. Saransaari and J. Albrecht, Neurochem. Int., 2007, 50, 883–889; (b) K. Q. Do, A. Trabesinger, H. M. Kirsten-Krüger, C. J. Lauer, U. Dydak, D. Hell, F. Holsboer, P. Boesiger and M. Cuénod, Eur. J. Neurosci., 2000, 12, 3721–3728. Chemical Science

L. Du, Y. Liu, M. Sun, D. Huang and S. Wearg, Article. Published on 22 February 2024. Downloaded on 5.16:56 AM. This article. The Creative Commons Attribution-Non-Commons Attribution-Non-Commons Attributi
	- 12 (a) Z. Liu, X. Zhou, Y. Miao, Y. Hu, N. Kwon, X. Wu and J. Yoon, Angew. Chem., Int. Ed., 2017, 56, 5812–5816; (b) K. Aoyama, Int. J. Mol. Sci., 2021, 22, 5010.
	- 13 W. Wang, W. Feng, J. Du, W. Xue, L. Zhang, L. Zhao, Y. Li and X. Zhong, Adv. Mater., 2018, 30, 1705746.
- 14 J. Sabaté del Río, O. Y. F. Henry, P. Jolly and D. E. Ingber, Nat. Nanotechnol., 2019, 14, 1143–1149.
- 15 M. Zhang, K. Liu, L. Xiang, Y. Lin, L. Su and L. Mao, Anal. Chem., 2007, 79, 6559–6565.
- 16 A. Finegersh, C. Avedissian, S. Shamim, I. Dustin, P. M. Thompson and W. H. Theodore, Epilepsia, 2011, 52, 689–697.
- 17 P. Vincent and C. Mulle, Neuroscience, 2009, 158, 309–323.
- 18 (a) R. Shapiro, Mutagen, 1977, 39, 149–175; (b) D. Zhang, X. Wang, X. Tian, L. Zhang, G. Yang, Y. Tao, C. Liang, K. Li, X. Yu, X. Tang, et al., Front. Immunol., 2018, 9, 00882.
- 19 H. F. Jin, Y. Wang, X. B. Wang, Y. Sun, C. S. Tang and J. B. Du, Nitric Oxide, 2013, 32, 56–61.