

Cite this: *Chem. Sci.*, 2024, 15, 4114

All publication charges for this article have been paid for by the Royal Society of Chemistry

## Tunable C–H functionalization and dearomatization enabled by an organic photocatalyst†

Bohang An,<sup>a</sup> Hao Cui,<sup>a</sup> Chao Zheng,<sup>b</sup> Ji-Lin Chen,<sup>a</sup> Feng Lan,<sup>a</sup> Shu-Li You and Xiao ZhangReceived 7th January 2024  
Accepted 5th February 2024

DOI: 10.1039/d4sc00120f

rsc.li/chemical-science

C–H functionalization and dearomatization constitute fundamental transformations of aromatic compounds, which find wide applications in various research areas. However, achieving both transformations from the same substrates with a single catalyst by operating a distinct mechanism remains challenging. Here, we report a photocatalytic strategy to modulate the reaction pathways that can be directed toward either C–H functionalization or dearomatization under redox-neutral or net-reductive conditions, respectively. Two sets of indoles and indolines bearing tertiary alcohols are divergently furnished with good yields and high selectivity. The key to success is the introduction of isoazatruxene ITN-2 as a novel photocatalyst (PC), which outperforms the commonly used PCs. The ready synthesis and high modulability of isoazatruxene type PCs indicate their great application potential.

Aromatic compounds are among the most fundamental feedstock chemicals, with global production on a million metric ton scale annually. The development of practical methods to increase the valorization of these abundant substances is thus in continuing demand. As classical transformations based on aromatic compounds, C–H functionalization such as Friedel–Crafts alkylation and Minisci-type reaction constitutes a straightforward strategy to incorporate diverse functionalities on the aromatic skeletons.<sup>1</sup> In a complementary fashion, dearomatization converts planar aromatics into three-dimensional molecules with increased sp<sup>3</sup>-hybridized carbons and stereocenters by disturbing the aromaticity (Scheme 1, top). Notable strategies include Birch reduction, oxidative dearomatization of phenols, arene–alkene photocycloadditions, and others.<sup>2</sup> Both types of transformations have found broad applications in areas ranging from pharmaceuticals and agrochemicals to materials science.<sup>3</sup> Despite the great achievements made to date, the utilization of a single catalyst to promote tunable C–H functionalization and dearomatization processes from the same starting materials is underexplored. Elegant reports mainly

involve the substrate control.<sup>4</sup> The difficulties might arise from multiple functions required for one catalyst.


On the other hand, indoles and indolines are privileged scaffolds in alkaloid natural products and biologically active molecules.<sup>5</sup> Meanwhile, tertiary alcohols are important structural units that widely exist in a variety of biologically active natural products, pharmaceuticals, and agrochemicals (Scheme 1A).<sup>6</sup> The molecules containing both indole/indolines and tertiary alcohols might have great potential applications. In this regard, divergent syntheses of such compounds are highly desirable. We questioned whether photoredox catalysis<sup>7</sup> could engineer C–H functionalization and dearomatization selectively to afford two sets of targets in a catalytic and divergent fashion. It is envisioned that the excited photoredox catalyst (PC) could donate an electron to the ketone moiety. The resultant ketyl radical<sup>8</sup> would add to indole nuclei (*via* **I**), which is followed by protonation. Then, intermediate **II** might undergo single-electron oxidation to achieve aromatization upon loss of a proton. Alternatively, single-electron reduction of **II** would occur instead to deliver dearomatization products after protonation (Scheme 1B). Although conceptually simple in design, this strategy must contend with problems involving both reactivity and selectivity. Specifically, SET reduction of the ketone moieties is challenging, owing to the significantly negative reduction potentials [*e.g.*  $E_{1/2}^{\text{red}} = -2.11$  V vs. SCE for acetophenone].<sup>9</sup> A powerful photocatalyst is required to be capable of participating in both SET reduction and SET oxidation events for the C–H functionalization process as well as sequential SET reductions for the dearomatization process. In addition, the reaction might encounter the chemoselective

<sup>a</sup>Fujian Key Laboratory of Polymer Materials, Fujian Provincial Key Laboratory of Advanced Materials Oriented Chemical Engineering, College of Chemistry and Materials Science, Fujian Normal University, Fuzhou 350007, China. E-mail: zhangxiao@fjnu.edu.cn

<sup>b</sup>State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, Shanghai 200032, China. E-mail: zhengchao@sioac.ac.cn; slyou@sioac.ac.cn

† Electronic supplementary information (ESI) available: Experimental and copies of NMR spectra. CCDC 2271792, 2271797, 2271798, 2271810 and 2309597. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d4sc00120f>






Scheme 1 Tunable C–H functionalization and dearomatization.

competition from reduction<sup>10</sup> or pinacol C–C coupling.<sup>11</sup> Furthermore, the diastereoselective issues should be addressed when multiple stereocenters are generated.

Herein, we describe the first photocatalytic method for selective C–H functionalization and dearomatization by operating a distinct mechanism. Starting from the same indole-tethered ketones, intramolecular alkylation and reductive dearomatization have been achieved under redox-neutral and net-reductive conditions, respectively. Two sets of important nitrogen-containing heterocycles are obtained in good yields and excellent selectivity. A new organic photocatalyst, isoazatruxene **ITN-2**, is discovered to guarantee the success.

Our study was initiated with the investigation of C–H functionalization reaction using indole derivative **1a** as the substrate under irradiation with a Kessil lamp (Scheme 2A). Due to the significantly negative reduction potential of a ketone moiety, the highly reducing Ir(ppy)<sub>3</sub> [ $E_{1/2}^{M^+/M^*} = -1.73$  V vs. SCE]<sup>7c</sup> was first attempted. However, this photocatalyst failed to give the desired C2-alkylation product **2a**, which confirmed that direct SET reduction of a ketone moiety by excited-state Ir(ppy)<sub>3</sub><sup>\*</sup> is not feasible. Next, the frequently-used transition-metal-based and



Scheme 2 Performance of various photocatalysts in C–H functionalization and characterization of ITNs.

organic photocatalysts including Ir(ppy)<sub>2</sub>(dtbbpy)PF<sub>6</sub>, Ir[(dFCF<sub>3</sub>ppy)<sub>2</sub>(dtbbpy)]PF<sub>6</sub>, Ru(bpy)<sub>3</sub>Cl<sub>2</sub>·6H<sub>2</sub>O, 4CzIPN [1,2,3,5-tetrakis(carbazol-9-yl)-4,6-dicyanobenzene], PTH (N-




phenylphenothiazine), rhodamine B, eosin Y, fluorescein, rose bengal, methylene blue, and thioxanthen-9-one were systematically screened, unfortunately all leading to poor results (<15% yield).

Given our interest in organic electronic materials,<sup>12</sup> we turned attention to isoazatruxenes (ITNs), which can be readily synthesized *via* cyclotrimerization of indoles.<sup>13</sup> Despite the easy access, the application of ITNs in photocatalysis remained elusive. Thus, the photophysical and electrochemical properties of isoazatruxenes (ITNs) possessing varied substituents on the nitrogen atoms (ITN-1: R = H; ITN-2: R = <sup>n</sup>Bu; ITN-3: R = Ph) were measured and are summarized in Scheme 2B (Fig. S1–S18†).<sup>14</sup> In addition, DFT calculations (B3LYP-D3(BJ)/def2-SVP) were performed to obtain the HOMO/LUMO energies of ITNs (Fig. S20†).<sup>14</sup> While ITN-1 shows weak absorption in the visible region, the introduction of alkyl- or aryl groups results in a bathochromic shift with respect to the unsubstituted counterpart based on the ultraviolet-visible (UV-vis) spectroscopy, possibly due to the inductive effect of the alkyl groups and the conjugative effect of the aryl groups (Scheme 2C). Electron-excitation analyses on the basis of TD-DFT calculations suggested that the S<sub>0</sub>-to-S<sub>1</sub> excitation of ITNs was mainly contributed by the electron transition from the corresponding HOMOs to LUMOs. The calculated vertical excitation energies of ITNs qualitatively reproduced the bathochromic shift order of the highest absorption peaks (ITN-1 < ITN-3 < ITN-2) (Fig. S21†).<sup>14</sup> Notably, ITNs exhibit highly negative excited state reductive potentials (spanning from –2.17 to –2.54 V vs. SCE) as well as comparable excited-state lifetimes (4.15–5.84 ns), which allowed them to be promising candidates for promoting the above transformation (Scheme 2E).

To our delight, the C2 alkylation proceeded when ITNs were used as the photocatalysts. Particularly, ITN-2 bearing <sup>n</sup>butyl substituents on the nitrogen atoms outperformed all the other photocatalysts, providing the corresponding polycyclic indole **2a** in 86% yield with toluene as the solvent. The superior catalytic efficiency of ITN-2 in comparison with other ITNs might correspond to its enhanced visible-light absorption, which resulted in better visible photon harvesting. Furthermore, control experiments demonstrated that the reaction requires a photocatalyst and visible light.<sup>14</sup>


Having determined the optimal reaction conditions, we next examined the scope of C–H functionalization. As shown in Scheme 3, the tethered aromatic ketones bearing either electron-withdrawing or electron-donating substituents (–Cl, –F, –Ph, –Me, –Et, –<sup>n</sup>Pr, –<sup>i</sup>Pr, –<sup>t</sup>Bu, –OMe) on the *para*-position of arenes were well tolerated, delivering the corresponding polycyclic indoles (**2b–2j**) in good to excellent yields (71–92%). The structure of **2a** was determined unambiguously by X-ray crystallographic analysis.<sup>14</sup> Interestingly, the more challenging substrates with multi-electron-donating groups [2,4-(Me)<sub>2</sub>, 2,5-(Me)<sub>2</sub>, 3,4-(Me)<sub>2</sub>, 3,4-(OMe)<sub>2</sub>] that possess lower reduction potentials could be accommodated, highlighting the power of ITN-2 (**2k–2n**, 52–83% yields). Moreover, both naphthyl and thienyl-derived ketones were converted to their corresponding polycyclic indoles bearing tertiary alcohols (**2o** and **2p**) with high efficiency under the catalysis of ITN-2. This method is also compatible with the substrates bearing ethyl ester or acetyl



Scheme 3 Substrate scope for C–H functionalization.<sup>a,b</sup> <sup>a</sup>Reaction conditions: a solution of ITN-2 (5.2 mg, 5.0 mol%) and **1** (0.2 mmol, 1.0 equiv.) in toluene (2.0 mL, 0.1 M) was irradiated by a 427 nm Kessil lamp (40 W) at room temperature under a nitrogen atmosphere for 4–36 h. <sup>b</sup>Isolated yield.

groups at the C-3 position of indole skeletons (**2q–2r**, 82–87% yields). Finally, the reactions of substrates bearing cyano, fluorine, and methyl substituents at the indole cores also proceeded smoothly, providing **2s–2u** in 76–90% yields.




Table 1 Optimization of the reaction conditions for dearomatization<sup>a</sup>

Entry	Variations from standard conditions	Yield <sup>b</sup> (%)	Dr <sup>c</sup>
1	None	91 (90) <sup>d</sup>	>20 : 1
2	No DMAP	91	1.7 : 1
3	<b>ITN-1</b> instead of <b>ITN-2</b>	0	n.d.
4	<b>ITN-3</b> instead of <b>ITN-2</b>	59	>20 : 1
5	Acetone instead of $\text{CH}_2\text{Cl}_2$	37	16 : 1
6	$\text{CH}_3\text{CN}$ instead of $\text{CH}_2\text{Cl}_2$	13	n.d.
7	DMSO instead of $\text{CH}_2\text{Cl}_2$	8	n.d.
8	DIPEA instead of <b>HEH</b>	45	>20 : 1
9	No <b>ITN-2</b>	0	n.d.
10	No <b>HEH</b>	0	n.d.
11	No light	0	n.d.

<sup>a</sup> Reaction conditions: a solution of **ITN-2** (2.6 mg, 5.0 mol%), **1a** (0.1 mmol, 1.0 equiv.), **HEH** (0.15 mmol, 1.5 equiv.), and **DMAP** (1.2 mg, 10.0 mol%) in  $\text{CH}_2\text{Cl}_2$  (1.0 mL, 0.1 M) was irradiated by blue LEDs (30 W) at room temperature under a nitrogen atmosphere for 24 h. <sup>b</sup> Determined by <sup>1</sup>H NMR yield using  $\text{CH}_2\text{Br}_2$  as an internal standard. <sup>c</sup> Ratios of diastereoisomers were determined by crude <sup>1</sup>H NMR analysis. <sup>d</sup> Isolated yield. n.d. = not determined. DMAP = 4-dimethylaminopyridine. DIPEA = *N,N*-diisopropylethylamine.

Then, we continued to investigate the intramolecular dearomatization of indoles with ketones (Table 1). In fact, this process remains problematic because of energetic barriers caused by aromaticity and competition between dearomatization and reduction or pinacol C–C coupling (**1a** → **4a–4c**). The formation of multiple stereocenters also faces the challenge of diastereoselective control. With *N*-acylated indole derivative **1a** as the model substrate, our optimization revealed that the use of **ITN-2** as the photocatalyst, 4-dimethylaminopyridine (**DMAP**) as the additive, and Hantzsch ester (**HEH**) as the reductant in dichloromethane led to the formation of expected dearomatized product **3a** in 90% yield with excellent diastereoselectivity under blue LED irradiation at room temperature (entry 1). Notably, the presence of a catalytic amount of **DMAP** significantly improved the diastereoselectivity, likely a result of thermodynamic control (entry 2). Consistently, **ITN-1** and **ITN-3** afforded inferior efficiency (entries 3 and 4). Changing dichloromethane to other commonly used solvents such as acetone,  $\text{CH}_3\text{CN}$ , or DMSO resulted in dramatically decreased yields (8–37%) for the desired product **3a** (entries 5–7), demonstrating the important role of solvents. When **DIPEA** was used as the external reductant instead, the dearomatization reaction also occurred well, albeit with a lower yield of **3a** (entry



Scheme 4 Substrate scope for dearomatization.<sup>a,b</sup> Reaction conditions: a solution of **ITN-2** (5.2 mg, 5.0 mol%), **1** (0.2 mmol, 1.0 equiv.), **HEH** (0.3 mmol, 1.5 equiv.), and **DMAP** (2.4 mg, 10.0 mol%) in  $\text{CH}_2\text{Cl}_2$  (2.0 mL, 0.1 M) was irradiated by blue LEDs (30 W) at room temperature under a nitrogen atmosphere for 12–48 h. <sup>a</sup> Isolated yield. Ratios of diastereoisomers were determined by crude <sup>1</sup>H NMR analysis.




8). Lastly, the control experiments indicated that photocatalyst **ITN-2**, reductant HEH, and visible light are all indispensable (entries 9–11).

With the optimized conditions in hand, we explored the scope of photocatalytic intramolecular dearomatization of indole derivatives with ketones (Scheme 4). It was observed that the aromatic ketones bearing either electron-withdrawing (e.g. 4-Cl, 4-F) or electron-donating [e.g. 4-Me, 4-Et, 4<sup>n</sup>Pr, 4<sup>l</sup>Pr, 4<sup>t</sup>Bu, 4-OMe, 2,4-(Me)<sub>2</sub>, 2,5-(Me)<sub>2</sub>, 3,4-(Me)<sub>2</sub>, 3,4-(OMe)<sub>2</sub>] groups on the arene skeletons were well accommodated. In all cases, highly functionalized benzannulated indolizidines possessing three consecutive stereocenters were furnished in good to excellent yields with remarkable diastereoselectivity (**3a–3n**: 75–98% yields, 18 : 1 → >20 : 1 dr). Similarly, the structure and relative configuration of **3k** were confirmed by X-ray crystallographic analysis.<sup>14</sup> In addition, indole derivatives tethered with naphthyl and thienyl ketones were compatible with this transformation (**3o–3p**: 85–87% yields, >20 : 1 dr). Changing the substituents at the C-3 position of indoles to ethyl ester or acetyl groups also proved feasible (**3q–3r**: 73–85% yields, >20 : 1 dr). Furthermore, the reactions of substrates **1s–1u** containing varied substituents (–CN, –F, –Me) on the indole skeletons occurred smoothly, delivering the desired indolizidines **3s–3u** in good yields with extraordinary diastereoselectivity (86–96% yields, 18 : 1 → >20 : 1 dr).

In order to demonstrate the synthetic practicality, gram-scale syntheses were carried out (Scheme 5). With **ITN-2** as the photocatalyst, the C–H functionalization and dearomatization reactions of substrate **1a** on 4.47 mmol scale were selectively achieved under the optimized reaction conditions, yielding polycyclic indole **2a** and indoline derivative **3a** in 85% yield (1.28 g) and 91% yield (1.36 g), respectively.

Several experiments were performed to probe the possible mechanism (Scheme 6A). First, both reactions were subjected to ultraviolet light irradiation (365 nm) without the addition of **ITN-2**. It was observed neither of the reactions proceeded, thus excluding the operation of the energy transfer mechanism. Upon the addition of TEMPO (2,2,6,6-tetramethylpiperidine *N*-oxyl), the expected transformations were completely inhibited, which suggested the involvement of radicals. Accordingly, the plausible mechanism for photocatalytic C–H functionalization and dearomatization was proposed as exemplified by the reactions of **1a** (Scheme 6B).

Irradiation of **ITN-2** affords its excited state. Oxidative quenching of **ITN-2**\* by indole derivative **1a** generates **ITN-2**<sup>•+</sup> and the corresponding ketyl radical (**I**), which is supposed to undergo addition to electron-deficient indole. After protonation, intermediate **II** proceeds diversely in the two transformations. In the C–H functionalization reaction, single-



Scheme 6 Proposed mechanism.


electron oxidation of **II** by **ITN-2**<sup>•+</sup> delivers aromatization product **2a** upon the release of a proton and completes the catalytic cycle. With respect to the dearomatization reaction, single-electron reduction of **II** by excited state **ITN-2**\* occurs instead, yielding the benzannulated indolizidine **3a** after accepting a proton. **ITN-2**<sup>•+</sup> can be further converted back to **ITN-2** in the presence of an external reductant. While C–H functionalization is a redox-neutral transformation, dearomatization is a net-reductive process requiring a stoichiometric external reductant to turn over the photocatalytic cycle.

## Conclusions

In summary, we have developed a photocatalytic protocol to engineer C–H functionalization and dearomatization processes selectively. Crucial to the strategy are the different redox events promoted by a new organic photocatalyst – isoazatruxene **ITN-2**. Under redox-neutral conditions, intramolecular C–H functionalization is directed, providing polycyclic indoles bearing tertiary alcohols in good yields. Under net-reductive conditions, reductive dearomatization of indole derivatives with ketones can be realized instead, in which the corresponding benzannulated indolizidines possessing three consecutive stereocenters are furnished with remarkable diastereoselectivity. We anticipate this study will stimulate the development of novel photocatalysts for previously challenging transformations.

## Data availability

All experimental data is available in the ESI.†



Scheme 5 Gram-scale reaction.



## Author contributions

X. Z. conceived the project. B. A., H. C., J.-L. C., and F. L. performed the experiments under S.-L. Y. and X. Z.'s supervision. C. Z. performed the DFT calculations. B. A., C. Z., S.-L. Y., and X. Z. cowrote the paper.

## Conflicts of interest

A patent has been filed on the use of isoazatruxenes and their derivatives in photocatalysis.

## Acknowledgements

We thank NSFC (22071024, 22271047), and the Natural Science Foundation of Fujian Province (2021J06020) for generous financial support.

## Notes and references

- For selected reviews, see: (a) K. Murakami, S. Yamada, T. Kaneda and K. Itami, *Chem. Rev.*, 2017, **117**, 9302–9332; (b) A. A. Festa, L. G. Voskressensky and E. V. Van der Eycken, *Chem. Soc. Rev.*, 2019, **48**, 4401–4423; (c) G. Evano and C. Theunissen, *Angew. Chem., Int. Ed.*, 2019, **58**, 7558–7598; (d) R. S. J. Proctor and R. J. Phipps, *Angew. Chem., Int. Ed.*, 2019, **58**, 13666–13699; (e) P. Kumar, P. J. Nagtilak and M. Kapur, *New J. Chem.*, 2021, **45**, 13692–13746; (f) N. Holmberg-Douglas and D. A. Nicewicz, *Chem. Rev.*, 2022, **122**, 1925–2016; (g) S. Brunen, B. Mitschke, M. Leutzsch and B. List, *J. Am. Chem. Soc.*, 2023, **145**, 15708–15713.
- For selected reviews, see: (a) P. W. Rabideau and Z. Marcinow, *Org. React.*, 1992, **42**, 1–334; (b) R. M. Moriarty and O. M. Prakash, *Org. React.*, 2001, **57**, 327–415; (c) S. P. Roche and J. A. Porco Jr, *Angew. Chem., Int. Ed.*, 2011, **50**, 4068–4093; (d) C.-X. Zhuo, W. Zhang and S.-L. You, *Angew. Chem., Int. Ed.*, 2012, **51**, 12662–12686; (e) L. M. Repka and S. E. Reisman, *J. Org. Chem.*, 2013, **78**, 12314–12320; (f) S. P. Roche, J.-J. Y. Tendoung and B. Tréguier, *Tetrahedron*, 2015, **71**, 3549–3591; (g) M. J. James, P. O'Brien, R. J. K. Taylor and W. P. Unsworth, *Chem.–Eur. J.*, 2016, **22**, 2856–2881; (h) R. Remy and C. G. Bochet, *Chem. Rev.*, 2016, **116**, 9816–9849; (i) M. Okumura and D. Sarlah, *Synlett*, 2018, **29**, 845–855; (j) J. Bariwal, L. G. Voskressensky and E. V. Van der Eycken, *Chem. Soc. Rev.*, 2018, **47**, 3831–3848; (k) W. C. Wertjes, E. H. Southgate and D. Sarlah, *Chem. Soc. Rev.*, 2018, **47**, 7996–8017; (l) G. Huang and B. Yin, *Adv. Synth. Catal.*, 2019, **361**, 405–425; (m) F.-T. Sheng, J.-Y. Wang, W. Tan, Y.-C. Zhang and F. Shi, *Org. Chem. Front.*, 2020, **7**, 3967–3998; (n) M. Okumura and D. Sarlah, *Eur. J. Org. Chem.*, 2020, **2020**, 1259–1273; (o) Y.-Z. Cheng, Z. Feng, X. Zhang and S.-L. You, *Chem. Soc. Rev.*, 2022, **51**, 2145–2170.
- For selected examples, see: (a) A. Taheri, B. Lai, C. Cheng and Y. Gu, *Green Chem.*, 2015, **17**, 812–816; (b) S. P. Pitre, J. C. Scaiano and T. P. Yoon, *ACS Catal.*, 2017, **7**, 6440–6444; (c) K. Wu, Y. Du and T. Wang, *Org. Lett.*, 2017, **19**, 5669–5672; (d) D. Alpers, M. Gallhof, J. Witt, F. Hoffmann and M. Brasholz, *Angew. Chem., Int. Ed.*, 2017, **56**, 1402–1406; (e) D. Chen, L. Xu, T. Long, S. Zhu, J. Yang and L. Chu, *Chem. Sci.*, 2018, **9**, 9012–9017; (f) E. C. Gentry, L. J. Rono, M. E. Hale, R. Matsuura and R. R. Knowles, *J. Am. Chem. Soc.*, 2018, **140**, 3394–3402; (g) F. Strieth-Kalthoff, C. Henkel, M. Teders, A. Kahnt, W. Knolle, A. Gómez-Suárez, K. Dirian, W. Alex, K. Bergander, C. G. Daniliuc, B. Abel, D. M. Guldi and F. Glorius, *Chem.*, 2019, **5**, 2183–2194; (h) J. Yang, D.-W. Ji, Y.-C. Hu, X.-T. Min, X. Zhou and Q.-A. Chen, *Chem. Sci.*, 2019, **10**, 9560–9564; (i) Y.-Z. Cheng, Q.-R. Zhao, X. Zhang and S.-L. You, *Angew. Chem., Int. Ed.*, 2019, **58**, 18069–18074; (j) M. Zhu, C. Zheng, X. Zhang and S.-L. You, *J. Am. Chem. Soc.*, 2019, **141**, 2636–2644; (k) M. Zhu, X.-L. Huang, H. Xu, X. Zhang, C. Zheng and S.-L. You, *CCS Chem.*, 2020, **3**, 652–664; (l) W.-J. Zhou, Z.-H. Wang, L.-L. Liao, Y.-X. Jiang, K.-G. Cao, T. Ju, Y. Li, G.-M. Cao and D.-G. Yu, *Nat. Commun.*, 2020, **11**, 3263; (m) H. Li, L. Guo, X. Feng, L. Huo, S. Zhu and L. Chu, *Chem. Sci.*, 2020, **11**, 4904–4910; (n) J. Ma, F. Schäfers, C. Daniliuc, K. Bergander, C. A. Strassert and F. Glorius, *Angew. Chem., Int. Ed.*, 2020, **59**, 9639–9645; (o) M. S. Oderinde, E. Mao, A. Ramirez, J. Pawluczyk, C. Jorge, L. A. M. Cornelius, J. Kempson, M. Vetrichevan, M. Pitchai, A. Gupta, A. K. Gupta, N. A. Meanwell, A. Mathur and T. G. M. Dhar, *J. Am. Chem. Soc.*, 2020, **142**, 3094–3103; (p) K. Sun, A. Shi, Y. Liu, X. Chen, P. Xiang, X. Wang, L. Qu and B. Yu, *Chem. Sci.*, 2022, **13**, 5659–5666; (q) E. Georgiou, D. Spinnato, K. Chen, P. Melchiorre and K. Muñiz, *Chem. Sci.*, 2022, **13**, 8060–8064; (r) X. Chang, F. Zhang, S. Zhu, Z. Yang, X. Feng and Y. Liu, *Nat. Commun.*, 2023, **14**, 3876; (s) X. Lv, Y.-N. Qi, J. Wang, X. Zhao and Z. Jiang, *Org. Lett.*, 2023, **25**, 3114–3119; (t) T.-T. Song, Y.-K. Mei, Y. Liu, X.-Y. Wang, S.-Y. Guo, D.-W. Ji, B. Wan, W. Yuan and Q.-A. Chen, *Angew. Chem., Int. Ed.*, 2023, e202314304; (u) J. Qin, Z. Zhang, Y. Lu, S. Zhu and L. Chu, *Chem. Sci.*, 2023, **14**, 12143–12151.
- For selected examples, see: (a) G. Tan, Q. You, J. Lan and J. You, *Angew. Chem., Int. Ed.*, 2018, **57**, 6309–6313; (b) Y. Dong, A. W. Schuppe, B. K. Mai, P. Liu and S. L. Buchwald, *J. Am. Chem. Soc.*, 2022, **144**, 5985–5995; (c) X. Li, L. Hu, S. Ma, H. Yu, G. Lu and T. Xu, *ACS Catal.*, 2023, **13**, 4873–4881.
- For selected reviews, see: (a) Y. Yao, M. Alami, A. Hamze and O. Provot, *Org. Biomol. Chem.*, 2021, **19**, 3509–3526; (b) S. M. Umer, M. Solangi, K. M. Khan and R. S. Z. Saleem, *Molecules*, 2022, **27**, 7586; (c) D. P. Zlotos, Y. M. Mandour and A. A. Jensen, *Nat. Prod. Rep.*, 2022, **39**, 1910–1937; for selected examples, see: (d) R. B. Woodward, M. P. Cava, W. D. Ollis, A. Hunger, H. U. Daeniker and K. Schenker, *Tetrahedron*, 1963, **19**, 247–288; (e) L. Randriambola, J.-C. Quirion, C. Kan-Fan and H.-P. Husson, *Tetrahedron Lett.*, 1987, **28**, 2123–2126; (f) P. J. Stephens, J.-J. Pan, F. J. Devlin, M. Urbanová and J. Hájíček, *J. Org. Chem.*, 2007, **72**, 2508–2524; (g) X. Zhang and J. C. Anderson, *Angew. Chem., Int. Ed.*, 2019, **58**, 18040–18045; (h) S. Al-



- Rashed, A. Baker, S. S. Ahmad, A. Syed, A. H. Bahkali, A. M. Elgorban and M. S. Khan, *Bioorg. Chem.*, 2021, **107**, 104626.
- 6 (a) S.-T. Chen and J.-M. Fang, *J. Org. Chem.*, 1997, **62**, 4349–4357; (b) D. Ameena and T. J. Snape, *Med. Chem. Commun.*, 2013, **4**, 893–907.
- 7 For selected reviews, see: (a) J. M. R. Narayanam and C. R. J. Stephenson, *Chem. Soc. Rev.*, 2011, **40**, 102–113; (b) J. Xuan and W.-J. Xiao, *Angew. Chem., Int. Ed.*, 2012, **51**, 6828–6838; (c) C. K. Prier, D. A. Rankic and D. W. C. MacMillan, *Chem. Rev.*, 2013, **113**, 5322–5363; (d) T. P. Nicholls, D. Leonori and A. C. Bissember, *Nat. Prod. Rep.*, 2016, **33**, 1248–1254; (e) M. H. Shaw, J. Twilton and D. W. C. MacMillan, *J. Org. Chem.*, 2016, **81**, 6898–6926; (f) K. L. Skubi, T. R. Blum and T. P. Yoon, *Chem. Rev.*, 2016, **116**, 10035–10074; (g) N. A. Romero and D. A. Nicewicz, *Chem. Rev.*, 2016, **116**, 10075–10166; (h) M. Silvi and P. Melchiorre, *Nature*, 2018, **554**, 41–49; (i) F. Strieth-Kalthoff, M. J. James, M. Teders, L. Pitzer and F. Glorius, *Chem. Soc. Rev.*, 2018, **47**, 7190–7202; (j) L. Marzo, S. K. Pagire, O. Reiser and B. König, *Angew. Chem., Int. Ed.*, 2018, **57**, 10034–10072; (k) A. A. Festa, L. G. Voskressensky and E. V. Van der Eycken, *Chem. Soc. Rev.*, 2019, **48**, 4401–4423; (l) X. Huang and E. Meggers, *Acc. Chem. Res.*, 2019, **52**, 833–847; (m) T.-Y. Shang, L.-H. Lu, Z. Cao, Y. Liu, W.-M. He and B. Yu, *Chem. Commun.*, 2019, **55**, 5408–5419; (n) Q.-Q. Zhou, Y.-Q. Zou, L.-Q. Lu and W.-J. Xiao, *Angew. Chem., Int. Ed.*, 2019, **58**, 1586–1604; (o) Y. Chen, L.-Q. Lu, D.-G. Yu, C.-J. Zhu and W.-J. Xiao, *Sci. China: Chem.*, 2019, **62**, 24–57; (p) A. Hossain, A. Bhattacharyya and O. Reiser, *Science*, 2019, **364**, eaav9713; (q) Z. Zhang, J.-H. Ye, T. Ju, L.-L. Liao, H. Huang, Y.-Y. Gui, W.-J. Zhou and D.-G. Yu, *ACS Catal.*, 2020, **10**, 10871–10885; (r) M. Latrache and N. Hoffmann, *Chem. Soc. Rev.*, 2021, **50**, 7418–7435; (s) J. D. Bell and J. A. Murphy, *Chem. Soc. Rev.*, 2021, **50**, 9540–9685; (t) F.-D. Lu, J. Chen, X. Jiang, J.-R. Chen, L.-Q. Lu and W.-J. Xiao, *Chem. Soc. Rev.*, 2021, **50**, 12808–12827; (u) G.-Q. Xu and P.-F. Xu, *Chem. Commun.*, 2021, **57**, 12914–12935; (v) A. Vega-Peñalozza, J. Mateos, X. Companyó, M. Escudero-Casao and L. Dell'Amico, *Angew. Chem., Int. Ed.*, 2021, **60**, 1082–1097; (w) A. Tlili and S. Lakhdar, *Angew. Chem., Int. Ed.*, 2021, **60**, 19526–19549; (x) Y. Sakakibara and K. Murakami, *ACS Catal.*, 2022, **12**, 1857–1878; (y) T. Bortolato, S. Cuadros, G. Simionato and L. Dell'Amico, *Chem. Commun.*, 2022, **58**, 1263–1283; (z) A. Y. Chan, I. B. Perry, N. B. Bissonnette, B. F. Buksh, G. A. Edwards, L. I. Frye, O. L. Garry, M. N. Lavagnino, B. X. Li, Y. Liang, E. Mao, A. Millet, J. V. Oakley, N. L. Reed, H. A. Sakai, C. P. Seath and D. W. C. MacMillan, *Chem. Rev.*, 2022, **122**, 1485–1542; (aa) K. P. S. Cheung, S. Sarkar and V. Gevorgyan, *Chem. Rev.*, 2022, **122**, 1543–1625; (ab) S. P. Pitre and L. E. Overman, *Chem. Rev.*, 2022, **122**, 1717–1751; (ac) D. A. Corbin and G. M. Miyake, *Chem. Rev.*, 2022, **122**, 1830–1874.
- 8 For selected reviews, see: (a) K. N. Lee and M.-Y. Ngai, *Chem. Commun.*, 2017, **53**, 13093–13112; (b) Q. Xia, J. Dong, H. Song and Q. Wang, *Chem.-Eur. J.*, 2019, **25**, 2949–2961; (c) Á. Péter, S. Agasti, O. Knowles, E. Pye and D. J. Procter, *Chem. Soc. Rev.*, 2021, **50**, 5349–5365; for selected examples, see: ; (d) S. Gross and H.-U. Reissig, *Org. Lett.*, 2003, **5**, 4305–4307; (e) K. T. Tarantino, P. Liu and R. R. Knowles, *J. Am. Chem. Soc.*, 2013, **135**, 10022–10025; (f) Z. Qu, T. Tian, Y. Tan, X. Ji, G.-J. Deng and H. Huang, *Green Chem.*, 2022, **24**, 7403–7409; (g) C. N. Rao and H.-U. Reissig, *Eur. J. Org. Chem.*, 2022, e202200264; (h) V. J. Mayerhofer, M. Lippolis and C. J. Teskey, *Angew. Chem., Int. Ed.*, 2023, **63**, e202314870; (i) W.-Y. Zhang, H.-C. Wang, Y. Wang, C. Zheng and S.-L. You, *J. Am. Chem. Soc.*, 2023, **145**, 10314–10321; (j) M. Wang, C. Zhang, C. Ci, H. Jiang, P. H. Dixneuf and M. Zhang, *J. Am. Chem. Soc.*, 2023, **145**, 10967–10973.
- 9 H. G. Roth, N. A. Romero and D. A. Nicewicz, *Synlett*, 2016, **27**, 714–723.
- 10 For selected examples, see: (a) A. Call, C. Casadevall, F. Acuña-Parés, A. Casitas and J. Lloret-Fillol, *Chem. Sci.*, 2017, **8**, 4739–4749; (b) J. Cai, Y. Li, Z. Ye, W. Wang, Y. Lin and L. Gong, *Green Synth. Catal.*, 2023, **4**, 253–257.
- 11 For selected examples, see: (a) M. Nakajima, E. Fava, S. Loescher, Z. Jiang and M. Rueping, *Angew. Chem., Int. Ed.*, 2015, **54**, 8828–8832; (b) K. Li, Q. Wan, C. Yang, X.-Y. Chang, K.-H. Low and C.-M. Che, *Angew. Chem., Int. Ed.*, 2018, **57**, 14129–14133; (c) A. Caron, É. Morin and S. K. Collins, *ACS Catal.*, 2019, **9**, 9458–9464; (d) N. Noto, Y. Hyodo, M. Yoshizawa, T. Koike and M. Akita, *ACS Catal.*, 2020, **10**, 14283–14289; (e) M. Liu, L. Tan, R. T. Rashid, Y. Cen, S. Cheng, G. Botton, Z. Mi and C.-J. Li, *Chem. Sci.*, 2020, **11**, 7864–7870; (f) T. Luo, L. Li, Y. Chen, J. An, C. Liu, Z. Yan, J. H. Carter, X. Han, A. M. Sheveleva, F. Tuna, E. J. L. McInnes, C. C. Tang, M. Schröder and S. Yang, *Nat. Commun.*, 2021, **12**, 3583; (g) H. Wang, J.-P. Qu and Y.-B. Kang, *Org. Lett.*, 2021, **23**, 2900–2903; (h) F. Lan, C.-S. Liu, C. Zhou, X. Huang, J.-Y. Wu and X. Zhang, *J. Mater. Chem. A*, 2022, **10**, 16578–16584; (i) S. Kundu, L. Roy and M. S. Maji, *Org. Lett.*, 2022, **24**, 9001–9006; (j) Y. Yan, G. Li, J. Ma, C. Wang, J. Xiao and D. Xue, *Green Chem.*, 2023, **25**, 4129–4136.
- 12 For selected reviews, see: (a) R. Ali and S. Alvi, *Tetrahedron*, 2020, **76**, 131345; (b) C. Zhou, B. An, F. Lan and X. Zhang, *Chem. Commun.*, 2023, **59**, 13245–13257.
- 13 (a) C. Ruiz, E. M. García-Frutos, D. A. da Silva Filho, J. T. López Navarrete, M. C. R. Delgado and B. Gómez-Lor, *J. Phys. Chem. C*, 2014, **118**, 5470–5477; (b) T. Deng, W. Yan, X. Liu, G. Hu, W. Xiao, S. Mao, J. Lin, Y. Jiao and Y. Jin, *Org. Lett.*, 2022, **24**, 1502–1506; (c) R. Gao, B. An, C. Zhou and X. Zhang, *Molecules*, 2022, **27**, 8722.
- 14 See the ESI† for more details.

