



Cite this: *Chem. Sci.*, 2024, **15**, 20421

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 22nd July 2024  
Accepted 13th November 2024

DOI: 10.1039/d4sc04871g  
[rsc.li/chemical-science](http://rsc.li/chemical-science)

## 1 Introduction

Thiol redox reactivity is central to many aspects of biology. Cysteine, the unique thiol-containing amino acid, provides proteins and peptides (*i.e.* glutathione) with unique catalytic,<sup>1,2</sup> structural,<sup>3</sup> signaling,<sup>4,5</sup> and antioxidant<sup>6</sup> functions. In many cases, these functions are underpinned by thiol/disulfide exchange reactions, while in others, oxidation of the thiol to a reactive intermediate is believed to be key. Of particular interest are the reactions of cysteine thiols with  $H_2O_2$ , the pervasive 'reactive oxygen species' that abounds the cell – and at elevated levels in pathological contexts. Disulfides are primarily formed in these reactions, but sulfenic acids ( $RSO_2H$ ) and sulfonic acids ( $RSO_3H$ ) can also be observed depending on conditions.<sup>7</sup> All of these products are believed to result from the initial formation of a sulfenic acid ( $RSOH$ ) from substitution of the thiolate on  $H_2O_2$ ; the sulfenic acid then either reacts as an electrophile, undergoing substitution by another equivalent of thiolate to yield disulfide, or as a nucleophile, where it is oxidized by another equivalent of  $H_2O_2$  to yield  $RSO_2H$  and then

$RSO_3H$ . While thiol oxidation to  $RSO_2H$  is reversible in the cell,  $RSO_3H$  formation is irreversible, abolishing the function of the thiol.<sup>8</sup>

The reactivity of  $RSOH$  as both electrophile and nucleophile leads to their rapid self-reaction, making isolation and determination of accurate physicochemical properties and/or reaction kinetics challenging – especially under physiologically-relevant conditions (*i.e.* in aqueous solutions).<sup>9</sup> Although some protein sulfenic acids are stable and can be observed by mass spectrometry,<sup>10</sup> the protein structure can make mechanistic studies and generalizations about  $RSOH$  reactivity difficult.<sup>11,12</sup> To provide insight to their reactivity, several persistent small-molecule  $RSOH$  have been prepared,<sup>13,14</sup> including a sterically-encumbered cysteine sulfenic acid.<sup>15</sup> Unfortunately, the hydrophobicity of the scaffold restricted the study of its chemical properties to largely organic solutions (*i.e.* 20 : 1 THF/ $H_2O$ ), which cannot be buffered for studies of the pH-dependence of the reactivity of thiols and sulfenic acids – a key consideration given that physiologically-relevant thiol chemistry is dominated by the more reactive conjugate base (thiolate), which is present to a substantial extent at pH 7.4 (thiol  $pK_a$ s range from 5.0 to 8.8).<sup>16</sup>

We previously used a triptycene scaffold functionalized with a bridgehead fluorine atom to monitor the reaction of thiol and  $H_2O_2$  in buffered methanol directly using  $^{19}F$  NMR.<sup>13</sup> The

Department of Chemistry and Biomolecular Sciences, University of Ottawa, 10 Marie Curie Pvt, Ottawa, ON K1N6N5, Canada. E-mail: [drattt@uottawa.ca](mailto:drattt@uottawa.ca)

† Electronic supplementary information (ESI) available: Experimental procedures, characterization data, NMR spectra, additional kinetic data, computational data. See DOI: <https://doi.org/10.1039/d4sc04871g>



tryptcene moiety precludes electrophilic reactions of the sulfenic acid because a nucleophile cannot access the  $\sigma_{S-O}^*$  orbital localized on the sulfur atom, unequivocally confirming it as the initial product. This approach not only enabled detailed kinetic measurements and unambiguous characterization of the specific-base catalysis mechanism responsible, but also the successive oxidations to sulfinic and sulfonic acids which follow.<sup>17</sup> Unfortunately, studies of the reaction of thiols and sulfenic acids with stronger oxidants, including other biologically-relevant 'reactive oxygen species' such as hypochlorous acid (HOCl) and singlet oxygen ( $^1O_2$ ), were not possible due to their competitive reactions with the (amine-based) buffers. Similarly to the reactions of thiols with  $H_2O_2$ , reactions of HOCl and  $^1O_2$  have been shown to yield disulfides, but the mechanisms remain unclear due, in part, to the inability to directly observe reaction intermediates. We surmised that substitution of the 9-triptcenethiol scaffold to increase water solubility would enable direct observation of any intermediates formed in the reactions of thiols with HOCl,  $^1O_2$  and/or other high energy oxidants. Moreover, it would permit analogous studies on sulfenic acids and the investigation of other reactions which have been proposed to involve RSOH intermediates. Herein we describe the preparation of such a scaffold, its use to provide insights to the reactions of thiols and sulfenic acids with  $H_2O_2$ , HOCl and  $^1O_2$ , and describe other applications to study sulfenic acid reactivity.

## 2 Results & discussion

### 2.1. Synthesis of thiol 1 and its oxidation products (6, 7, 8)

To improve the water-solubility of the fluorinated 9-triptcenethiol used in our previous work, we incorporated pendant

tetraethylene glycol (TEG) units on the triptycene backbone as in **1**. The synthesis of **1** was carried out in 7 steps from 9-bromoanthracene with an overall yield of 9% (Fig. 1A). Lithium-bromide exchange on 9-bromo-10-fluorotriptycene (**2**)<sup>17</sup> and substitution of the resulting triptycenylium lithium on methyl disulfide afforded methyl sulfide **3** in 92% yield. Selective bromination of **3** was challenging; the Lewis base-catalyzed approach developed by Nishii and Miura<sup>18</sup> was most successful, leading to a mixture of dibrominated products **4a** (*meta*, *meta* to the bridgehead C-S bond) and **4b** (*meta*, *para*). Interestingly, despite its striking similarity, the fluorinated triptycene **3** did not assist the reaction on its own and the TpSMe used by Nishii and Miura<sup>18</sup> had to be added for the reaction to proceed to completion. Indeed, the presence of the fluorine atom also precluded access to the tribrominated triptycene, which was readily obtained under these reaction conditions in its absence. Both isomers of **4** were then subjected to Ullmann-type coupling conditions in the presence of Cu(OAc)<sub>2</sub> using TEG as both nucleophile and solvent, sluggishly affording a corresponding isomeric mixture of **5** in 78% yield. The methyl sulfide was then cleaved using freshly prepared lithium thioethoxide in absolute DMPU at 120 °C to obtain the target thiol **1** in 51% yield. Reverse phase HPLC could resolve **1a** from **1b**, enabling its unequivocal characterization, but most of our experiments were carried out on the mixture as it afforded a second data set in each independent experiment (*vide infra*).

The sulfenic acids (**6**) and sulfinic acids (**7**) derived from **1** were prepared in 46% and 22% yield, respectively, from the direct oxidation of thiols **1** by  $H_2O_2$  (2 eq.) in  $H_2O/THF$  (1 : 1) containing  $K_2HPO_4$  and separated by flash column chromatography. The sulfonic acids (**8**) were prepared similarly by the overnight reaction of **1** with 5 eq. of  $H_2O_2$  (Fig. 1B).




Fig. 1 (A) Preparation of thiols **1**, and (B) corresponding sulfenic acids **6**, sulfinic acids **7** and sulfonic acids **8**.



## 2.2. Reactions of **1** with $\text{H}_2\text{O}_2$

To obtain reaction kinetics for the oxidation of thiols **1** by  $\text{H}_2\text{O}_2$  in water, **1** was treated with an excess of  $\text{H}_2\text{O}_2$  in phosphate-buffered  $\text{D}_2\text{O}/\text{THF}$  (3 : 1, v/v) at pD 7.8 (pH 7.4),<sup>19</sup> and their consumption was monitored by  $^{19}\text{F}$  NMR along with the corresponding appearance of product sulfenic, sulfinic and sulfonic acids (quantified relative to added 2,2,2-trifluoroethanol) as shown in Fig. 2. Since the two regioisomers of **1** and their corresponding sulfur oxyacids have distinct  $^{19}\text{F}$  NMR shifts, two data sets were collected per experiment (Fig. 2B). These data sets were fit to a simple kinetic model for successive bimolecular reactions of **1**, **6** and **7** with  $\text{H}_2\text{O}_2$  (Fig. 2C), from which observed rate constants ( $k_{\text{obs}}$ ) of 0.10, 0.013 and 0.0014  $\text{M}^{-1} \text{s}^{-1}$  were derived for the reactions of thiol, sulfenic acid and sulfonic acid,

respectively. Not only is this the first example of these reactions being followed directly in aqueous solution but, as far as we are aware, these are the first rate constants for RSOH and sulfonic acid oxidation by  $\text{H}_2\text{O}_2$  measured in aqueous solution.<sup>‡</sup>

Previously measured  $k_{\text{obs}}$  for the reactions of the water-insoluble triptycene thiol, sulfenic acid and sulfonic acid with  $\text{H}_2\text{O}_2$  in methanol at  ${}^s\text{pH} = 9.9$  (ca. pH = 7.66 in  $\text{H}_2\text{O}$ )<sup>20</sup> were  $2.5 \times 10^{-3}$ ,  $9.5 \times 10^{-4}$  and  $7.3 \times 10^{-4} \text{ M}^{-1} \text{s}^{-1}$ , respectively. Given that these reactions are specific-base catalyzed (Fig. 2A), we suspected that the origin of the large solvent effects on the reactions of the thiol (40×) and sulfenic acid (13×) is the more favourable pre-equilibrium in aqueous solution relative to methanol. To derive absolute rate constants, the  $\text{pK}_{\text{a}}$ s of thiol **1** ( $8.3 \pm 0.1$ ) and sulfenic acid **6** ( $10.4 \pm 0.1$ ) were measured in




Fig. 2 (A) Reaction scheme for oxidation of thiols **1** by  $\text{H}_2\text{O}_2$ , including  $\text{pK}_{\text{a}}$  values determined for each of **1**, **6** and **7** and absolute rate constants obtained for the reactions of the conjugate bases of each of **1**, **6**, and **7** with  $\text{H}_2\text{O}_2$  in 50 mM phosphate-buffered pD 7.8  $\text{D}_2\text{O}/\text{THF}$  (3 : 1). The rate constants are derived from the mean of the individual rate constants determined for each of the **a** and **b** isomers (see ESI†). (B) Representative time-resolved  $^{19}\text{F}$  NMR spectra obtained upon treatment of **1** (5 mM) with 25 mM  $\text{H}_2\text{O}_2$  in 50 mM phosphate-buffered pD 7.8  $\text{D}_2\text{O}/\text{THF}$  (3 : 1). (C) Representative reaction profiles of **1a** (2.8 mM) and **1b** (2.2 mM) with 5 eq.  $\text{H}_2\text{O}_2$  in pD 7.8  $\text{D}_2\text{O}/\text{THF}$  (3 : 1), fits of the data to a simple kinetic model for successive bimolecular reactions of **1**, **6** and **7** with  $\text{H}_2\text{O}_2$  and the corresponding observed rate constants ( $\text{M}^{-1} \text{s}^{-1}$ ) derived from the fits.



$\text{D}_2\text{O}/\text{THF}$  (3 : 1) (the values for **1a** and **6a** were within the margin of error on the measurements made with the mixtures and corresponding values in  $\text{H}_2\text{O}$  afforded  $8.0 \pm 0.1$  and  $10.1 \pm 0.1$ , respectively). Expectedly, deprotonation of thiol and sulfenic acid were found to be much more favourable in water than in methanol ( $\text{p}K_a = 11.6$  and  $12.8$ , respectively).<sup>17</sup> With these  $\text{p}K_a$  values at hand, we derived absolute (pH-independent) rate constants for the reactions of thiolate ( $k = 0.43 \text{ M}^{-1} \text{ s}^{-1}$ , compared to  $0.13 \text{ M}^{-1} \text{ s}^{-1}$  in methanol) and sulfenate ( $k = 5.2 \text{ M}^{-1} \text{ s}^{-1}$ , compared to  $0.75 \text{ M}^{-1} \text{ s}^{-1}$  in methanol) with  $\text{H}_2\text{O}_2$ . As expected from the alpha effect<sup>21</sup> and our previous data,<sup>17</sup> sulfenate anions were found to be more reactive nucleophiles than thiolate anions. Sulfinate was found to be much less reactive as a nucleophile ( $k = 1.4 \times 10^{-3} \text{ M}^{-1} \text{ s}^{-1}$  compared to  $5.8 \times 10^{-4} \text{ M}^{-1} \text{ s}^{-1}$  in methanol) consistent with the greater relative stability of the anion ( $\text{p}K_a \sim 2^{17}$  vs. 10 for the corresponding RSOH). Overall, the rate constant determined for **1** is toward the low end of the range of those which have previously been reported (derived from consumption of thiol(ate)s),<sup>8</sup> which span from  $0.16 \text{ M}^{-1} \text{ s}^{-1}$  (for *N*-acetylcysteine) to  $4.5 \text{ M}^{-1} \text{ s}^{-1}$  (for penicillamine) at pH 7.4,<sup>12</sup> which is presumably due to the electron-withdrawing effect of the 10-fluorotriptyceny scaffold.

### 2.3. Reactions of **1** with HOCl

At physiological pH, thiols react rapidly with HOCl ( $k > 10^8 \text{ M}^{-1} \text{ s}^{-1}$ ),<sup>22</sup> with sulfenyl chlorides being proposed as the initial products, but which are not observed due to their rapid reaction with thiols to yield disulfides.<sup>23</sup> We observed instant formation of a single product (**13a**) upon addition of limiting NaOCl to a solution of **1a** in pD 7.8  $\text{D}_2\text{O}/\text{THF}$  (Fig. 3A). The  $^{19}\text{F}$  NMR chemical shift ( $-205.9$  ppm) did not correspond to the sulfenic acid **6a** ( $-206.6$  ppm), implying formation of the sulfenyl chloride. However, when thiol **1a** was treated with *N*-chlorosuccinimide in chloroform, a product (**14a**) with a distinct  $^{19}\text{F}$  NMR chemical shift ( $-206.4$  ppm) was obtained (Fig. 3B).<sup>24</sup> High-resolution mass spectrometry indicated that **14a** was the sulfenyl chloride. While **14a** was stable over extended periods in pD 7.8  $\text{D}_2\text{O}/\text{THF}$  buffer, addition of **1a** readily yielded **13a**, which we have characterized as the disulfide of **1a** (Fig. 3C). Since substitution of thiolate **1a** on sulfenyl chloride **14a** to form the disulfide is unlikely (the nucleophile cannot access the  $\sigma_{\text{S}-\text{Cl}}^*$  orbital localized on the sulfur atom), the thiolate presumably reacts with the sulfenyl chloride by electron transfer, resulting in the formation of thiyl radicals which dimerize to give the disulfide (Fig. 3B). An analogous mechanism has been proposed for the observation of disulfides from the reaction of sulfenyl chlorides and amines.<sup>25</sup> The intervention of thiyl radicals in the reaction of non-hindered thiols with HOCl has previously been demonstrated by Davies,<sup>26</sup> who characterized DMPO adducts of thiyl radicals by EPR, suggesting that this pathway is not unique to the reaction of 9-tritycenethiols.

### 2.4. Reactions of **1** with $^1\text{O}_2$

Similarly to their reactions with  $\text{H}_2\text{O}_2$  and HOCl, thiols have been shown to yield disulfides upon reaction with  $^1\text{O}_2$ .<sup>27</sup> To

provide insight to the mechanism of this reaction, we investigated the reaction of **1** with  $^1\text{O}_2$  derived from decomposition of the endoperoxide of sodium naphthalene-1,4-dipropionate (NDPO<sub>2</sub>) at 37 °C ( $t_{1/2} = 23$  min).<sup>28,29</sup> The reactions were again monitored by  $^{19}\text{F}$  NMR at 37 °C in pD 7.8 phosphate buffered  $\text{D}_2\text{O}/\text{THF}$  (3 : 1, v/v), but now in the presence of 4 eq. of NDPO<sub>2</sub> in lieu of either  $\text{H}_2\text{O}_2$  or HOCl. After 33 minutes, the conversion of thiol was  $\sim 45\%$ , giving primarily sulfinate **7** ( $\sim 25\%$ ). Importantly, disulfide **13a** was not observed. Other products included two presumed pairs of unknown isomers ( $\sim 10\%$ ) downfield of **8** at  $\delta = -204.55$  ppm and  $-205.25$  ppm (Fig. 4A). Control experiments suggest that the formation of these unknowns can be attributed to a reaction occurring in THF in the presence of  $^1\text{O}_2$  (see ESI Fig. S6†). The missing  $\sim 10\%$  in the mass balance suggests that some signal suppression from the formation of persistent paramagnetic species is taking place (see ESI Fig. S3†).<sup>30,31</sup>

Consistent with the foregoing experiments with **1**, when thiol **9** was subjected to Rose Bengal-photosensitized oxygenation in 15 : 1  $\text{CD}_3\text{OD} : \text{H}_2\text{O}$  containing 3 eq. NaOH, the corresponding sulfinate **11** and sulfonate **12** were formed (Fig. 4B) (see Fig. S4 of the ESI† for corresponding  $^1\text{H}$  NMR spectra). Traces of disulfide **15** could be detected under these conditions (as a precipitate), suggesting that thiyl radical formation is possible under the reaction conditions, albeit as a minor reaction pathway.¶ Likewise, while RSOH **6** was not detected in  $\text{D}_2\text{O}/\text{THF}$ , a minuscule amount of **10** could be detected in  $\text{CD}_3\text{OD}/\text{D}_2\text{O}$  (Fig. 4B).

The drastically different product distribution arising from the reaction of **1** (and **9**) with  $^1\text{O}_2$  compared to the precedent<sup>32</sup> suggests that nucleophilic substitution on an oxidized sulfur intermediate is responsible for disulfide formation from reactions of thiols and  $^1\text{O}_2$ , in general. While at first glance it would seem that RSOH is that intermediate, and although we found that RSOH **10** was readily photooxidized to RSO<sub>2</sub>H **11** (see ESI Fig. S7†), it is difficult to provide a compelling mechanistic rationale for the direct formation of RSOH from RSH and  $^1\text{O}_2$ . Instead, we posit that a sulfenyl hydroperoxide (RSOOH) is the immediate product, and when the reaction with a thiolate cannot take place at sulfur to give a disulfide (*i.e.* for **1** and **9**), the reaction takes place at oxygen to give two equivalents of RSOH. Alternatively, RSOOH can rearrange to RSO<sub>2</sub>H. If formed, RSOH can react with  $^1\text{O}_2$  to afford a sulfinyl hydroperoxide (RS(O)OOH) which can oxidize another equivalent of RSOH to yield two equivalents of RSO<sub>2</sub>H (Fig. 4C).

Formation of RSOO<sup>−</sup> from thiolate (RS<sup>−</sup>) and  $^1\text{O}_2$  is predicted by CBS-QB3 (ref. 33) to be highly exergonic ( $\Delta G^\circ = -25.1 \text{ kcal mol}^{-1}$  for *t*-BuS<sup>−</sup> as a model thiolate). No TS could be computed for the nucleophilic addition pathway, presumably due to the use of a single determinant wavefunction approach. A TS could, however, be computed for the addition of RSH to  $^1\text{O}_2$ , which was associated with a barrier of  $\Delta G^\ddagger = 14.7 \text{ kcal mol}^{-1}$  (Fig. 4D). This may be taken as an upper bound for the thiolate reaction since the barrier to addition of the better nucleophile is sure to be lower. Another possibility involves electron transfer from RS<sup>−</sup> to  $^1\text{O}_2$  to form a radical pair that can combine or, if cage escape occurs, oxygenation of the thiyl radical to yield






Fig. 3 (A) Representative  $^{19}\text{F}$  NMR spectra obtained before and after treatment of **1a** (17 mM) with 0.35 eq. of NaOCl in 50 mM phosphate-buffered pD 7.8  $\text{D}_2\text{O}/\text{THF}$  (3 : 1). (B) Preparation of sulfenyl chloride **14a** from **1a** and representative  $^{19}\text{F}$  NMR in  $\text{CDCl}_3$ . (C) Representative  $^{19}\text{F}$  NMR spectra obtained before and after treatment of **14a** (17 mM) with 0.65 eq. of **1a** in 50 mM phosphate-buffered pD 7.8  $\text{D}_2\text{O}/\text{THF}$  (3 : 1). (D) Proposed mechanism of disulfide formation upon thiol oxidation by hypochlorous acid.

a thiylperoxy radical ( $\text{RSOO}^\cdot$ ) that could be subsequently reduced *via* either H-atom transfer or electron transfer. H-atom transfer from a thiol to  $\text{RSOO}^\cdot$  is predicted to be exergonic ( $\Delta G^\circ = -4.1 \text{ kcal mol}^{-1}$ ) and proceed *via* an accessible barrier ( $\Delta G^\ddagger = 18.8 \text{ kcal mol}^{-1}$ ). However, our calculations indicate that the highly exergonic ( $\Delta G^\circ = -76.8 \text{ kcal mol}^{-1}$ ) rearrangement of  $\text{RSOO}^\cdot$  to a sulfonyl radical ( $\text{RSO}_2^\cdot$ ) is much more likely. Earlier calculations had suggested this would require traversing a prohibitively high barrier ( $\Delta G^\ddagger = 45 \text{ kcal mol}^{-1}$ ),<sup>34</sup> but we have found this barrier to be significantly lower ( $\Delta G^\ddagger = 13.7 \text{ kcal mol}^{-1}$ ) using the higher accuracy CBS-QB3 methodology. Regardless, the lack of disulfide formed in our experiments argues that thiyl radicals are not key intermediates.

The sulfinate observed when disulfide formation is suppressed could also arise from concerted rearrangement of  $\text{RSOOH}$  – a highly exergonic reaction ( $\Delta G^\circ = -60.6 \text{ kcal mol}^{-1}$ ) with an easily accessible barrier ( $\Delta G^\ddagger = 13.6 \text{ kcal mol}^{-1}$ ). Again, this barrier is likely an upper bound given that the H-bond between the migrating OH group and an explicit solvent molecule will become stronger in the TS (this cannot be captured in the polarizable solvent continuum employed in the calculation). Alternatively, a step-wise rearrangement can be envisioned. The calculated O–O BDE of  $\text{RSOOH}$  was found to be a meagre

10.2 kcal mol<sup>-1</sup>, driven by the stability of the sulfinyl radical, suggesting O–O bond homolysis followed by in-cage recombination to form a sulfinic acid is possible. Earlier spin-trapping studies during thiol photooxygenation suggested the formation of  $^\cdot\text{OH}$ , presumably due to cage-escape following  $\text{RSOOH}$  thermolysis (Fig. 4E).<sup>30</sup> The lack of significant signal suppression in our NMR experiments, suggests that homolysis does not occur to a significant extent and that nucleophilic chemistry predominates. Regardless, the predicted rapid rearrangement chemistry presumably prevents the direct observation of  $\text{RSOOH}$  on the timescale of the NMR experiments.

The implication of  $\text{RSOOH}$  in disulfide formation from thiols exposed to  $^1\text{O}_2$  compels consideration that  $\text{RSOOH}$  could also be intermediates in disulfide formation in the radical-mediated oxidation (autoxidation) of thiols, given the rapid reaction of thiyl radicals with  $^3\text{O}_2$  ( $\sim 10^8 \text{ M}^{-1} \text{ s}^{-1}$ ).<sup>35</sup> Indeed, sulfur oxyacids are major products under some conditions,<sup>36,37</sup> and may derive from  $\text{RSOOH}$  as a common intermediate to disulfide formation. However, as indicated above, given that the (unimolecular) rearrangement of  $\text{RSOO}^\cdot$  to  $\text{RSO}_2^\cdot$  is predicted to proceed with a smaller barrier than the (biomolecular) H-atom transfer to  $\text{RSOO}^\cdot$  from  $\text{RSH}$  (13.7 kcal mol<sup>-1</sup> vs. 18.8 kcal mol<sup>-1</sup>, respectively), the formation of  $\text{RSOOH}$  under




Fig. 4 (A) Representative time-resolved <sup>19</sup>F NMR spectra obtained upon treatment of **1** (5 mM) with 4 eq. of NDPO<sub>2</sub> in 50 mM phosphate-buffered pD 7.8 D<sub>2</sub>O/THF (3:1). (B) Representative time-resolved <sup>19</sup>F NMR spectra obtained of the reaction of **9** (20 mM) with photochemically generated <sup>1</sup>O<sub>2</sub> in CD<sub>3</sub>OD/H<sub>2</sub>O (15:1) containing 60 mM NaOH. (C) Scheme accounting for the different product distributions arising from reaction of hindered and unhindered thiols with <sup>1</sup>O<sub>2</sub>. (D) Mechanistic possibilities and CBS-QB3 calculated free energies (298 K) of the reaction of a model (t-butyl) thiol with <sup>1</sup>O<sub>2</sub> (SET = single electron transfer; HAT = hydrogen atom transfer). Data calculated using a polarizable continuum model using the integral equation formalism (IEFPCM) parameterized for water are given; associated gas phase values are given in the ESI.†

radical conditions is unlikely. Instead, the partitioning of RS<sup>·</sup> and RSOO<sup>·</sup> between thiyl dimerization and thiylperoxyl rearrangement pathways is likely key in determining whether disulfides or sulfur oxyacids are the major products.

## 2.5. Other applications of the new scaffold

Since comparably very little remains known of RSOH reactivity (relative to RSH, RSO<sub>2</sub>H and RSO<sub>3</sub>H), we surmised that **6** may prove generally useful as a mechanistic probe. It has been shown previously that ascorbate acts as the cofactor for 1-Cys peroxiredoxins (Prx), and it was proposed to reduce the RSOH of 1-Cys Prx back to its active thiol form after oxidation.<sup>38</sup> However, other protein sulfenic acids are resistant to reduction by ascorbate.<sup>39</sup> Previously employed indirect bifunctional assays estimated rate constants ranging from 2.2 to 4 × 10<sup>2</sup> M<sup>-1</sup> s<sup>-1</sup>, depending on the protein RSOH. To provide additional insight

to this reactivity, we incubated **6** with 3 eq. of ascorbate<sup>||</sup> in pD 7.8 D<sub>2</sub>O/THF (3:1) and monitored the reaction by <sup>19</sup>F NMR (Fig. 5A). No reaction was observed over 72 hours, suggesting an amino acid sidechain in the Prx active site is required for activation of the sulfenic acid to reduction (the conjugate acid of an essential histidine residue is a possibility), or that a more sophisticated redox cascade is at play.

Labile iron(II) is often invoked as key to the auto-initiation of cellular (phospho)lipid peroxidation and associated ferroptosis due to its ability to reduce lipid hydroperoxides to form initiating lipid alkoxyl radicals.<sup>40,41</sup> Since thiyl radicals are significantly more thermodynamically stable than alkoxyl radicals (*i.e.* t-BuSH and t-BuOH BDEs are 86 and 106 kcal mol<sup>-1</sup>, respectively),<sup>42</sup> we anticipated that RSOH would readily undergo the same reaction (Fig. 5B). To our great surprise, **10** did not react with Fe(II) (or Fe(III) in the presence of ascorbic acid) in methanol. At first glance, we surmised that this may have to do with




Fig. 5 (A) 1-Cys peroxiredoxins (Prx) reduce hydrogen peroxide and hydroperoxides by nucleophilic substitution by the cysteine thiol(ate). The resultant sulfenic acids are converted back to the starting thiol by ascorbate to complete the catalytic cycle. The lack of reactivity of RSOH 6 to ascorbate implies that active site residues are essential to enzyme turnover. (B) Lipid peroxidation is auto-initiated in the presence of  $\text{Fe}^{2+}$  due to reductive cleavage of the O–O bonds in product hydroperoxides, which yields initiating radicals. No reaction is observed for RSOH 10, implying that the process is less favourable for RSOH than it is for the corresponding ROOH.

some impairment of the reaction by the triptycene backbone. However, CBS-QB3 calculations predict that the S–O bond in a model (t-butyl) RSOH is 27 kcal mol<sup>-1</sup> stronger than the O–O bond in the corresponding hydroperoxide – despite the 20 kcal mol<sup>-1</sup> difference in the stabilities of the product thyl and alkoxy radicals. The significant stabilization of RSOH relative to ROOH can be rationalized by lesser lone pair/lone pair repulsion in RSOH due to the longer S–O bond and more diffuse S lone pairs and also greater lone-pair/ $\sigma_{\text{O}-\text{H}}^*$  hyperconjugation in RSOH compared to ROOH. Based on these calculated thermodynamics and the lack of reactivity of 6 and ascorbate with or without added iron, it is clear that reductive heterolysis of RSOH is a highly unfavourable process in instances where there is no acid available to activate the RSOH.

### 3 Conclusions

A newly synthesized 10-fluorotriptycene thiol enables visualization of transient reactive intermediates (e.g., RSOH and RS<sup>+</sup>)

and can be used to directly observe transformations of thiols and sulfenic acids in aqueous buffer by <sup>19</sup>F NMR. Reactions of thiols and sulfenic acids with  $\text{H}_2\text{O}_2$  were found to proceed with kinetics that were slightly faster than in methanol, largely due to more favourable ionization to the reactive conjugate bases. Reaction of thiol with HOCl did not yield a sulfenic acid, but presumably a sulfenyl chloride. The sulfenyl chloride could be independently prepared by reaction of the thiol with N-chlorosuccinimide and was found to rapidly yield disulfide when treated with thiol(ate) in buffer – attributed to dimerization of thyl radicals formed upon electron transfer from thiolate to sulfenyl chloride. Product distributions of the reaction of thiols with <sup>1</sup>O<sub>2</sub> enable us to propose sulfenyl and sulfinyl hydroperoxides as intermediates en route to disulfides, sulfenic and sulfonic acids. Through product analyses and CBS-QB3 calculated reaction energetics, the myriad of one- and two-electron processes that these short-lived intermediates undergo are herein summarized. A lack of reactivity of RSOH 6 and 10 to one-electron reduction strongly supports that reductive heterolysis of RSOH does not occur to a significant extent without enzyme (acid) catalysis and highlights the energetic contribution of lone pair repulsion in hydroperoxides to their relative instability. We anticipate that this scaffold will be a useful tool compound for mechanistic investigations of thiol and sulfenic acid reactivity.

## 4 Experimental section

### 4.1. General

Reagents and solvents were obtained from commercial suppliers and used as received, unless indicated otherwise. Hydrogen peroxide concentration was determined by manometric titration. Sodium hypochlorite concentration was determined by iodometry. <sup>1</sup>H, <sup>13</sup>C and <sup>19</sup>F NMR spectra were recorded on a Bruker AVANCE spectrometer at the specified frequency. <sup>19</sup>F NMR spectra in  $\text{D}_2\text{O}$  were referenced using 2,2,2-trifluoroethanol (TFE) ( $\delta$  –77.0 ppm) as an internal standard, <sup>19</sup>F NMR spectra in  $\text{CDCl}_3$ ,  $\text{CD}_3\text{OD}$ , and  $\text{C}_6\text{D}_6$  were referenced to trifluorotoluene ( $\delta$  –63.72 ppm). Product ratios were obtained through integration of their <sup>19</sup>F NMR signals. For quantification, <sup>19</sup>F NMR spectra were phased and baseline corrected. For NMR characterization of isomeric mixtures, the signals corresponding to each isomer were assigned based on those observed for HPLC-purified isomer 1a, and the corresponding signals observed when it was oxidized with  $\text{H}_2\text{O}_2$ . Preparative HPLC was carried on a Waters 2996 HPLC system, using a preparative Waters XBridge Prep C18 5  $\mu\text{m}$  (19  $\times$  150 mm) column. Column chromatography was carried out with 40–63  $\mu\text{m}$ , 230–400 mesh silica gel purchased from SiliCycle. All calculations were carried out using the Gaussian 09 quantum chemistry package using the CBS-QB3 complete basis set approach.<sup>33</sup>

### 4.2. Synthesis

**4.2.1 10-Fluoro-9,10-[1',2']benzenoanthracen-9(10H)-yl(methyl)sulfane (3).** Warning: a significant amount of



methanethiol is produced in this reaction. To an oven-dried flask charged with a solution of 9-bromo-10-fluorotriptycene<sup>17</sup> (3.0 g, 8.57 mmol) in neat THF (40.0 mL) and cooled to  $-78^{\circ}\text{C}$ , was slowly added *n*BuLi (4.12 mL, 10.28 mmol) as a 2.5 M solution in hexanes. The reaction was stirred at  $-78^{\circ}\text{C}$  for 30 minutes, after which methyl disulfide (0.91 mL, 10.28 mmol) was added and the suspension was left to warm up to room temperature, upon which it became a homogeneous solution. After 1 h, the reaction was quenched with a saturated solution of NH<sub>4</sub>Cl. The crude product was extracted using EtOAc (3  $\times$  50 mL), dried over MgSO<sub>4</sub> and concentrated *in vacuo*. The residue was then purified by column chromatography using hexanes as the eluent to yield **3** (2.51 g, 92%) as a colourless solid: <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.65–7.59 (m, 6H),  $\delta$  7.17–7.09 (m, 6H),  $\delta$  2.58 (s, 3H). <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  144.2, 144.0, 141.7, 141.6, 125.7, 125.6, 122.5, 118.7, 97.6, 60.5, 14.1. HRMS (EI) *m/z* calculated for [M] C<sub>21</sub>H<sub>15</sub>FS 318.0879, found 318.0866.

**4.2.2 Dibromo-10-fluoro-9,10-[1',2']benzenoanthracen-9(10H)-yl(methyl)sulfane (4).** An oven-dried Schlenk tube was charged with **3** (1.74 g, 5.47 mmol), TpsMe<sup>18</sup> (82 mg, 0.274 mmol) and DCE (25 mL). Under a flow of nitrogen, *N*-bromo-succinimide (3.51 g, 19.69 mmol) and AgSbF<sub>6</sub> (93 mg, 0.274 mmol) were both added as solids. The reaction was stirred for 48 h at  $50^{\circ}\text{C}$  before being poured into a saturated NaHCO<sub>3</sub> solution (50 mL). The product was extracted using CHCl<sub>3</sub> (3  $\times$  30 mL), concentrated *in vacuo*, and dried over MgSO<sub>4</sub>. The residue was purified by column chromatography using hexanes as the eluent to give a mixture of dibrominated isomers as an off-white foam (1.58 g, 61% total yield). The monobrominated derivative was also isolated.

**4.2.2.1 2,7-Dibromo-10-fluoro-9,10-[1',2']benzenoanthracen-9(10H)-yl(methyl)sulfane (4a).** 33% yield. <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  7.74–7.71 (m, 2H),  $\delta$  7.61–7.57 (m, 2H),  $\delta$  7.47–7.43 (m, 2H),  $\delta$  7.31–7.24 (m, 2H),  $\delta$  7.19–7.13 (m, 2H),  $\delta$  2.56 (s, 3H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  145.9, 143.6, 143.2, 142.9, 142.8, 142.7, 140.7, 140.5, 128.9, 126.2, 126.1, 122.8, 122.4, 120.5, 119.0, 97.8, 59.9, 14.1. HRMS (EI) *m/z* calculated for [M] C<sub>21</sub>H<sub>13</sub>Br<sub>2</sub>FS 473.9087, found 473.9089.

**4.2.2.2 2,6-Dibromo-10-fluoro-9,10-[1',2']benzenoanthracen-9(10H)-yl(methyl)sulfane (4b).** 28% yield. <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  7.74–7.71 (m, 2H),  $\delta$  7.61–7.57 (m, 2H),  $\delta$  7.47–7.43 (m, 2H),  $\delta$  7.31–7.24 (m, 2H),  $\delta$  7.19–7.13 (m, 2H),  $\delta$  2.54 (s, 3H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  145.7, 143.4, 143.1, 142.6, 140.7, 140.5, 128.9, 126.2, 126.1, 124.4, 122.8, 122.4, 120.5, 119.9, 118.9, 97.0, 59.9, 14.0. HRMS (EI) *m/z* calculated for [M] C<sub>21</sub>H<sub>13</sub>Br<sub>2</sub>FS 473.9087, found 473.9089.

**4.2.3 Di(tetraethylene glycol)-10-fluoro-9,10-[1',2']benzenoanthracen-9(10H)-yl(methyl)sulfane (5).** To an oven-dried Schlenk tube was added a mixture of the isomers **4** (1.10 g, 2.32 mmol) suspended in tetraethylene glycol (20.0 mL). Copper(II) acetate hydrate (0.91 g, 4.58 mmol) and potassium carbonate (3.20 g, 23.15 mmol) were then both added as solids. The tube was sealed, degassed, and refilled with nitrogen. The blue solution was heated at  $150^{\circ}\text{C}$  for 5 days, during which the reaction became brown. Once the reaction mixture was cooled to ambient temperature, it was poured into distilled water (100 mL) and extracted (3  $\times$  25 mL) with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer

was washed with water (3  $\times$  100 mL) to remove excess TEG, dried over MgSO<sub>4</sub>, and concentrated *in vacuo* to obtain the crude product as a dark-orange oil, which was then purified by column chromatography (gradient 1–3% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to obtain the mixture of isomers **5** (**5a** : **5b** = 57 : 43) as a viscous yellow oil (1.27 g, 1.81 mmol, 78% total yield).

**4.2.3.1 2,7-Di(tetraethylene glycol)-10-fluoro-9,10-[1',2']benzenoanthracen-9(10H)-yl(methyl)sulfane (5a).** <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  7.58–7.52 (m, 2H), 7.46–7.40 (m, 2H), 7.43 (d, *J* = 8.2 Hz, 2H), 7.23–7.16 (m, 2H), 7.14–7.06 (m, 2H), 6.64–6.57 (m, 2H), 4.08 (t, *J* = 4.8 Hz, 4H), 3.80 (t, *J* = 4.8 Hz, 4H), 3.73–3.61 (m, 20H), 3.57 (t, *J* = 4.5 Hz, 4H), 2.56 (s, 3H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  156.9, 143.3, 141.5, 137.0, 136.1, 125.7, 125.5, 122.5, 122.2, 119.3, 118.2, 111.3, 110.8, 110.2, 106.2, 97.0, 72.5, 70.8, 70.6, 70.3, 69.7, 67.8, 61.7, 60.6, 60.0, 14.2. HRMS (ESI<sup>+</sup>) *m/z* calculated for [M] + Na<sup>+</sup> C<sub>36</sub>H<sub>45</sub>FNaO<sub>10</sub>S 725.2772, found 725.2767.

**4.2.3.2 2,6-Di(tetraethylene glycol)-10-fluoro-9,10-[1',2']benzenoanthracen-9(10H)-yl(methyl)sulfane (5b).** <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  7.58–7.52 (m, 2H), 7.46–7.40 (m, 2H), 7.43 (d, *J* = 8.2 Hz, 2H), 7.23–7.16 (m, 2H), 7.14–7.06 (m, 2H), 6.64–6.57 (m, 2H), 4.08 (t, *J* = 4.8 Hz, 4H), 3.80 (t, *J* = 4.8 Hz, 4H), 3.73–3.61 (m, 20H), 3.57 (t, *J* = 4.5 Hz, 4H), 2.54 (s, 3H). <sup>13</sup>C NMR (151 MHz, CDCl<sub>3</sub>)  $\delta$  157.0, 156.9, 143.3, 137.0, 125.7, 125.5, 123.6, 122.5, 122.2, 119.5, 119.3, 118.3, 111.3, 110.8, 110.0, 97.0, 72.5, 70.8, 70.6, 70.3, 69.7, 67.8, 61.7, 60.6, 60.0, 14.2. HRMS (ESI<sup>+</sup>) *m/z* calculated for [M] + Na<sup>+</sup> C<sub>36</sub>H<sub>45</sub>FNaO<sub>10</sub>S 725.2772, found 725.2767.

**4.2.4 Di(tetraethylene glycol)-10-fluoro-9,10-[1',2']benzenoanthracen-9(10H)-ylmercaptan (1).** In a sealable tube, methyl sulfide **5** (349 mg, 0.497 mmol) was dissolved in anhydrous DMPU (2.0 mL), and the solution was purged with N<sub>2</sub> for 2 minutes before freshly prepared lithium ethylmercaptide<sup>43</sup> (169 mg, 2.485 mmol) was quickly added and the tube sealed. The reaction was heated at  $120^{\circ}\text{C}$  for 2 days. Once cooled to room temperature, it was poured into 0.1 M aqueous HCl (100 mL) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3  $\times$  20 mL). The organic fraction was then washed with 0.5 M NaOH (3  $\times$  75 mL), the aqueous washings were collected and acidified to pH  $\sim$ 2 using 12 M HCl, and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3  $\times$  50 mL), dried over MgSO<sub>4</sub>, and concentrated *in vacuo* to give the crude product as a red oil that was purified by column chromatography (gradient elution, 2–4% MeOH/CHCl<sub>3</sub>) to give the mixture of thiols **1** (**1a** : **1b** = 57 : 43) as a viscous pale-yellow oil (174 mg, 51% total yield). The **1a** isomer could be separated by reverse phase HPLC using 3 : 1 MeOH : H<sub>2</sub>O as the mobile phase.

**4.2.4.1 2,7-Di(tetraethylene glycol)-10-fluoro-9,10-[1',2']benzenoanthracen-9(10H)-ylmercaptan (1a).**  $\delta$  156.9 (s), 144.6 (d, *J* = 6.0 Hz), 143.8 (d, *J* = 20 Hz), 136.2 (d, *J* = 20 Hz), 126.1 (s), 125.5 (s), 121.4 (d, *J* = 2 Hz), 119.1 (d, *J* = 6.0 Hz), 118.0 (d, *J* = 6 Hz), 110.4 (s), 110.2 (s), 97.5 (d, *J* = 205 Hz), 72.5 (s), 70.8 (s), 70.6 (s), 70.3 (s), 69.7 (s), 67.8 (s), 61.7 (s), 57.5 (d, *J* = 2 Hz). <sup>19</sup>F NMR (471 MHz, D<sub>2</sub>O/THF)  $\delta$  –207.1. HRMS (ESI<sup>+</sup>) *m/z* calculated for [M] + Na<sup>+</sup> C<sub>36</sub>H<sub>45</sub>FNaO<sub>10</sub>S 711.2632, found 711.2615.

**4.2.4.2 2,6-Di(tetraethylene glycol)-10-fluoro-9,10-[1',2']benzenoanthracen-9(10H)-ylmercaptan (1b).** <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  7.63 (m, 1H), 7.57–7.49 (m, 2H), 7.43 (m, 1H), 7.31 (m,



1H), 7.18 (d,  $J$  = 2.4 Hz, 1H), 7.17–7.04 (m, 2H), 6.62 (dd,  $J$  = 8.4, 2.4 Hz, 1H), 6.58 (dd,  $J$  = 8.4, 2.4 Hz, 1H), 4.09 (t,  $J$  = 4.8 Hz, 4H), 3.80 (t,  $J$  = 4.8 Hz, 4H), 3.73–3.61 (m, 20H), 3.57 (t,  $J$  = 4.5 Hz, 4H), 2.42 (s, 1H).  $^{13}\text{C}$  NMR (151 MHz,  $\text{CDCl}_3$ )  $\delta$  157.1 (s), 157.0 (s), 145.0 (d,  $J$  = 20 Hz), 143.5 (d,  $J$  = 20 Hz), 143.2 (d,  $J$  = 6 Hz), 135.4 (d,  $J$  = 20 Hz), 125.8 (s), 125.7 (s), 122.5 (d,  $J$  = 2 Hz), 121.1 (d,  $J$  = 2 Hz), 119.3 (d,  $J$  = 2 Hz), 118.2 (d,  $J$  = 2 Hz), 110.6 (s), 110.5 (s), 110.3 (s), 110.0 (d,  $J$  = 2 Hz), 97.5 (d,  $J$  = 205 Hz), 72.5 (s), 70.8 (s), 70.6 (s), 70.3 (s), 69.7 (s), 67.8 (s), 61.7 (s), 57.5 (d,  $J$  = 2 Hz).  $^{19}\text{F}$  NMR (471 MHz,  $\text{D}_2\text{O}/\text{THF}$ )  $\delta$  –206.3. HRMS (ESI $+$ )  $m/z$  calculated for  $[\text{M} + \text{Na}]^+$   $\text{C}_{36}\text{H}_{45}\text{FNaO}_{10}\text{S}$  711.2632, found 711.2615.

**4.2.5 Di(tetraethylene glycol)-10-fluoro-9,10-[1',2']benzenoanthracen-9(10H)-ylsulfenic acid (6a) and potassium di(tetraethylene glycol)-10-fluoro-9,10-[1',2']benzenoanthracen-9(10H)-ylsulfinate (7).** Thiols **1** (60 mg, 0.087 mmol, as the isomeric mixture of **1a** : **1b** 57 : 43) were dissolved in THF (1 mL) and added to a solution of dibasic potassium phosphate (30.4 mg, 0.194 mmol) in HPLC-grade  $\text{H}_2\text{O}$  (900  $\mu\text{L}$ ). The mixture was thoroughly mixed to ensure homogeneity before 100  $\mu\text{L}$  of  $\text{H}_2\text{O}_2$  as a 1.94 M solution in  $\text{D}_2\text{O}/\text{H}_2\text{O}$  was slowly added while stirring. Progress of the reaction was monitored by  $^{19}\text{F}$  NMR. Upon complete consumption of the starting material the reaction was concentrated *in vacuo* until almost no water remained. The residue was purified by column chromatography (gradient elution 4–10% MeOH/CHCl<sub>3</sub>) to obtain the isomeric sulfenic acids **6** (28.2 mg, 46% total yield, **6a** : **6b** = 58 : 42) and sulfinites **7** (19.1 mg, 22% total yield, **7a** : **7b** = 57 : 43). Isomerically pure **6a** was obtained using the same protocol starting from thiol **1a**. To obtain isomerically pure **7a**, sulfenic acid **6a** (5.8 mg, 0.00823 mmol) was dissolved in a 10 mM solution of NaOH in  $\text{CD}_3\text{OD}$  (0.75 mL) in an NMR tube. 25  $\mu\text{L}$  of  $\text{H}_2\text{O}_2$  as a 300 mM solution in  $\text{CD}_3\text{OD}$  was then added in 5  $\mu\text{L}$  portions until quantitative conversion of **6a** to **7a** was observed by  $^{19}\text{F}$  NMR.

**4.2.5.1 2,7-Di(tetraethylene glycol)-10-fluoro-9,10-[1',2']benzenoanthracen-9(10H)-ylsulfenic acid (6a).**  $^1\text{H}$  NMR (600 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  7.53–7.48 (m, 2H), 7.41–7.38 (d,  $J$  = 8.2 Hz, 2H), 7.23 (t,  $J$  = 2.1 Hz, 2H), 7.14–7.05 (m, 2H), 6.67 (dd,  $J$  = 8.2, 2.1 Hz, 2H), 4.08 (m, 4H), 3.77 (m, 4H), 3.67–3.63 (m, 4H), 3.63–3.57 (m, 12H), 3.56–3.52 (m, 4H), 3.50–3.46 (m, 4H).  $^{13}\text{C}$  NMR (151 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  158.4 (s), 146.2 (d,  $J$  = 20 Hz), 144.5 (d,  $J$  = 6 Hz), 142.7 (d,  $J$  = 6 Hz), 138.3 (d,  $J$  = 20 Hz), 128.8 (s), 126.6 (d,  $J$  = 6 Hz), 123.3 (d,  $J$  = 2.5 Hz), 119.8 (d,  $J$  = 6 Hz), 118.8 (d,  $J$  = 6 Hz), 112.2 (d,  $J$  = 2.5 Hz), 111.4 (s), 111.4 (s), 99.0 (d,  $J$  = 200 Hz), 73.6 (s), 71.7 (s), 71.6 (s), 71.5 (s), 71.3 (s), 71.3 (s), 70.8 (s), 68.9 (s), 64.5 (d,  $J$  = 2.5 Hz), 62.2 (s).  $^{19}\text{F}$  NMR (471 MHz,  $\text{D}_2\text{O}/\text{THF}$ )  $\delta$  –206.65. HRMS (ESI $+$ )  $m/z$  calculated for  $[\text{M} + \text{Na}]^+$   $\text{C}_{36}\text{H}_{45}\text{FNaO}_{11}\text{S}$  727.2564, found 727.2592.

**4.2.5.2 2,6-Di(tetraethylene glycol)-10-fluoro-9,10-[1',2']benzenoanthracen-9(10H)-ylsulfenic acid (6b).**  $^1\text{H}$  NMR (600 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  7.53–7.47 (m, 2H), 7.50–7.43 (m, 2H), 7.31–7.27 (m, 1H), 7.17–7.05 (m, 3H), 6.68 (dd,  $J$  = 8.2, 2.4 Hz, 1H), 6.62 (dd,  $J$  = 8.2, 2.4 Hz, 1H), 4.16 (m, 4H), 3.85 (m, 4H), 3.74–3.57 (m, 20H), 3.54 (t,  $J$  = 4.5 Hz, 4H).  $^{13}\text{C}$  NMR (151 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  158.6 (s), 158.3 (s), 147.5 (d,  $J$  = 20 Hz), 145.7 (d,  $J$  = 20 Hz), 145.0 (d,  $J$  = 6 Hz), 143.2 (d,  $J$  = 6 Hz), 137.8 (d,  $J$  = 6 Hz), 135.3 (d,  $J$  = 20 Hz),

126.8 (s), 126.5 (s), 125.4 (s), 124.5 (d,  $J$  = 2.5 Hz), 123.1 (d,  $J$  = 2.5 Hz), 120.0 (d,  $J$  = 6 Hz), 119.0 (d,  $J$  = 6 Hz), 111.9 (d,  $J$  = 2.5 Hz), 111.7 (s), 111.3 (s), 99.0 (d,  $J$  = 200 Hz), 74.5 (s), 72.6 (s), 72.4 (s), 72.2 (s), 71.7 (s), 69.8 (s), 64.1 (d,  $J$  = 2.5 Hz), 63.1 (s).  $^{19}\text{F}$  NMR (471 MHz,  $\text{D}_2\text{O}/\text{THF}$ )  $\delta$  –205.95. HRMS (ESI $+$ )  $m/z$  calculated for  $[\text{M} + \text{Na}]^+$   $\text{C}_{36}\text{H}_{45}\text{FNaO}_{11}\text{S}$  727.2564, found 727.2592.

**4.2.5.3 2,7-Di(tetraethylene glycol)-10-fluoro-9,10-[1',2']benzenoanthracen-9(10H)-ylsulfinate (7a).**  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.22 (br, 2H), 7.96 (br, 1H), 7.46 (m, 1H), 7.37 (d,  $J$  = 8.2 Hz, 2H), 7.11–7.04 (m, 2H), 6.67 (dd,  $J$  = 8.2 Hz, 2.1 Hz, 2H), 4.15 (m, 4H), 3.81 (m, 4H), 3.71–3.56 (m, 20H), 3.54–3.49 (m, 4H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  157.3 (s), 157.2 (s), 149.1 (d,  $J$  = 20 Hz), 147.7 (d,  $J$  = 6 Hz), 143.8 (d,  $J$  = 6 Hz), 142.6 (d,  $J$  = 6 Hz), 140.2 (d,  $J$  = 20 Hz), 128.8 (s), 126.0 (d,  $J$  = 20 Hz), 125.4 (d,  $J$  = 2.5 Hz), 119.7 (d,  $J$  = 6 Hz), 118.6 (d,  $J$  = 6 Hz), 113.9 (d,  $J$  = 2.5 Hz), 111.1 (s), 99.1 (d,  $J$  = 200 Hz), 72.3 (d,  $J$  = 2 Hz), 72.1 (s), 70.3 (s), 70.2 (s), 70.1 (s), 70.0 (s), 69.3 (s), 67.5 (s), 60.6 (s).  $^{19}\text{F}$  NMR (471 MHz,  $\text{D}_2\text{O}/\text{THF}$ )  $\delta$  –206.1. HRMS (ESI $-$ )  $m/z$  calculated for  $[\text{M} - \text{H}]^-$   $\text{C}_{36}\text{H}_{44}\text{FO}_{12}\text{S}$  719.2538, found 719.2526.

**4.2.5.4 2,6-Di(tetraethylene glycol)-10-fluoro-9,10-[1',2']benzenoanthracen-9(10H)-ylsulfinate (7b).**  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.50 (br, 1H), 7.97 (br, 1H), 7.47 (m, 1H), 7.36 (m, 1H), 7.14–7.04 (m, 3H), 6.71–6.62 (m, 2H), 4.15 (m, 4H), 3.81 (m, 4H), 3.71–3.56 (m, 20H), 3.54–3.49 (m, 4H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  157.5 (s), 157.4 (s), 145.1 (d,  $J$  = 20 Hz), 144.6 (d,  $J$  = 20 Hz), 142.0 (d,  $J$  = 6 Hz), 141.6 (d,  $J$  = 6 Hz), 137.1 (d,  $J$  = 20 Hz), 126.1 (s), 125.1 (s), 125.0 (d,  $J$  = 2.5 Hz), 119.4 (d,  $J$  = 6 Hz), 118.0 (d,  $J$  = 6 Hz), 117.0 (d,  $J$  = 6 Hz), 116.8 (d,  $J$  = 6 Hz), 114.3 (d,  $J$  = 2.5 Hz), 113.1 (d,  $J$  = 2.5 Hz), 110.5 (s), 110.3 (s), 99.9 (d,  $J$  = 200 Hz), 72.4 (d,  $J$  = 2.5 Hz), 72.1 (s), 70.3 (s), 70.2 (s), 70.1 (s), 70.0 (s), 69.3 (s), 67.5 (s), 60.6 (s).  $^{19}\text{F}$  NMR (471 MHz,  $\text{D}_2\text{O}/\text{THF}$ )  $\delta$  –205.4. HRMS (ESI $-$ )  $m/z$  calculated for  $[\text{M} - \text{H}]^-$   $\text{C}_{36}\text{H}_{44}\text{FO}_{12}\text{S}$  719.2538, found 719.2526.

**4.2.6 Potassium di(tetraethylene glycol)-10-fluoro-9,10-[1',2']benzenoanthracen-9(10H)-ylsulfonate (8).** Thiols **1** (30 mg, 0.044 mmol, as a 53 : 47 isomeric mixture of **1a** : **1b**) or purified **1a** were dissolved in THF (1 mL) and added to a solution of dibasic potassium phosphate (15 mg, 0.086 mmol) in HPLC-grade  $\text{H}_2\text{O}$  (900  $\mu\text{L}$ ). 114  $\mu\text{L}$  of  $\text{H}_2\text{O}_2$  as a 1.94 M solution in  $\text{D}_2\text{O}/\text{H}_2\text{O}$  was added all at once. After stirring overnight, the reaction was concentrated *in vacuo* until almost no water remained. The residue was purified by column chromatography (10% MeOH/CHCl<sub>3</sub>) to obtain the isomeric sulfonates **8** (30 mg, quantitative yield, **8a** : **8b** = 53 : 47).

**4.2.6.1 2,7-Di(tetraethylene glycol)-10-fluoro-9,10-[1',2']benzenoanthracen-9(10H)-ylsulfonate (8a).**  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.23 (m, 1H), 7.91 (t,  $J$  = 2.1 Hz, 2H), 7.51–7.45 (m, 1H), 7.41–7.35 (m, 2H), 7.14–7.04 (m, 2H), 6.71–6.63 (m, 2H), 4.15 (m, 4H), 3.81 (m, 4H), 3.71–3.56 (m, 20H), 3.54–3.49 (m, 4H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  156.5 (s), 145.1 (d,  $J$  = 20 Hz), 141.5 (d,  $J$  = 6 Hz), 140.0 (d,  $J$  = 6 Hz), 137.5 (d,  $J$  = 20 Hz), 125.1 (s), 124.9 (d,  $J$  = 2 Hz), 124.8 (s), 117.9 (d,  $J$  = 6 Hz), 116.9 (d,  $J$  = 6 Hz), 113.0 (d,  $J$  = 2 Hz), 110.4 (s), 97.5 (d,  $J$  = 200 Hz), 73.6 (d,  $J$  = 2 Hz), 73.4 (s), 71.6 (s), 71.5 (s), 71.4 (s), 71.3 (s), 71.2 (s), 68.9 (s), 62.2 (s), 62.0 (s).  $^{19}\text{F}$  NMR (471 MHz,  $\text{D}_2\text{O}/\text{THF}$ )  $\delta$  –205.9. HRMS (ESI $-$ )  $m/z$  calculated for  $[\text{M} - \text{H}]^-$   $\text{C}_{36}\text{H}_{44}\text{FO}_{13}\text{S}$  735.2503, found 735.2491.



**4.2.6.2 2,6-Di(tetraethylene glycol)-10-fluoro-9,10-[1',2']benzenoanthracen-9(10H)-ylsulfonate (8b).**  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  8.23 (m, 1H), 8.12 (dd,  $J$  = 8.4, 1.4 Hz, 1H), 7.89 (t,  $J$  = 2.1 Hz, 1H), 7.51–7.45 (m, 1H), 7.41–7.35 (m, 1H), 7.14–7.04 (m, 3H), 6.71–6.63 (m, 2H), 4.15 (m, 4H), 3.81 (m, 4H), 3.71–3.56 (m, 20H), 3.54–3.49 (m, 4H).  $^{13}\text{C}$  NMR (126 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  156.7 (s), 156.6 (s), 146.4 (d,  $J$  = 20 Hz), 144.5 (d,  $J$  = 20 Hz), 142.0 (d,  $J$  = 6 Hz), 140.0 (d,  $J$  = 6 Hz), 137.0 (d,  $J$  = 20 Hz), 132.5 (d,  $J$  = 6 Hz), 126.2 (d,  $J$  = 2 Hz), 125.0 (s), 124.9 (d,  $J$  = 2 Hz), 124.7 (d,  $J$  = 2 Hz), 118.0 (d,  $J$  = 6 Hz), 117.0 (d,  $J$  = 6 Hz), 112.7 (d,  $J$  = 2 Hz), 110.4 (s), 110.0 (s), 97.5 (d,  $J$  = 200 Hz), 73.6 (d,  $J$  = 2 Hz), 73.4 (s), 71.6 (s), 71.5 (s), 71.4 (s), 71.3 (s), 71.2 (s), 68.9 (s), 62.2 (s), 62.0 (s).  $^{19}\text{F}$  NMR (471 MHz,  $\text{D}_2\text{O}/\text{THF}$ )  $\delta$  –205.25. HRMS (ESI $+$ )  $m/z$  calculated for  $[\text{M} + \text{Na}]^+$   $\text{C}_{36}\text{H}_{44}\text{FO}_{13}\text{S}$  735.2503, found 735.2487.

**4.2.7 2,7-Di(tetraethylene glycol)-10-fluoro-9,10-[1',2']benzenoanthracen-9(10H)-yl disulfide (13a).** In an NMR tube, thiol **1a** (8.2 mg, 11.9  $\mu\text{mol}$ ) was dissolved in THF (175  $\mu\text{L}$ ) containing 240 mM of TFE (internal standard). Phosphate-buffered saline (490  $\mu\text{L}$  of a 50 mM pD 7.8 solution in  $\text{D}_2\text{O}$ ) was then added. Upon reaching homogeneity, 17  $\mu\text{L}$  of NaOCl (as a 700 mM solution in  $\text{D}_2\text{O}/\text{H}_2\text{O}$ ) was added portion-wise, monitoring the consumption of **1a**. The tube was mixed, and the reaction was complete upon acquiring the next set of scans (*ca.* 45 s). The contents of the NMR tube were diluted with distilled  $\text{H}_2\text{O}$  (10 mL) and extracted with  $\text{CHCl}_3$  ( $3 \times 5$  mL), dried over  $\text{MgSO}_4$ , and concentrated *in vacuo* to give **13a** (8.2 mg, 11.9  $\mu\text{mol}$ ).  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  8.11 (d,  $J$  = 7.5 Hz, 2H), 7.55 (d, 2H), 7.39 (d,  $J$  = 8.2 Hz, 4H), 7.33 (t,  $J$  = 2.1 Hz, 4H), 7.13 (td,  $J$  = 7.5 Hz, 1.3 Hz, 2H), 7.04 (td,  $J$  = 7.5 Hz, 1.3 Hz, 2H), 6.56 (d,  $J$  = 8.2 Hz, 2.1 Hz, 4H), 4.15 (m, 8H), 3.81 (m, 8H), 3.71–3.56 (m, 40H), 3.54–3.49 (m, 4H), 2.78 (br s, exchangeable Hs).  $^{13}\text{C}$  NMR (151 MHz,  $\text{CDCl}_3$ )  $\delta$  156.4 (s), 144.7 (d,  $J$  = 20 Hz), 142.8 (d,  $J$  = 6 Hz), 141.5 (s), 136.3 (d,  $J$  = 20 Hz), 126.1 (s), 125.3 (s), 123.1 (s), 119.0 (d,  $J$  = 2.5 Hz), 118.1 (d,  $J$  = 2.5 Hz), 118.1 (d,  $J$  = 2.5 Hz), 111.7 (s), 111.4 (s), 97.2 (d,  $J$  = 200 Hz), 72.5 (s), 70.6 (s), 70.5 (s), 70.3 (s), 69.5 (s), 67.0 (s), 62.8 (d,  $J$  = 2.5 Hz), 61.7 (s).  $^{19}\text{F}$  NMR (471 MHz,  $\text{D}_2\text{O}/\text{THF}$ )  $\delta$  –205.9. HRMS (ESI $+$ )  $m/z$  calculated for  $[\text{M} + \text{Na}]^+$   $\text{C}_{72}\text{H}_{88}\text{F}_2\text{O}_{20}\text{NaS}_2$  1397.5176, found 1397.5193.

**4.2.8 2,7-Di(tetraethylene glycol)-10-fluoro-9,10-[1',2']benzenoanthracen-9(10H)-ylsulfenyl chloride (14a).** In an NMR tube, thiol **1a** (6.9 mg, 10.0  $\mu\text{mol}$ ) was dissolved in  $\text{CDCl}_3$  (700  $\mu\text{L}$ ) containing TFE (internal standard). *N*-Chlorosuccinimide as a 200 mM solution in  $\text{CDCl}_3$  was added portion-wise every 5 minutes until the starting thiol was completely consumed (*ca.* 50  $\mu\text{L}$ , 1 eq.). The contents of the NMR tube were washed with distilled  $\text{H}_2\text{O}$  (5 mL) and extracted with  $\text{CHCl}_3$  ( $3 \times 3$  mL), dried over  $\text{MgSO}_4$ , and concentrated *in vacuo* to give **14a** (6.8 mg, 94% yield).  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  7.56 (dd,  $J$  = 7.4 Hz, 1.0 Hz, 2H), 7.45 (d,  $J$  = 8.2 Hz, 2H), 7.39 (d,  $J$  = 8.2 Hz, 2H), 7.23 (br m, 2H), 7.17–7.07 (m, 2H), 6.65 (d,  $J$  = 8.2 Hz, 2.1 Hz, 2H), 4.09 (m, 4H), 3.81 (m, 4H), 3.72–3.61 (m, 20H), 3.58–3.54 (m, 4H), 2.61 (br s, exchangeable H's).  $^{13}\text{C}$  NMR (151 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  156.9 (s), 144.4 (d,  $J$  = 20 Hz), 141.3 (d,  $J$  = 6 Hz), 139.8 (d,  $J$  = 6 Hz), 136.6 (d,  $J$  = 20 Hz), 126.2 (s), 125.6 (s), 122.1 (br), 119.4 (d,  $J$  = 6 Hz), 118.4 (d,  $J$  = 6 Hz), 110.8 (s), 97.4 (d,  $J$  = 200 Hz), 72.5 (s), 70.8 (s), 70.7 (s), 70.6 (s), 70.3 (s), 69.6 (s), 67.8 (s), 63.8 (d,  $J$  = 2 Hz), 61.7

(s).  $^{19}\text{F}$  NMR (471 MHz,  $\text{D}_2\text{O}/\text{THF}$ )  $\delta$  –206.4. HRMS (ESI $+$ )  $m/z$  calculated for  $[\text{M} + \text{Na}]^+$   $\text{C}_{36}\text{H}_{44}\text{FCIO}_{10}\text{S}$  745.2225, found 745.2249.

**4.2.9 10-Fluoro-9,10-[1',2']benzenoanthracen-9(10H)-yl disulfide (15).** Thiol **9** (ref. 17) (30 mg, 0.1 mmol) was suspended in MeOH (5 mL), potassium carbonate (45 mg, 0.3 mmol) was added as a solid, and the mixture was stirred for 10 minutes at room temperature. Iodine (50 mg, 0.2 mmol) was then added directly and the reaction was stirred for 2 hours. The reaction was then poured into an aqueous solution (50 mL) of sodium thiosulfate pentahydrate (100 mg, 0.4 mmol), and subsequently extracted with EtOAc ( $3 \times 25$  mL). The organic layer was dried over  $\text{MgSO}_4$  and concentrated *in vacuo*. The residue was purified by column chromatography (10% EtOAc/hexanes) to obtain disulfide **15** (11 mg, 37%). Although conversion looked to be complete by TLC, the isolated yield was reduced due to low solubility of the compound and corresponding losses up during chromatography.  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  7.99 (dd,  $J$  = 7.6, 1.1 Hz, 6H), 7.61 (dd,  $J$  = 7.6, 1.1 Hz, 6H), 7.11 (td,  $J$  = 7.6, 1.1 Hz, 6H), 6.94 (td,  $J$  = 7.6, 1.1 Hz, 6H).  $^{13}\text{C}$  NMR (151 MHz,  $\text{CDCl}_3$ )  $\delta$  146.1, 143.5, 143.3, 141.5, 126.0, 125.5, 123.3, 118.0, 96.6, 62.8.  $^{19}\text{F}$  NMR (282 MHz,  $\text{C}_6\text{D}_6$ )  $\delta$  –206.6. HRMS (EI)  $m/z$  calculated for  $[\text{M}]^+$   $\text{C}_{40}\text{H}_{24}\text{F}_2\text{S}_2$  606.1288, found 606.1287.

### 4.3 $^{19}\text{F}$ NMR-monitored reactions

**4.3.1 Buffer preparation.** Mixtures of  $\text{K}_2\text{HPO}_4$  and  $\text{KH}_2\text{PO}_4$  were dissolved in  $\text{D}_2\text{O}$  to a final concentration of 50 mM. Either KOH or HCl was added to adjust the pH to the desired value. Observed pH (pD) was converted to pH by subtracting 0.4.<sup>19</sup>

**4.3.2 Oxidation of **1** with  $\text{H}_2\text{O}_2$ .** Solutions of **1** (typically 15–20 mM) in THF containing 240 mM TFE were prepared before each experiment. 175  $\mu\text{L}$  of this solution was transferred to a NMR tube. 525  $\mu\text{L}$  of phosphate-buffered  $\text{D}_2\text{O}$  was added to the NMR tube and the mixture was vortexed to homogeneity. An initial set of scans was acquired and then  $\text{H}_2\text{O}_2$  was added as a 1.94 M solution in  $\text{D}_2\text{O}/\text{H}_2\text{O}$ . The NMR tube was mixed before the reaction was monitored for 15 hours. COPASI<sup>44</sup> was used to fit the data to a simple kinetic scheme for successive bimolecular reactions of thiol, sulfenic and sulfinic acids with  $\text{H}_2\text{O}_2$ , which yielded the observed rate constants given in the text.

**4.3.3 Oxidation of **1a** with HOCl.** A solution of **1a** (68 mM) in THF containing 240 mM TFE was prepared. An aliquot (175  $\mu\text{L}$ ) of this solution was transferred to an NMR tube and 490  $\mu\text{L}$  of phosphate-buffered  $\text{D}_2\text{O}$  (pD 7.8) was added. An initial spectrum was recorded, after which 0.35 eq. of NaOCl (as a solution in  $\text{D}_2\text{O}$ ) was added at once. The tube was mixed and the spectrum was recorded, revealing immediate quantitative conversion to the disulfide **13a**.

**4.3.4 Reduction of sulfenyl chloride **14a** with **1a**.** A solution of **14a** (50 mM) in THF containing 240 mM TFE was prepared. An aliquot (175  $\mu\text{L}$ ) of this solution was transferred to an NMR tube and 490  $\mu\text{L}$  of phosphate-buffered  $\text{D}_2\text{O}$  (pD 7.8) was added. The NMR tube was left overnight under ambient conditions to rule out hydrolysis as a decomposition pathway. After 18 h an initial set of spectra was acquired and a solution of **1a** (20 mM)



in  $\text{D}_2\text{O}/\text{THF}$  (3 : 1) was added dropwise, while mixing, revealing stoichiometric conversion of **14a** to **13a**.

**4.3.5 Oxidation of **1** with thermally generated  $^1\text{O}_2$ .** A 20 mM solution of **1** in THF containing 240 mM TFE was prepared before the experiment. 175  $\mu\text{L}$  of this solution was transferred to an NMR tube and 525  $\mu\text{L}$  of an ice-cooled freshly prepared 27 mM solution of NDPO<sub>2</sub> (prepared using a reported procedure<sup>45</sup>) in phosphate-buffered  $\text{D}_2\text{O}$  (pD 7.8) was added to the NMR tube. The NMR tube was mixed and an initial set of scans at 25 °C was acquired before the NMR spectrometer was heated to 37 °C using a thermostat. The reaction was then monitored for 30 min. Due to significant deviation in chemical shift values from the conditions involving oxidation of **1** by  $\text{H}_2\text{O}_2$ , a reaction mixture obtained by reacting thiol **1** with  $\text{H}_2\text{O}_2$  was added to the reaction mixture after completion to identify the observed products.

**4.3.6 Oxidation of **9** with photochemically generated  $^1\text{O}_2$ .** In an NMR tube, **9** (6 mg, 0.02 mmol) was suspended in  $\text{CD}_3\text{OD}$  (690  $\mu\text{L}$ ), to this suspension was added sodium hydroxide as a 1 M solution in water (60  $\mu\text{L}$ , 0.06 mmol, 3 eq.) and the mixture became homogeneous. 10  $\mu\text{L}$  of a 1.5 mM aqueous solution of Rose Bengal (0.0002 mmol, 0.01 eq.) was added and the reaction mixture was cooled to *ca.* 4 °C. Trifluorotoluene (10 mM) was used as an internal standard for NMR quantification. Oxygen was bubbled through the solution after which the reaction vessel was irradiated with a 400 W HPS lamp. Time points were taken to analyze reaction progress. The reaction was continued to the point that **9** was fully consumed, after which the reaction was concentrated *in vacuo*. Following this, the crude product was concentrated *in vacuo*, suspended in  $\text{C}_6\text{D}_6$ , and analyzed by <sup>19</sup>F NMR, which resulted in the observation of disulfide **14** as a trace reaction product, **11** and **12** were not observed despite being major products as they are insoluble in  $\text{C}_6\text{D}_6$  (Fig. S5†). Products were assigned after adding authentic **12** to the NMR tube ( $\text{CD}_3\text{OD}$ ). Monitoring of reaction progress is shown in Fig. 3B of the manuscript. <sup>1</sup>H NMR spectra of the initial and final (after 6 min. of irradiation) time points are shown in Fig. S4.†

**4.3.7 Reduction of sulfenic acid **6** with ascorbate.** A 5 mM solution of **6** in THF containing 10 mM TFE was prepared before the experiment. 175  $\mu\text{L}$  of this solution was transferred to an NMR tube and 300  $\mu\text{L}$  of phosphate-buffered pD 7.8  $\text{D}_2\text{O}$  was added to the NMR tube before an initial set of scans was acquired. No spectroscopic changes were observed after 24 h. An additional time point was taken after 3 days, and no changes were observed.

**4.3.8 Reduction of sulfenic acid **10** with ascorbate and iron.** Sulfenic acid **10** (6.4 mg, 0.02 mmol) was dissolved in MeOH (8 mL). 1 mL of a freshly prepared 200 mM solution of ascorbic acid in MeOH (1 eq.) was then added, followed by addition of a 20 mM solution of  $\text{Fe}_2(\text{SO}_4)_3$  in MeOH (1 mL, 0.2 eq.). The mixture was stirred for 24 hours, after which it was diluted with  $\text{CH}_2\text{Cl}_2$  (50 mL), filtered through a silica plug and washed with 20% MeOH/ $\text{CH}_2\text{Cl}_2$  (50 mL). The filtrate was concentrated *in vacuo* and analyzed by <sup>19</sup>F NMR. No changes were observed, with the mass balance indicating that no reaction had occurred.

## Data availability

Additional experimental procedures and reaction monitoring data, NMR spectra and computational data are available in the ESI.†

## Author contributions

D. D. S. designed and executed the experiments and carried out quantum chemical calculations. D. A. P. conceived the project and contributed to experimental design. D. D. S. and D. A. P. wrote the manuscript.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

This work was supported by grants from the Natural Sciences and Engineering Council of Canada and the Canada Foundation for Innovation to D. A. P. Computational investigations were enabled by Compute Ontario and the Digital Research Alliance of Canada. D. D. S. thanks the Government of Ontario for support in the form of an Ontario Graduate Scholarship.

## Notes and references

‡ Kinetic studies on reactions of protein sulfenic acids have been conducted, however, the approach did not involve direct visualization of the RSOH intermediate. See ref. 39.

§ It was assumed that the sulfinic acid is fully deprotonated at this pD, based on previously measured  $\text{pK}_a$  values.<sup>17</sup>

¶ Reduction potentials in water ( $E^0(\text{RS}^\bullet/\text{RS}^-) = +0.84$  V vs. NHE<sup>46</sup> and  $E^0(\text{O}_2/\text{O}_2^\bullet^-) = +0.81$  vs. NHE<sup>47</sup>) suggest that the reaction of  $^1\text{O}_2$  and  $\text{RS}^-$  is essentially thermoneutral. This may indicate an even stronger oxidant than  $^1\text{O}_2$  is formed in the reaction. Earlier studies support this notion, as it has been shown that thiols significantly exacerbate the damage to DNA caused by  $^1\text{O}_2$  and that hydroxyl radicals may be formed in the reaction.<sup>48</sup>

|| Three equivalents of ascorbate were selected for this reaction based on the reported rate constants.<sup>49</sup> If reduction were to occur, it would be possible to monitor by <sup>19</sup>F NMR under these conditions.

- 1 B. Morgan, D. Ezeriña, T. N. E. Amoako, J. Riemer, M. Seedorf and T. P. Dick, *Nat. Chem. Biol.*, 2013, **9**, 119–125.
- 2 G. J. McBean, M. Aslan, H. R. Griffiths and R. C. Torrão, *Redox Biol.*, 2015, **5**, 186–194.
- 3 J.-M. Lv, S.-Q. Lü, Z.-P. Liu, J. Zhang, B.-X. Gao, Z.-Y. Yao, Y.-X. Wu, L. A. Potempa, S.-R. Ji, M. Long and Y. Wu, *Sci. Rep.*, 2018, **8**, 1494.
- 4 S. García-Santamarina, S. Boronat and E. Hidalgo, *Biochemistry*, 2014, **53**, 2560–2580.
- 5 S. G. Rhee, *Science*, 2006, **312**, 1882–1883.
- 6 Z. Wu, U. Barayeu, D. Schilling, T. P. Dick and D. A. Pratt, *Curr. Opin. Chem. Biol.*, 2023, **76**, 102353.
- 7 Y.-C. Chang, C.-N. Huang, C.-H. Lin, H.-C. Chang and C.-C. Wu, *Proteomics*, 2010, **10**, 2961–2971.



8 S. Akter, L. Fu, Y. Jung, M. Lo Conte, J. R. Lawson, W. T. Lowther, R. Sun, K. Liu, J. Yang and K. S. Carroll, *Nat. Chem. Biol.*, 2018, **14**, 995–1004.

9 J. M. M. Pople and J. M. Chalker, *Curr. Opin. Chem. Biol.*, 2021, **60**, 55–65.

10 N. J. Kettenhofen and M. J. Wood, *Chem. Res. Toxicol.*, 2010, **23**, 1633–1646.

11 V. Gupta and K. S. Carroll, *Biochim. Biophys. Acta, Gen. Subj.*, 2014, **1840**, 847–875.

12 C. C. Winterbourn and D. Metodiewa, *Free Radical Biol. Med.*, 1999, **27**, 322–328.

13 N. Nakamura, *J. Am. Chem. Soc.*, 1983, **105**, 7172–7173.

14 T. Yoshimura, E. Tsukurimichi, S. Yamazaki, S. Soga, C. Shimasaki and K. Hasegawa, *J. Chem. Soc. Chem. Commun.*, 1992, 1337–1338.

15 T. Sano, R. Masuda, S. Sase and K. Goto, *Chem. Commun.*, 2021, **57**, 2479–2482.

16 C. C. Winterbourn and M. B. Hampton, *Free Radical Biol. Med.*, 2008, **45**, 549–561.

17 J.-P. R. Chauvin and D. A. Pratt, *Angew. Chem., Int. Ed.*, 2017, **56**, 6255–6259.

18 Y. Nishii, M. Ikeda, Y. Hayashi, S. Kawauchi and M. Miura, *J. Am. Chem. Soc.*, 2020, **142**, 1621–1629.

19 A. Kręzel and W. Bal, *J. Inorg. Biochem.*, 2004, **98**, 161–166.

20 I. Canals, F. Z. Oumada, M. Rosés and E. Bosch, *J. Chromatogr. A*, 2001, **911**, 191–202.

21 G. Klopman, K. Tsuda, J. B. Louis and R. E. Davis, *Tetrahedron*, 1970, **26**, 4549–4554.

22 C. Storkey, M. J. Davies and D. I. Pattison, *Free Radical Biol. Med.*, 2014, **73**, 60–66.

23 X. L. Armesto, M. Canle L, M. I. Fernández, M. V García and J. A. Santaballa, *Tetrahedron*, 2000, **56**, 1103–1109.

24 P. Nagy and M. T. Ashby, *J. Am. Chem. Soc.*, 2007, **129**, 14082–14091.

25 N. T. Ioffe, M. I. Kalinkin, I. N. Rozhkov and I. L. Knunyants, *Bull. Acad. Sci. USSR, Div. Chem. Sci.*, 1969, **18**, 1929–1930.

26 M. J. Davies and C. L. Hawkins, *Free Radical Res.*, 2000, **33**, 719–729.

27 T. P. A. Devasagayam, A. R. Sundquist, P. Di Mascio, S. Kaiser and H. Sies, *J. Photochem. Photobiol., B*, 1991, **9**, 105–116.

28 A. W. M. Nieuwint, J. M. Aubry, F. Arwert, H. Kortbeek, S. Herzberg and H. Joenje, *Free Radical Res. Commun.*, 1985, **1**, 1–9.

29 P. Di Mascio and H. Sies, *J. Am. Chem. Soc.*, 1989, **111**, 2909–2914.

30 G. R. Buettner, *FEBS Lett.*, 1984, **177**, 295–299.

31 A. J. McGrath, G. E. Garrett, L. Valgimigli and D. A. Pratt, *J. Am. Chem. Soc.*, 2010, **132**, 16759–16761.

32 T. P. A. Devasagayam, A. R. Sundquist, P. Di Mascio, S. Kaiser and H. Sies, *J. Photochem. Photobiol., B*, 1991, **9**, 105–116.

33 J. A. Montgomery Jr, M. J. Frisch, J. W. Ochterski and G. A. Petersson, *J. Chem. Phys.*, 2000, **112**, 6532–6542.

34 C. Chatgilialoglu and M. Guerra, in *Sulfur-Centered Reactive Intermediates in Chemistry and Biology*, ed. C. Chatgilialoglu and K.-D. Asmus, Springer US, Boston, MA, 1990, pp. 31–36.

35 J. Mönig, K.-D. Asmus, L. G. Forni and R. L. Willson, *Int. J. Radiat. Biol. Relat. Stud. Phys., Chem. Med.*, 1987, **52**, 589–602.

36 A. V Kachur, C. J. Koch and J. E. Biaglow, *Free Radical Res.*, 1998, **28**, 259–269.

37 H. Berger, *Recl. Trav. Chim. Pays-Bas*, 1963, **82**, 773–789.

38 G. Monteiro, B. B. Horta, D. C. Pimenta, O. Augusto and L. E. S. Netto, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 4886–4891.

39 L. Turell, H. Botti, S. Carballal, G. Ferrer-Sueta, J. M. Souza, R. Durán, B. A. Freeman, R. Radi and B. Alvarez, *Biochemistry*, 2008, **47**, 358–367.

40 M. Conrad and D. A. Pratt, *Nat. Chem. Biol.*, 2019, **15**, 1137–1147.

41 D. D. Saraev, Z. Wu, H.-Y. H. Kim, N. A. Porter and D. A. Pratt, *ACS Chem. Biol.*, 2023, **18**, 2073–2081.

42 W. M. Haynes, *CRC Handbook of Chemistry and Physics*, Taylor & Francis, 93rd edn, 2012.

43 J. N. Cvengroš Stefan, B. Anne and S. Hans-Günther, *Synlett*, 2008, **2008**, 1993–1998.

44 S. Hoops, S. Sahle, R. Gauges, C. Lee, J. Pahle, N. Simus, M. Singhal, L. Xu, P. Mendes and U. Kummer, *Bioinformatics*, 2006, **22**, 3067–3074.

45 J. M. Aubry, *J. Am. Chem. Soc.*, 1985, **107**, 5844–5849.

46 P. S. Surdhar and D. A. Armstrong, *J. Phys. Chem.*, 1987, **91**, 6532–6537.

47 W. H. Koppenol, D. M. Stanbury and P. L. Bounds, *Free Radical Biol. Med.*, 2010, **49**, 317–322.

48 T. P. A. Devasagayam, P. Di Mascio, S. Kaiser and H. Sies, *Biochim. Biophys. Acta, Gene Struct. Expression*, 1991, **1088**, 409–412.

49 V. Anschau, G. Ferrer-Sueta, R. L. Aleixo-Silva, R. Bannitz Fernandes, C. A. Tairum, C. C. C. Tonoli, M. T. Murakami, M. A. de Oliveira and L. E. S. Netto, *Free Radical Biol. Med.*, 2020, **156**, 207–216.

